Copper complex formed with neurokinin B: binding stoichiometry, redox property, self-assembly and cytotoxicity

Yaliang Huang^{a,b}, Ning Xia^b, Miao Chen^a, Ping Yang^b, Lin Liu^{b,*}, Jianxiu Wang^{a,*}

^aHunan Provincial Key Laboratory of Micro & Nano Materials Interface Science, College of Chemistry and Chemical Engineering, Central South University, Changsha, Hunan 410083, People's Republic of China

^bHenan Province of Key Laboratory of New Optoelectronic Functional Materials, College of Chemistry and Chemical Engineering, Anyang Normal University, Anyang, Henan 455000, People's Republic of China



Fig. S1 Mass spectra of NKB (A and C) and NKB-Cu²⁺ complex (B and D) at pH 6.0 (A and B) and 8.0 (C and D).



Fig. S2 AFM image of *D*-type of NKB peptide after incubation for 24 h.



Fig. S3 (A) ThT assay of NKB, neurokinin A and substance P. (B) AFM images of neurokinin A and substance P after incubation for 24 h.



Fig. S4 AFM images for the fibrous <u>A</u>MHDFFVGLM-NH₂ (a) and DMH<u>A</u>FFVGLM-NH₂ (b) after incubation with Cu^{2+} at 37 °C for 1 h.