

Supporting Information

(New Journal of Chemistry)

Synthesis of one dimensional Cu₂S nanorods using self-grown sacrificial template for electrocatalytic oxygen evolution reaction (OER)

S. B. Kale^{1,2}, P. T. Babar², J. H. Kim^{2*}, C. D. Lokhande^{1*}

¹Centre for Interdisciplinary Research, D. Y. Patil Education Society (Institution Deemed to Be University), Kolhapur, 416 004, MS, India.

²Optoelectronic Convergence Research Center, Department of Materials Science and Engineering, Chonnam National University, Gwangju 500-757, South Korea

Figure S1: XRD patterns of Cu₂S/Cu samples prepared for different sulfurization time.

Figure S2: XRD patterns of Cu₂S/Cu samples prepared for different Na₂S concentration.

Figure S3: The FE-SEM images of Cu₂S/Cu samples prepared for different sulfurization time at 25kX magnification.

Figure S4: The FE-SEM images of Cu₂S/Cu samples prepared for different Na₂S concentration at 25kX magnification.

Figure S5: XRD pattern of Cu₂S/Cu after stability test.

Figure S6: LSV curves of Cu₂S/Cu samples prepared for different sulfurization time.

Figure S7: LSV curves of Cu₂S/Cu samples prepared for different Na₂S concentration.

Figure S8: Cyclic voltammetry curves in non-faradic region for (a) $\text{Cu}(\text{OH})_2/\text{Cu}$ and (b) $\text{Cu}_2\text{S}/\text{Cu}$




Fig. S1




Fig. S2




Fig. S3




Fig. S4




Fig. S5




Fig. S6




Fig. S7




Fig. S8