

Electronic Supplementary Information (ESI) for New Journal of Chemistry

**Delineating the role of crystallinity on the electrocatalytic activity of
colloidally synthesized MoP nanocrystals**

Siyabonga S. Nkabinde, Zakhele B. Ndala, Ndivhuwo P. Shumbula, Tshwarela Kolokoto,
Obakeng Nchoe, Grace N. Ngubeni, Kalenga P. Mubiayi, Nosipho Moloto*

*Molecular Sciences Institute, School of Chemistry, University of the Witwatersrand, Private
Bag 3, Wits, 2050, South Africa*

*Corresponding authors: Nosipho Moloto

Email: Nosipho.Moloto@wits.ac.za

Tel: +2711 7176774

Fax: +2711 7176749

Supporting information




Fig. S1: The size distributions of (a) A-MoP and (b) C-MoP nanocrystals as determined from TEM images ($N = 100$).




Fig. S2: EDX spectra of (a) amorphous A-MoP and (b) crystalline C-MoP(2 h) nanocrystals.




Fig. S3: XRD pattern of long term stored amorphous A-MoP.




Fig. S4: XRD pattern of long term stored amorphous C-MoP.




Fig. S5: High-resolution spectrum of C 1s of A-MoP.




Fig. S6: High-resolution spectrum of C 1s of C-MoP.