Supporting Information

Near-Infrared Triggered Release of Tirofiban from Nanocarriers for Inhibition of Platelet Integrin αIIbβ3 to Decrease Early Stage Neointima Formation

Zhen Zhao, ^{‡a} Peng Qiu, ^{‡a} Huaxiang Lu, ^{‡a} Minyi Yin, ^{*a} Xiaobing Liu, ^{*a} Fengshi Li, ^a

Kai Liu,^{a,b} Dalin Li,^b Xinwu Lu,*a and Bo Li*a

University Medical College, Qingdao 266000, China

^aDepartment of Vascular Surgery, Shanghai Ninth People's Hospital, Shanghai Jiao Tong University School of Medicine, Shanghai 200011, China E-mail: <u>yinminyi9@126.com; benny_liuxb@163.com; luxinwu@shsmu.edu.cn;</u> <u>boli@shsmu.edu.cn</u> ^bDepartment of Vascular Surgery, Qingdao Municipal Hospital Affiliated to Qingdao



Figure S1. TEM image of CuS nanoparticles.



Figure S2. The size (A) and zeta potential (B) of CuS, CuS/CTAB and CuS@mSiO₂-PEG nanoparticles.



Figure S3. Dynamic light scattering (DLS) data of as-prepared (blue line) and stored (in PBS solution for 5 days) CuS@mSiO₂-PEG core-shell nanoparticles.



Figure S4. H&E stained slices of main organs. Magnification: 200 times.