Novel synthetic strategy towards subphthalocyanine-functionalized acetylenic scaffolds via various dibromo-enynes

Line Broløs, Martin Drøhse Kilde and Mogens Brøndsted Nielsen*

Department of Chemistry, University of Copenhagen, Universitetsparken 5, DK-2100 Copenhagen Ø, Denmark.

E-mail: mbn@chem.ku.dk

Electronic Supplementary Information

Table of Contents

NMR and mass spectra	S2
·	
UV-Vis absorption spectra	S26

NMR and mass spectra



Figure S2: ¹³C-APT NMR (CDCl₃, 126 MHz) spectrum of 1a.



Figure S3: High-resolution mass spectrum of 1a.



Figure S4: ¹H-NMR (CDCl₃, 500 MHz) spectrum of 2a.



Figure S5: (left) COSY (500 MHz) and (right) ¹H/¹³C-APT HSQC (500/126 MHz) spectra of **2a** recorded in CDCl₃.



Figure S6: ¹³C-APT NMR (CDCl₃, 126 MHz) spectrum of 2a.



Figure S7: High-resolution mass spectrum of 2a.



Figure S8: ¹H-NMR (CDCl₃, 500 MHz) spectrum of **3a**.



Figure S9: (left) COSY (500 MHz) and (right) ¹H/¹³C HSQC (500/126 MHz) spectra of **3a** recorded in CDCl₃.



Figure S10: ¹³C-NMR (CDCl₃, 126 MHz) spectrum of 3a.



Figure S11: High-resolution mass spectrum of 3a.



Figure S12: ¹H-NMR (CDCl₃, 500 MHz) spectrum of 4a.



Figure S13: ¹³C-NMR (CDCl₃, 126 MHz) spectrum of 4a.



Figure S14: High-resolution mass spectrum of 4a.





Figure S15: ¹H-NMR (CDCl₃, 500 MHz) spectrum of 1b.



Figure S16: ¹³C-NMR (CDCl₃, 126 MHz) spectrum of 1b.



Figure S17: High-resolution mass spectrum of 1b.



Figure S18: ¹H-NMR (CDCl₃, 500 MHz) spectrum of 2b.



Figure S19: (left) COSY (500 MHz) and (right) ¹H/¹³C HSQC (500/126 MHz) spectra of 2b recorded in CDCl₃.



Figure 20: ¹³C-NMR (CDCl₃, 126 MHz) spectrum of **2b**.



Figure S21: High-resolution mass spectrum of 2b.



Figure S22: ¹H-NMR (CDCl₃, 500 MHz) spectrum of **3b**.



Figure S23: ¹³C-NMR (CDCl₃, 126 MHz) spectrum of **3b**.



Figure S24: High-resolution mass spectrum of 3b.

Figure S25: High-resolution mass spectrum of 3b.



Figure S26: ¹H-NMR (CDCl₃, 500 MHz) spectrum of 4b.



Figure S27: ¹³C-NMR (CDCl₃, 126 MHz) spectrum of 4b.



Figure S28: High-resolution mass spectrum of 1c.



Figure S30: ¹³C-NMR (CDCl₃, 126 MHz) spectrum of 1c.



Figure S31: High-resolution mass spectrum of 1c.



Figure S33: ¹³C-NMR (CDCl₃, 126 MHz) spectrum of 2c.



Figure S34: High-resolution mass spectrum of 2c.

UV-Vis absorption spectra



Figure S35: UV-Vis absorption spectrum of compound 1a recorded in CH₂Cl₂ at 25 °C.



Figure S36: UV-Vis absorption spectrum of compound 2a recorded in CH₂Cl₂ at 25 °C.



Figure S37: UV-Vis absorption spectrum of compound 3a recorded in CH₂Cl₂ at 25 °C.



Figure S38: UV-Vis absorption spectrum of compound 4a recorded in CH₂Cl₂ at 25 °C.



Figure S39: UV-Vis absorption spectrum of compound 1b recorded in CH₂Cl₂ at 25 °C.



Figure S40: UV-Vis absorption spectrum of compound 2b recorded in CH₂Cl₂ at 25 °C.



Figure S41: UV-Vis absorption spectrum of compound 3b recorded in CH₂Cl₂ at 25 °C.



Figure S42: UV-Vis absorption spectrum of compound 4b recorded in CH₂Cl₂ at 25 °C.



Figure S43: UV-Vis absorption spectrum of compound 1c recorded in CH₂Cl₂ at 25 °C.



Figure S44: UV-Vis absorption spectrum of compound 2c recorded in CH₂Cl₂ at 25 °C.