

Electrochemical-induced synthesis of quinazolinones via cathode hydration of *o*-aminobenzonitriles in aqueous solution

Li Yang,^a Huiqing Hou,^b Lan Li,^a Jin Wang,^b Sunying Zhou,^b Mei Wu,^b and Fang Ke^{*}
^{a,b}

^a. Faculty of Material and Chemical Engineering, Yibin University, Yibin 644000, China.

^b. Institute of MateriaMedica, School of Pharmacy, Fujian Provincial Key Laboratory of Natural Medicine Pharmacology, Fujian Medical University, Fuzhou 350122, China. E-mail: kefang@mail.fjmu.edu.cn; Fax: +86-591-22862016; Tel: +86-591-22862016, +86-831-3545095.

Supporting information

1.	General information	1
2.	General procedure for the catalytic reactions.....	1
3.	General procedure for the gram scale experiment	1
4.	Cyclic voltammetry experiment.....	1
5.	Synthesis of N-(2,5-dichlorophenyl)-6-(2,2,2-trifluoroethoxy) pteridin-4-amine (D3).	1
	Experimental procedures and characterization data	2
	¹ H NMR and ¹³ C NMR spectra for the products.....	9

1. General information

All reagents were purchased from commercial sources and used without further purification. All solvents were dried in a standard manner. Reactions were monitored by TLC on silica gel plates. Column chromatography was performed over silica gel (200-300 mesh) and a petroleum ether/ethyl. Shanghai chenhua CHI600E electrochemical workstation was used in the standard configuration as delivered, including proprietary software. All products were characterized by NMR. ^1H NMR spectra were recorded at 400 MHz and ^{13}C NMR spectra were recorded at 101 MHz (Bruker DPX) with CDCl_3 and DMSO-d_6 as solvent. Chemical shifts are reported in ppm using TMS as internal standard. NMR by the services provided at the Shandong Liaocheng University. HPLC were recorded on an SHIMDZU LC-20A instrument with a HP5-MS 30 m x 0.25 mm capillary apolar columns. A549, HCT-116 and SGC-7901 cells were all obtained from the Cell Bank of the Chinese Academy of Sciences.

2. General procedure for the catalytic reactions


A undivided cell equipped was charged with *o*-aminobenzonitrile (0.70 mmol) and benzaldehyde (0.50 mmol), I_2 (0.4 mmol), NaOH (2.0 mmol) and water (3 mL). The mixture was stirred at room temperature for 12 h in the 80 mA circuit. After the reaction was completed the solution of the crude product was concentrated in vacuo, and the residue was purified by column chromatography on a silica gel (petroleum ether/ethyl acetate=3/1) to afford the target product as a white solid.

3. General procedure for the gram scale experiment


In 50 mL undivided cell equipped with *o*-aminobenzonitriles (7.0 mmol), benzaldehydes (5.0 mmol), I_2 (2.0 mmol), NaOH (10 mmol) and water (10 mL).The mixture was stirred at room temperature for 12 h in the 80mA circuit. After the reaction was completed the solution of the crude product was concentrated in vacuo, and the residue was purified by column chromatography on a silica gel (petroleum ether/ethyl acetate=3/1) to afford the target product as a white solid.

4. Cyclic voltammetry experiment

Cyclic voltammograms were measured using Shanghai chenhua CHI600E electrochemical workstation with electrochemical analysis software, using a conventional three-electrode cell. The working electrode was a glassy carbon working electrode, the counter and reference electrodes consisted of a Pt wire and a SCE, respectively. The glassy carbon working electrode was polished with a polishing cloth before each measurement. The concentration of all tested compounds was 1 mmol/L. The scan rate was 0.1 V/s.




5. Synthesis of N-(2,5-dichlorophenyl)-6-(2,2,2-trifluoroethoxy) pteridin-4-amine (D3).



- (1) The reaction of 3-amino-6-(2,2,2-trifluoroethoxy)pyrazine-2-carbonitrile with formaldehyde (1.2 mmol) was carried out for 12 hours under the irradiation of 80 mA circuit with I₂ (40 mmol%), NaOH (2 mmol) as base and 4 mL water as solvent, the yield of 6-(2,2,2-trifluoroethoxy)pteridin-4(3H)-one was 81%.
- (2) In a 100 mL flask, 6-trifluoroethoxy-4(3h)-Pteridine (2 mmol), 10 mL thionyl dichloride, and 1-2 drops of DMF were added. Vacuum distillation removes dichlorothionyl and is then dried in vacuum for 30 minutes before being cooled to a yellow solid. It was slowly added into 10 mL ice water, then reaction mixture was added NaHCO₃ to adjust pH value to 7, and stirred for 10 minutes, filtered, and dried in a vacuum to obtain 6-(2,2,2-trifluoroethoxy)-4-chloro-pteridine as pale yellow powder. The yield was 86%.
- (3) 6-trifluoroethoxy-4-chloro-pteridine (2 mmol), aromatic amine (2 mmol) and isopropanol were added into 50 mL single-port round-bottom flask and stirred evenly for several minutes. Then a drying tube of anhydrous calcium chloride was connected to the upper end of the spherical condensing tube, and the optimal reaction conditions for dichloromethane dissolution were optimized by adjusting the microwave power and reaction time. After the reaction was followed by thin layer chromatography (TLC), it was cooled to 25 °C and then extracted with 50 mL water and ethyl acetate (30 mL × 3) to concentrate the organic phase of the pale yellow crude. The crude product is separated by silica gel column Chromatography (gradient elution of petroleum ether from 1:4 to 1:2 Ethyl acetate) and the final product is obtained.


Experimental procedures and characterization data

2-phenylquinazolin-4(3H)-one (3aa)¹




¹H NMR (400 MHz, DMSO-d₆) δ 12.54 (s, 1H), 8.25-8.11 (m, 3H), 7.89-7.81 (m, 1H), 7.75 (d, *J* = 8.0 Hz, 1H), 7.65-7.48 (m, 4H). ¹³C NMR (101 MHz, DMSO-d₆) δ 162.74, 152.81, 149.18, 135.07, 133.20, 131.86, 129.08, 128.24, 127.94, 127.06, 126.33, 121.44. MS [EI, m/z]: 222 [M⁺].

2-(3-chlorophenyl)quinazolin-4(3H)-one (3ab)²




¹H NMR (400 MHz, DMSO-d₆+CDCl₃) δ 8.24 (t, *J* = 1.8 Hz, 1H), 8.17-8.12 (m, 2H), 7.78 (ddd, *J* = 8.4, 7.0, 1.5 Hz, 1H), 7.74-7.70 (m, 1H), 7.56-7.45 (m, 3H). ¹³C NMR (101 MHz, DMSO-d₆+CDCl₃) δ 162.72, 151.32, 148.88, 135.08, 134.72, 134.16, 131.35, 130.50, 128.03, 127.85, 126.99, 126.54, 126.22, 121.58. MS [EI, m/z]: 256 [M⁺].

2-(2-chlorophenyl)quinazolin-4(3H)-one (3ac)²




¹H NMR (400 MHz, DMSO-d₆) δ 12.65 (s, 1H), 8.19 (dd, *J* = 7.9, 1.2 Hz, 1H), 7.89-7.83 (m, 1H), 7.76-7.66 (m, 2H), 7.64-7.55 (m, 3H), 7.50 (td, *J* = 7.4, 1.4 Hz, 1H). **¹³C NMR (101 MHz, DMSO-d₆)** δ 161.90, 152.73, 149.06, 135.06, 134.29, 132.08, 131.97, 131.35, 130.07, 127.95, 127.70, 127.54, 126.32, 121.73. **MS [EI, m/z]:** 256[M⁺].

2-(2-fluorophenyl)quinazolin-4(3H)-one (3ad)³




¹H NMR (400 MHz, DMSO-d₆) δ 12.57 (s, 1H), 8.18 (dd, *J* = 7.9, 1.2 Hz, 1H), 7.90-7.83 (m, 1H), 7.80 (td, *J* = 7.6, 1.7 Hz, 1H), 7.74 (d, *J* = 7.7 Hz, 1H), 7.67-7.60 (m, 1H), 7.60-7.54 (m, 1H), 7.43-7.35 (m, 2H). **¹³C NMR (101 MHz, DMSO-d₆)** δ 161.96, 161.29, 158.80, 150.42, 149.15, 135.08, 133.36, 133.28, 131.52, 131.50, 127.98, 127.51, 126.33, 125.10, 125.07, 122.82, 122.69, 121.59, 116.75, 116.54. **MS [EI, m/z]:** 240 [M⁺].

2-(4-bromophenyl)quinazolin-4(3H)-one (3ae)²




¹H NMR (400 MHz, DMSO-d₆) δ 12.62 (s, 1H), 8.20-8.10 (m, 3H), 7.85 (ddd, *J* = 8.5, 7.1, 1.6 Hz, 1H), 7.81-7.71 (m, 3H), 7.57-7.51 (m, 1H). **¹³C NMR (101 MHz, DMSO-d₆)** δ 162.71, 151.98, 148.98, 135.18, 132.39, 132.11, 130.28, 127.95, 127.29, 126.36, 125.71, 121.45. **MS [EI, m/z]:** 300 [M⁺].

2-(2-hydroxyphenyl)quinazolin-4(3H)-one (3af)⁴




¹H NMR (400 MHz, DMSO-d₆) δ 13.78 (s, 1H), 12.47 (s, 1H), 8.24 (d, *J* = 8.0 Hz, 1H), 8.17 (d, *J* = 7.6 Hz, 1H), 7.86 (d, *J* = 7.0 Hz, 1H), 7.78 (d, *J* = 8.0 Hz, 1H), 7.57 (d, *J* = 7.8 Hz, 1H), 7.47 (t, *J* = 7.7 Hz, 1H), 7.05-6.94 (m, 2H). **¹³C NMR (101 MHz, DMSO-d₆)** δ 161.93, 160.51, 154.22, 151.01, 146.66, 135.52, 134.21, 128.20, 127.46, 126.53, 121.39, 119.32, 118.37, 114.24. **MS [EI, m/z]:** 239 [M⁺].

2-(2,4-dimethylphenyl)quinazolin-4(3H)-one (3ag)⁵




¹H NMR (400 MHz, DMSO-d₆) δ 12.37 (s, 1H), 8.16 (dd, *J* = 7.9, 1.2 Hz, 1H), 7.87-7.81 (m, 1H), 7.68 (d, *J* = 7.6 Hz, 1H), 7.57-7.51 (m, 1H), 7.41 (d, *J* = 7.7 Hz, 1H), 7.18-7.12 (m, 2H), 2.37 (s, 3H), 2.35 (s, 3H). **¹³C NMR (101 MHz, DMSO-d₆)** δ 162.38, 154.90, 149.22, 139.96, 136.53, 134.90, 131.88, 131.67, 129.65, 127.78, 126.99, 126.72, 126.24, 121.35, 21.31, 20.06. **MS [EI, m/z]:** 250 [M⁺].

2-(4-methoxyphenyl)quinazolin-4(3H)-one (3ah)²




¹H NMR (400 MHz, DMSO-d₆) δ 12.41 (s, 1H), 8.23-8.18 (m, 2H), 8.14 (dd, *J* = 7.9, 1.2 Hz, 1H), 7.82 (ddd, *J* = 8.5, 7.1, 1.6 Hz, 1H), 7.71 (d, *J* = 8.1 Hz, 1H), 7.52-7.46 (m, 1H), 7.12-7.07 (m, 2H), 3.86 (s, 3H). **¹³C NMR (101 MHz, DMSO-d₆)** δ 162.77, 162.35, 152.34, 149.41, 135.02, 129.93, 127.76, 126.60, 126.30, 125.28, 121.16, 114.47, 55.94. **MS [EI, m/z]:** 252 [M⁺].

2-(p-tolyl)quinazolin-4(3H)-one (3ai)²




¹H NMR (400 MHz, DMSO-d₆) δ 12.47 (s, 1H), 8.15 (dd, *J* = 7.9, 1.2 Hz, 1H), 8.11 (d, *J* = 8.3 Hz, 2H), 7.84 (ddd, *J* = 8.4, 7.2, 1.5 Hz, 1H), 7.73 (d, *J* = 7.6 Hz, 1H), 7.55-7.49 (m, 1H), 7.36 (d, *J* = 8.1 Hz, 2H), 2.40 (s, 3H). **¹³C NMR (101 MHz, DMSO-d₆)** δ 162.72, 152.70, 149.35, 141.93, 135.06, 130.36, 129.67, 128.16, 127.89, 126.88, 126.31, 121.37, 21.46. **MS [EI, m/z]:** 236 [M⁺].

2-benzylquinazolin-4(3H)-one (3aj)²




¹H NMR (400 MHz, DMSO-d₆) δ 12.42 (s, 1H), 8.08 (dd, *J* = 7.9, 1.3 Hz, 1H), 7.78 (ddd, *J* = 8.5, 7.2, 1.6 Hz, 1H), 7.61 (d, *J* = 7.7 Hz, 1H), 7.50-7.45 (m, 1H), 7.39 (d, *J* = 7.1 Hz, 2H), 7.33 (t, *J* = 7.5 Hz, 2H), 7.25 (t, *J* = 7.2 Hz, 1H), 3.95 (s, 2H). **¹³C NMR (101 MHz, DMSO-d₆)** δ 162.37, 156.51, 149.25, 137.01, 134.87, 129.35, 128.97, 127.35, 126.70, 126.19, 121.20, 41.24. **MS [EI, m/z]:** 236 [M⁺].

2-(thiophen-2-yl)quinazolin-4(3H)-one (3ak)²




¹H NMR (400 MHz, DMSO-d₆) δ 12.46 (s, 1H), 8.61 (dd, *J* = 2.9, 1.3 Hz, 1H), 8.14 (dd, *J* = 7.9, 1.3 Hz, 1H), 7.88 (dd, *J* = 5.1, 1.3 Hz, 1H), 7.82 (ddd, *J* = 8.5, 7.2, 1.6 Hz, 1H), 7.74-7.66 (m, 2H), 7.54-7.47 (m, 1H). **¹³C NMR (101 MHz, DMSO-d₆)** δ 162.54, 149.34, 148.76, 135.88, 135.05, 129.12, 127.83, 127.77, 127.52, 126.82, 126.35, 121.43. **MS [EI, m/z]:** 228 [M⁺].

2-(pyridin-3-yl)quinazolin-4(3H)-one (3al)⁶




¹H NMR (400 MHz, DMSO-d₆) δ 12.74 (s, 1H), 9.30 (d, *J* = 1.7 Hz, 1H), 8.77 (dd, *J* = 4.8, 1.5 Hz, 1H), 8.54-8.48 (m, 1H), 8.18 (dd, *J* = 7.9, 1.2 Hz, 1H), 7.91-7.84 (m, 1H), 7.78 (d, *J* = 7.7 Hz, 1H), 7.63-7.54 (m, 2H). **¹³C NMR (101 MHz, DMSO-d₆)** δ 162.56, 152.30, 151.23, 149.23, 149.01, 135.87, 135.20, 129.20, 128.07, 127.45, 126.36, 124.01, 121.62. **MS [EI, m/z]:** 223 [M⁺].

6-methoxy-2-phenylquinazolin-4(3H)-one (3am)¹




¹H NMR (400 MHz, DMSO-d₆) δ 12.51 (s, 1H), 8.17 (dd, *J* = 8.0, 1.7 Hz, 2H), 7.70 (d, *J* = 8.9 Hz, 1H), 7.60-7.51 (m, 4H), 7.45 (dd, *J* = 8.9, 3.0 Hz, 1H), 3.90 (s, 3H). **¹³C NMR (101 MHz, DMSO-d₆)** δ 162.51, 158.22, 150.56, 143.68, 133.26, 131.52, 129.70, 129.05, 127.96, 124.59, 122.26, 106.33, 56.12. **MS [EI, m/z]:** 252 [M⁺].

8-methyl-2-phenylquinazolin-4(3H)-one (3an)⁷




¹H NMR (400 MHz, DMSO-d₆) δ 12.52 (s, 1H), 8.27-8.20 (m, 2H), 8.00 (d, *J* = 7.9 Hz, 1H), 7.70 (d, *J* = 7.2 Hz, 1H), 7.63-7.53 (m, 3H), 7.40 (t, *J* = 7.6 Hz, 1H), 2.63 (s, 3H). **¹³C NMR (101 MHz, DMSO-d₆)** δ 163.02, 151.50, 147.59, 136.08, 135.38, 133.43, 131.79, 129.09, 128.20, 126.53, 123.96, 121.36, 17.63. **MS [EI, m/z]:** 236 [M⁺].

7-nitro-2-phenylquinazolin-4(3H)-one (3ao)⁶




¹H NMR (400 MHz, DMSO-d₆) δ 12.92 (s, 1H), 8.45 (d, *J* = 2.2 Hz, 1H), 8.38 (d, *J* = 8.7 Hz, 1H), 8.26-8.21 (m, 3H), 7.66-7.57 (m, 3H). **¹³C NMR (101 MHz, DMSO-d₆)** δ 161.99, 155.10, 151.84, 132.64, 132.47, 130.12, 129.19, 128.68, 128.54, 125.82, 122.83, 120.53. **MS [EI, m/z]:** 267 [M⁺].

6-chloro-2-phenylquinazolin-4(3H)-one (3ap)²




¹H NMR (400 MHz, DMSO-d₆) δ 12.71 (s, 1H), 8.22-8.15 (m, 2H), 8.10 (d, *J* = 2.5 Hz, 1H), 7.87 (dd, *J* = 8.7, 2.5 Hz, 1H), 7.77 (d, *J* = 8.7 Hz, 1H), 7.62-7.53 (m, 3H). **¹³C NMR (101 MHz, DMSO-d₆)** δ 161.77, 153.31, 147.97, 135.18, 132.93, 132.07, 131.25, 130.23, 129.11, 128.32, 125.35, 122.72. **MS [EI, m/z]: 256 [M⁺].**

6-bromo-2-phenylquinazolin-4(3H)-one (3aq)¹




¹H NMR (400 MHz, DMSO-d₆) δ 12.72 (s, 1H), 8.24 (d, *J* = 2.3 Hz, 1H), 8.18 (dd, *J* = 6.8, 1.6 Hz, 2H), 7.99 (dd, *J* = 8.7, 2.4 Hz, 1H), 7.70 (d, *J* = 8.7 Hz, 1H), 7.61 (d, *J* = 7.0 Hz, 1H), 7.57 (d, *J* = 7.5 Hz, 2H). **¹³C NMR (101 MHz, DMSO-d₆)** δ 161.63, 153.42, 148.24, 137.91, 132.95, 132.10, 130.38, 129.12, 128.47, 128.33, 123.10, 119.43. **MS [EI, m/z]: 300 [M⁺].**

6-nitro-2-phenylquinazolin-4(3H)-one (3ar)²



¹H NMR (400 MHz, DMSO-d₆) δ 12.99 (s, 1H), 8.84 (d, *J* = 2.6 Hz, 1H), 8.56 (dd, *J* = 9.0, 2.7 Hz, 1H), 8.23 (d, *J* = 7.4 Hz, 2H), 7.92 (d, *J* = 9.0 Hz, 1H), 7.66 (t, *J* = 7.2 Hz, 1H), 7.59 (t, *J* = 7.3 Hz, 2H). **¹³C NMR (101 MHz, DMSO-d₆)** δ 162.05, 156.19, 153.39, 145.17, 132.76, 132.53, 129.65, 129.21, 128.99, 128.73, 122.52, 121.26. **MS [EI, m/z]: 267 [M⁺].**

2-(4-chlorophenyl)-8-methylquinazolin-4(3H)-one (3as)¹¹



¹H NMR (400 MHz, DMSO-d₆) δ 12.71 (s, 1H), 8.22-8.15 (m, 2H), 8.10 (d, *J* = 2.5 Hz, 1H), 7.87 (dd, *J* = 8.7, 2.5 Hz, 1H), 7.77 (d, *J* = 8.7 Hz, 1H), 7.62-7.53 (m, 3H), 2.51 (t, *J* = 1.9 Hz, 3H). **¹³C NMR (101 MHz, DMSO-d₆)** δ 161.77, 153.31, 147.97, 135.18, 132.93, 132.07, 131.25, 130.23, 129.11, 128.32, 127.80, 125.35, 122.72. **MS [EI, m/z]: 270 [M⁺].**


6-methoxy-2-(2-methoxyphenyl)quinazolin-4(3H)-one (3at)⁸



¹H NMR (400 MHz, DMSO-d₆) δ 12.02 (s, 1H), 7.71 (dd, *J* = 7.6, 1.8 Hz, 1H), 7.66 (d, *J* = 8.9 Hz, 1H), 7.57-7.49 (m, 2H), 7.44 (dd, *J* = 8.9, 3.0 Hz, 1H), 7.19 (d, *J* = 8.1 Hz, 1H), 7.09 (td, *J* = 7.5, 0.8 Hz, 1H), 3.90 (s, 3H), 3.87 (s, 3H). **¹³C NMR (101 MHz, DMSO-d₆)** δ 161.48, 158.17, 157.59, 150.53,


144.00, 132.47, 130.88, 129.58, 124.38, 123.14, 122.27, 120.91, 112.33, 106.25, 56.24, 56.12. **MS** [EI, m/z]: 282 [M⁺].

2-(3-methoxyphenyl)-6-methylquinazolin-4(3H)-one (3au)¹²




¹H NMR (400 MHz, DMSO-d₆) δ 12.45 (s, 1H), 7.96 (s, 1H), 7.78 (d, *J* = 7.8 Hz, 1H), 7.76-7.72 (m, 1H), 7.66 (s, 2H), 7.46 (t, *J* = 8.0 Hz, 1H), 7.14 (dd, *J* = 7.9, 2.2 Hz, 1H), 3.87 (s, 3H), 2.47 (s, 3H). **¹³C NMR (101 MHz, DMSO-d₆)** δ 162.61, 159.83, 151.67, 147.15, 136.85, 136.37, 134.57, 130.20, 127.91, 125.72, 121.25, 120.47, 117.92, 112.89, 55.86, 21.32. **MS** [EI, m/z]: 266 [M⁺].

6-methyl-2-phenylquinazolin-4(3H)-one (3av)¹




¹H NMR (400 MHz, DMSO-d₆) δ 12.45 (s, 1H), 8.18 (dd, *J* = 8.1, 1.6 Hz, 2H), 7.96 (s, 1H), 7.65 (s, 2H), 7.60-7.51 (m, 3H), 2.46 (s, 3H), 2.45 (s, 1H). **¹³C NMR (101 MHz, DMSO-d₆)** δ 162.62, 151.93, 147.22, 136.77, 136.33, 133.26, 131.69, 129.05, 128.11, 127.85, 125.71, 121.20, 21.32. **MS** [EI, m/z]: 236 [M⁺].

2-phenylpteridin-4(3H)-one (3aw)⁹




¹H NMR (400 MHz, DMSO-d₆) δ 11.32 (s, 1H), 8.40 (s, 1H), 8.01-7.89 (m, 3H), 7.72-7.65 (m, 2H), 7.60 (t, *J* = 7.4 Hz, 2H). **¹³C NMR (101 MHz, DMSO-d₆)** δ 169.07, 164.94, 164.82, 156.18, 155.81, 149.63, 147.28, 133.86, 133.53, 131.85, 131.42, 129.48, 128.20, 123.81. **MS** [EI, m/z]: 224 [M⁺].

6-methoxypteridin-4(3H)-one (3ax)¹⁰




¹H NMR (400 MHz, DMSO-d₆) δ 12.75 (s, 1H), 8.69 (s, 1H), 8.23 (s, 1H), 4.02 (s, 3H). **¹³C NMR (101 MHz, DMSO-d₆)** δ 160.59, 158.41, 151.32, 146.58, 143.50, 130.42, 54.76. **MS** [EI, m/z]: 178 [M⁺].

2-pentylquinazolin-4(3H)-one (3ay)¹⁴




¹H NMR (500 MHz, DMSO-d₆) δ 12.09 (s, 1H), 7.99 (dd, *J* = 7.9, 1.3 Hz, 1H), 7.70 - 7.66 (m, 1H), 7.51 (d, *J* = 7.8 Hz, 1H), 7.37 (t, *J* = 7.5 Hz, 1H), 2.53- 2.47 (m, 2H), 1.67 - 1.60 (m, 2H), 1.26-1.19 (m, 4H), 0.78 (t, *J* = 6.9 Hz, 3H); **¹³C NMR (126 MHz, DMSO-d₆)** δ 161.47, 157.17, 148.59, 133.93, 126.43, 125.56, 125.31, 120.39, 34.10, 30.39, 26.14, 21.48, 13.50; **MS [EI, m/z]:** 216 [M⁺].

N-(2,5-dichlorophenyl)-6-(2,2,2-trifluoroethoxy)pteridin-4-amine¹³




¹H NMR (400 MHz, DMSO-d₆) δ 10.01 (s, 1H), 9.06 (s, 1H), 8.71 (s, 1H), 8.13 (d, *J* = 2.5 Hz, 1H), 7.68 (d, *J* = 8.6 Hz, 1H), 7.42 (dd, *J* = 8.6, 2.5 Hz, 1H), 5.39 (q, *J* = 9.0 Hz, 2H). **¹³C NMR (101 MHz, DMSO-d₆)** δ 158.50, 156.39, 155.41, 150.88, 145.97, 136.68, 132.29, 131.52, 127.72, 127.38, 126.96, 122.91, 121.15, 63.54, 63.19, 62.84, 62.49. **MS [EI, m/z]:** 389 [M⁺].

o-aminobenzamide (4a)¹⁵



¹H NMR (400 MHz, DMSO-d₆) δ 7.74 (s, 1H), 7.55 (dd, *J* = 7.9, 1.6 Hz, 1H), 7.14 (ddd, *J* = 8.4, 7.0, 1.5 Hz, 2H), 6.69 (dd, *J* = 8.3, 1.4 Hz, 1H), 6.56 (s, 2H), 6.52 – 6.45 (m, 1H); **¹³C NMR (101 MHz, DMSO-d₆)** δ 171.83, 132.40, 129.23, 116.90, 114.89, 114.16; **MS [EI, m/z]:** 136 [M⁺].


(E)-2-(benzylideneamino)benzamide (5aa)¹⁶



¹H NMR (400 MHz, DMSO-d₆) δ 8.28 (s, 1H), 7.62 (dd, *J* = 7.8, 1.8 Hz, 1H), 7.51 (d, *J* = 1.9 Hz, 1H), 7.49 (d, *J* = 1.3 Hz, 1H), 7.42-7.38 (m, 2H), 7.37-7.34 (m, 1H), 7.25 (ddd, *J* = 8.8, 7.1, 1.6 Hz, 1H), 7.10 (d, *J* = 1.7 Hz, 1H), 6.75 (d, *J* = 8.3 Hz, 1H), 6.71-6.65 (m, 1H), 5.76 (t, *J* = 2.0 Hz, 1H).

¹³C NMR (101 MHz, DMSO-d₆) δ 164.05, 148.33, 142.11, 133.78, 128.91, 128.79, 127.81, 127.32, 117.58, 115.43, 114.87, 67.03; **MS [EI, m/z]:** 224 [M⁺].

2-phenyl-2,3-dihydroquinazolin-4(1H)-one (6aa)¹⁷




¹H NMR (600 MHz, DMSO-d₆) δ 8.35 (s, 1H), 7.67 (dd, *J* = 7.8, 1.8 Hz, 1H), 7.56-7.52 (m, 2H), 7.43- 7.35 (m, 3H), 7.27 (ddd, *J* = 8.6, 7.2, 1.7 Hz, 1H), 7.16 (s, 1H), 6.80 (d, *J* = 8.2 Hz, 1H), 6.73- 6.69 (m, 1H), 5.81 (s, 1H); **¹³C NMR (151 MHz, DMSO-d₆)** δ 164.17, 148.38, 142.10, 133.84, 128.96, 128.82, 127.88, 127.36, 117.64, 115.44, 114.92, 67.11; **MS [EI, m/z]:** 224 [M⁺].

References



1. F.-C. Jia , Z.-W. Zhou, C. Xu, Y.-D. Wu and A.-X. Wu, *Org. Lett.*, **2016**, *18*, 2942-2945.
2. J. Zhou and J. Fang, *J. Org. Chem.*, **2011**, *76*, 7730-7736.
3. S.Guo, Y. Li , L. Tao, W. Zhang and X. Fan, *RSC Adv.*, **2014**, *4*, 59289-59296.
4. S. Chatterjee, R.Srinath, S.Bera, K. Khamaru, A. Rahman and B. Banerji, *Org. Lett.*, **2019**, *21*, 9028-9032.
5. S. Maiti, J. Kim, J.-H. Park, D. Nam, J. B. Lee, Y.-J. Kim, J.-M. Kee, J. K. Seo, K. Myung, J.-U. Rohde, W. Choe, O.-H. Kwon and S. Y. Hong, *J. Org. Chem.*, **2019**, *84*, 6737-6751.
6. W. Xu , Y. Jin, H. Liu, Y. Jiang and H. Fu, *Org. Lett.*, **2011**, *13*, 1274-1277.
7. S. Das, S. Sinha, D. Samanta, R. Mondal, G. Chakraborty, P. Brandaõ and N. D. Paul, *J. Org. Chem.*, **2019**, *84*, 10160-10171.
8. M. Hayakawa, H. Kaizawa, H. Moritomo, T. Koizumi, T. Ohishi, M. Okada, M. Ohta, S.-i.Tsukamoto, P. Parker, P. Workman and M. Waterfield, *Bioorg. Med. Chem.*, **2006**, *14*, 6847-6858.
9. T. Okawa, M. Kawase and S. Eguchi, *Synthesis*, **1998**, *8*, 1185-9400.
10. C. Duan, J. Jia, R. Zhu and J. Wang, *J. Heterocycl. Chem.*, **2012**, *49*, 865-8722.
11. A. Nathubhai, P. J. Wood, M. D. Liroy, A. S. Thompson and M. D. Threadgill, *ACS Med. Chem. Lett.*, **2013**, *4*, 1173-1177.
12. Z.-Y. Yu, M.-Y. Chen, J.-X. He, D.-J. Tao, J.-J. Yuan, Y.-Y. Peng and Z.-B. Song, *Mol.Catal.*, **2017**, *434*,134-139.
13. J. Lin, P. Wang, Z. Zhang, G. Xue, D. Zha, J. Wang, X. Xu and Z. Li, *Synth. Commun.*, **2020**, *50*, 823-830.
14. X. Tian, L. Song, E. Li, Q. Wang, W. Yu and J. Chang, *RSC Adv.*, **2015**, *5*, 62194-62201.
15. W. Li, J. Artz, C. Broicher, K. Junge, H. Hartmann, A. Besmehn, R. Palkovits and M. Beller, *Catal. Sci. Technol.*, **2019**, *9*, 157-162.
16. J. K. Laha, K. S. S. Tummalaapalli, A. Nair and N. Patel, *J. Org. Chem.*, **2015**, *80*, 22, 11351-11359.
17. (a)W. Zhao, P. Liu and F. Liu, *ChemCatChem*, **2016**, *8*, 1523-1530; (b) W. Gong, X. Chen, H. Jiang, D. Chu, Y. Cui and Y. Liu, *J. Am. Chem. Soc.*; **2019**, *141*, 7498-7508; (c) H. Hikawa, Y. Ino, H. Suzuki and Y. Yokoyama, *J. Org. Chem.*, **2012**, *77*, 7046-7051.

¹H NMR and ¹³C NMR spectra for the products


2-phenylquinazolin-4(3H)-one (3aa)




2-(3-chlorophenyl)quinazolin-4(3H)-one (3ab)




2-(2-chlorophenyl)quinazolin-4(3H)-one (3ac)




2-(2-fluorophenyl)quinazolin-4(3H)-one (3ad)




2-(4-bromophenyl)quinazolin-4(3H)-one (3ae)




2-(2-hydroxyphenyl)quinazolin-4(3H)-one (3af)




2-(2,4-dimethylphenyl)quinazolin-4(3H)-one (3ag)




2-(4-methoxyphenyl)quinazolin-4(3H)-one (3ah)




2-(p-tolyl)quinazolin-4(3H)-one (3ai)






2-benzylquinazolin-4(3H)-one (3aj)






2-(thiophen-2-yl)quinazolin-4(3H)-one (3ak)




2-(pyridin-3-yl)quinazolin-4(3H)-one (3al)




6-methoxy-2-phenylquinazolin-4(3H)-one (3am)





8-methyl-2-phenylquinazolin-4(3H)-one (3an)




7-nitro-2-phenylquinazolin-4(3H)-one (3ao)




6-chloro-2-phenylquinazolin-4(3H)-one (3ap)




6-bromo-2-phenylquinazolin-4(3H)-one (3aq)




6-nitro-2-phenylquinazolin-4(3H)-one (3ar)




2-(4-chlorophenyl)-8-methylquinazolin-4(3H)-one (3as)




6-methoxy-2-(2-methoxyphenyl)quinazolin-4(3H)-one (3at)




2-(3-methoxyphenyl)-6-methylquinazolin-4(3H)-one (3au)




6-methyl-2-phenylquinazolin-4(3H)-one (3av)




2-phenylpteridin-4(3H)-one (3aw)




6-methoxypteridin-4(3H)-one (3ax)





2-pentylquinazolin-4(3H)-one (3ay)



N-(2,5-dichlorophenyl)-6-(2,2,2-trifluoroethoxy)pteridin-4-amine



o-aminobenzamide(4a)



(E)-2-(benzylideneamino)benzamide (5aa)



2-phenyl-2,3-dihydroquinazolin-4(1H)-one (6aa)

