

1 Mayenite-Based Electride C<sub>12</sub>A<sub>7</sub>e<sup>-</sup>: An  
2 Innovative Synthetic Method via Plasma  
3 Arc Melting - Supplementary Information

4

5 **Sebastian Weber <sup>#,a</sup>, Sebastian Schäfer <sup>a</sup>, Mattia Saccoccio <sup>a</sup>, Karsten Seidel <sup>b</sup>, Holger Kohlmann <sup>c</sup>,**  
6 **Roger Gläser <sup>d</sup> and Stephan A. Schunk <sup>a,b,\*</sup>**

7 <sup>a</sup> hte GmbH, Kurpfalzring 104, 69123 Heidelberg, Germany; stephan.schunk@hte-company.de.

8 <sup>b</sup> BASF SE, Carl-Bosch-Strasse 38, 67056 Ludwigshafen am Rhein, Germany; karsten.seidel@basf.com.

9 <sup>c</sup> Institute of Inorganic Chemistry, Leipzig University, Johannisallee 29, 04103 Leipzig, Germany;  
10 holger.kohlmann@uni-leipzig.de

11 <sup>d</sup> Institute of Chemical Technology, Leipzig University, Linnéstraße 3, 04103 Leipzig, Germany;  
12 roger.glaeser@uni-leipzig.de.


13 <sup>#</sup> New address: Institute for Chemical Technology and Polymer Chemistry (ITCP); Institute of Catalysis  
14 Research and Technology (IKFT), Karlsruhe Institute of Technology (KIT), Engesserstraße 20, 76131  
15 Karlsruhe, Germany; sebastian.weber@kit.edu

16 \*Corresponding author.

17

18 Calibration plasma power

19



20

21 Figure S1 Plasma power calculated from the measured current for different gas atmospheres (95%Ar/5%H<sub>2</sub> in red and Ar in  
22 blue) depending on the plasma intensity level of the arc furnace.

23