

Supplementary Information

Analysis for the effects of Cu-MOFs on fungal cell inactivation

Mayura Veerana, †^a Hyun-Chul Kim, †^b Sarmistha Mitra,^a Bishwa Chandra Adhikari,^a

*Gyungsoon Park,^{*a} Seong Huh,^b Sung-Jin Kim,^c and Youngmee Kim^{*c}*

^a Plasma Bioscience Research Center and Department of Electrical and Biological Physics, Kwangwoon University, Seoul 01897, Republic of Korea. E-mail: gyungp@kw.ac.kr

^b Department of Chemistry and Protein Research Centre for Bio-Industry, Hankuk University of Foreign Studies, Yongin 17035, Korea

^c Institute of Nano-Bio Technology and Department of Chemistry and Nano Science, Ewha Womans University, Seoul 120-750, Korea. E-mail: ymeekim@ewha.ac.kr; Fax: +82-2-3277-2385; Tel: +82-2-3277-4164

† These authors contributed equally to this work.

Correspondence and requests for materials should be addressed to Y.K. (email: ymeekim@ewha.ac.kr) or to G.P. (email: gyungp@kw.ac.kr).




Fig. S1. IR spectra of of Cu-MOFs 1 - 3.





Fig. S2. Effect of collision after shaking Cu-MOFs on viability of *C. albicans* (a) and *A. niger* (b) cells at 4 days.



Scale bar = 10 μ m





Fig. S3. TUNNEL assay showing apoptosis-like cell death in *C. albicans*.



Scale bar = 10 μ m




Fig. S4. TUNNEL assay showing apoptosis-like cell death in *A. niger*.