

## SUPPORTING INFORMATION

# Brønsted acidic ionic liquid anchored on magnetite nanoparticles as a novel recoverable heterogeneous catalyst in the Biginelli reaction

Hourieh Sadat Oboudatian, Hossein Naeimi, Mohsen Moradian\*

Department of Organic Chemistry, Faculty of Chemistry, University of Kashan, Kashan, I.R. Iran


### Table of Contents

#### Details of Experimental Section


2. The details for the synthesis of compound 4a-4o.....	S3
2.3. General procedure for the synthesis of 3,4-Dihydropyrimidinones .....	S3
Scheme 1. of Preparation of 3,4-dihydropyrimidinones by $\text{Fe}_3\text{O}_4@\text{SiO}_2/\text{TES-Mo}^+\text{HSO}_4^-$ ....	S4
The FT-IR spectra of compound 4a.....	S4
The $^1\text{H}$ NMR spectra of 4a.....	S4
The FT-IR spectra of compound 4b.....	S5
The $^1\text{H}$ NMR spectra of 4b.....	S5
The FT-IR spectra of compound 4c.....	S6
The $^1\text{H}$ NMR spectrum of 4c.....	S6
The FT-IR spectra of compound 4d.....	S7
The $^1\text{H}$ NMR spectrum of 4d.....	S7
The FT-IR spectra of compound 4e.....	S8

The <sup>1</sup> H NMR spectrum of <b>4e</b> .....	S8
The FT-IR spectra of compound <b>4f</b> .....	S9
The <sup>1</sup> H NMR spectrum of <b>4f</b> .....	S9
The FT-IR spectra of compound <b>4g</b> .....	S10
The <sup>1</sup> H NMR spectrum of <b>4g</b> .....	S10
The FT-IR spectra of compound <b>4h</b> .....	S11
The <sup>1</sup> H NMR spectrum of <b>4h</b> .....	S11
The FT-IR spectra of compound <b>4i</b> .....	S12
The <sup>1</sup> H NMR spectrum of <b>4i</b> .....	S12
The FT-IR spectra of compound <b>4j</b> .....	S13
The <sup>1</sup> H NMR spectrum of <b>4j</b> .....	S13
The FT-IR spectra of compound <b>4k</b> .....	S14
The <sup>1</sup> H NMR spectrum of <b>4k</b> .....	S14
The FT-IR spectra of compound <b>4l</b> .....	S15
The <sup>1</sup> H NMR spectrum of <b>4l</b> .....	S15
The <sup>1</sup> H NMR spectra of <b>4l</b> in DMSO-D <sub>6</sub> and D <sub>2</sub> O as NMR solvent.....	S16
The FT-IR spectra of compound <b>4m</b> .....	S17
The <sup>1</sup> H NMR spectrum of <b>4m</b> .....	S17
The FT-IR spectra of compound <b>4n</b> .....	S18
The <sup>1</sup> H NMR spectrum of <b>4n</b> .....	S18
The <sup>1</sup> H NMR spectra of <b>4n</b> in CHCl <sub>3</sub> -D <sub>6</sub> and D <sub>2</sub> O as NMR solvent.....	S19
The FT-IR spectra of compound <b>4o</b> .....	S20
The <sup>1</sup> H NMR spectrum of <b>4o</b> .....	S20


### The FT-IR spectra of compound 4a




### The <sup>1</sup>H NMR spectrum of 4a




### The FT-IR spectra of compound **4b**




### The <sup>1</sup>H NMR spectrum of **4b**




### The FT-IR spectra of compound 4c




### The <sup>1</sup>H NMR spectrum of 4c




### The FT-IR spectra of compound 4d




### The <sup>1</sup>H NMR spectrum of 4d




### The FT-IR spectra of compound 4e




The  $^1\text{H}$  NMR spectrum of 4e




### The FT-IR spectra of compound 4f




### The <sup>1</sup>H NMR spectrum of 4f




### The FT-IR spectra of compound 4g




The  $^1\text{H}$  NMR spectrum of **4g**




The FT-IR spectra of compound **4h**




The  $^1\text{H}$  NMR spectrum of **4h**




The FT-IR spectra of compound **4i**




The  $^1\text{H}$  NMR spectrum of **4i**




The FT-IR spectra of compound **4j**




The  $^1\text{H}$  NMR spectrum of **4j**




The FT-IR spectra of compound **4k**




The  $^1\text{H}$  NMR spectrum of 4k




The FT-IR spectra of compound 4l




The  $^1\text{H}$  NMR spectrum of **4I**




The  $^1\text{H}$  NMR spectra of **4I** in  $\text{DMSO-D}_6$  and  $\text{D}_2\text{O}$  as NMR solvent




The FT-IR spectra of compound **4m**



The <sup>1</sup>H NMR spectrum of **4m**



The FT-IR spectra of compound **4n**



The <sup>1</sup>H NMR spectrum of **4n**

8.981  
8.040  
7.895  
7.568  
7.768  
7.723  
7.706  
7.680  
7.663  
7.579  
7.548  
7.504  
7.451  
7.402  
7.367  
7.346  
7.326  
7.302  
7.165  
7.141  
7.106  
7.086  
7.034  
7.013  
6.842  
6.652  
6.631  
6.552  
6.525  
6.451  
5.458  
5.119  
5.016  
4.797  
4.237  
4.176  
4.140  
4.043  
3.987  
3.970  
3.956  
3.940  
3.905





Current Data Parameters  
NAME 3H-4  
EXPNO 1089  
PROCNO 1


F2 - Acquisition Parameters  
Date\_ 20160821  
Time\_ 11.35  
INSTRUM spect  
PROBHD 5 mm Multinu  
PULPROG zg  
TD 32768  
SOLVENT CDCl3  
NS 8  
DS C  
SWH 8012.020 Hz  
FIDRES 0.244532 Hz  
AQ 2.0447731 sec  
RG 2298.8  
DW 62.400 usec  
DE 6.00 usec  
TE 300.0 K  
D1 5.00000000 sec

===== CHANNEL f1 =====  
NUC1 1H  
P1 15.00 usec  
PL1 -6.00 dB  
SFO1 400.1336012 MHz


F2 - Processing parameters  
SI 32768  
SF 400.1300665 MHz  
WDW FM  
SSB 0  
LB 0.30 Hz  
GB 0  
FC 1.00



The  $^1\text{H}$  NMR spectra of **4n** in  $\text{CHCl}_3\text{-D}_6$  and  $\text{D}_2\text{O}$  as NMR solvent



The FT-IR spectra of compound **4o**



The  $^1\text{H}$  NMR spectrum of **4o**

