Supporting Information

Interfacial behavior of β -lactoglobulin aggregates at oil-water interface studied using particle tracking and dilatational rheology

Nan Yang^{a,b,c,*}, Jing Ye^{a,}, Jing Li^{a,}, Bing Hu^a, Robert L. Leheny^{c,*}, Katsuyoshi Nishinari^{a,b}, Yapeng Fang ^{a,b,*}

Figure S1



Figure S1 Schematic of the sample cell for particle-tracking studies of the protein adsorption process.

Figure S2



Figure S2 Schematic of the sample cell for particle-tracking studies of the displacement of proteins by bile salts.

Figure S3

^{a.} Glyn O. Phillips Hydrocolloid Research Centre, National "111" Center for Cellular Regulation and Molecular Pharmaceutics, Key Laboratory of Fermentation Engineering (Ministry of Education), Department of Bioengineering and Food Science, Hubei University of Technology, Wuhan 430068, China.

^{b.} Food Hydrocolloid International Science and Technology Cooperation Base of Hubei Province, Hubei University of Technology, Wuhan 430068, China.

^{c.} Department of Physics and Astronomy, Johns Hopkins University, Baltimore, MD 21218, USA

^{*}Corresponding authors: N.Y., nanyang@hbut.edu.cn; R.L., leheny@jhu.edu; Y.F., ypfang@sjtu.edu.cn



Figure S3 TEM image and length distribution of unlyophilized β -lg F.

Figure S4



Figure S4 Final individual MSDs at the decane/water interface, during the adsorption of β -lg with a concentration of 110 µg/mL. This graph is for the interface after 65 minutes

Figure S5



Figure S5 Individual MSDs of tracers during the adsorption of BSs to the decane/water interface at the times indicated. BS concentration was 0.1 mg/mL.