

Supplementary Data

Specific detection of *Cronobacter sakazakii* in powdered infant formula using ssDNA aptamer

Hye Ri Kim^{a,b}, Myunghye Kim^c and Byoung Chan Kim^{a,b,*}

^a Center for Environment, Health and Welfare Research, Korea Institute of Science and Technology (KIST), Hwarangno 14-gil 5, Seongbuk-gu, Seoul 02792, Republic of Korea

^b Division of Energy and Environment Technology, KIST School, University of Science and Technology (UST), Hwarangno 14-gil 5, Seongbuk-gu, Seoul 02792, Republic of Korea

^c Department of Food Science and Technology, Yeungnam University, 280 Daehak-ro, Gyeongsan-si, Gyeongsangbuk-do 38541, Republic of Korea

* Corresponding author

Byoung Chan Kim, Ph.D.

Tel: + 82 2 958 5877

Fax: + 82 2 958 5805

E-mail address: bchankim@kist.re.kr (B.C. Kim)




Figure S1. Schematic representation of a centrifugation-based partitioning method (CBPM).




Figure S2. The calibration curve of *Cronobacter sakazakii* detection in range of $10^2 - 10^6$ cfu/ml.




Figure S3. Images of seven *Cronobacter* species grown on CESA agar plates.