

Supplementary Materials

Dual-wavelength electrochemiluminescence ratiometry for hydrogen

sulphide detection based on Cd²⁺-doped g-C₃N₄ naonsheets

Jun-Tao Cao,*^a Yi-Zhuo Fu,^a Xiao-Long Fu,^a Shu-Wei Ren,^b and Yan-Ming Liu*^a

^a*College of Chemistry and Chemical Engineering, Xinyang key laboratory of functional nanomaterials for bioanalysis, Xinyang Normal University, Xinyang 464000, China*

^b*Xinyang Central Hospital, Xinyang 464000, China*

*Corresponding author: E-mail: liuym9518@sina.com (Y.-M. Liu) and jtcao11@163.com (J.-T. Cao), Tel & fax: +86-376-6392889.

Table S1. Analytical results of the proposed method for H₂S in human plasma samples.

Serum Samples No.	Found (μM)	RSDs (%) (%, n = 3)
1	0.53	7.6
2	0.46	5.5
3	0.59	5.6
4	0.65	4.8
5	0.61	3.9

Table S2. Recovery tests of H₂S in human serum samples.

Serum samples	Found (μM)	Added (μM)	Total Found (μM)	Recoveries (%)	RSDs (%), n = 3
1	0.46	0.05	0.50	80.0	3.8
		0.50	0.93	94.0	6.9
		5.00	5.48	100.4	4.3
2	0.65	0.05	0.70	100.0	7.4
		0.50	1.18	106.0	3.2
		5.00	5.63	99.6	5.9