

1 Preparative separation and purification of loliolide and
2 epiloliolide from *Ascophyllum nodosum* using amine-
3 based microporous organic polymer for solid phase
4 extraction coupled with macroporous resin and
5 prep-HPLC

6 Hao Chi,^a Xia Qi,^a Xiaohui Wang,^{c,d} Ying Wang,^{c,d} Xihong Han,^{c,d} Jiao Wang^{*b}
7 and Hongwei Wang^{*a}

8

9 Supporting Information

10 Summary

11 This supplementary material includes additional results and information as described in the text
12 of the main article. Including:

13 Contents

14 Page S-2: **Figure S1.** The standard curve of loliolide established using UPLC.

15 Page S-3: **Figure S2.** HRESI-MS of loliolide.

16 Page S-4: **Figure S3.** ¹H NMR of loliolide.

17 Page S-5: **Figure S4.** ¹³C NMR of loliolide.

18 Page S-6: **Figure S5.** NOSEY of loliolide.

19 Page S-7: **Figure S6.** COSY of loliolide.

20 Page S-8: **Figure S7.** HMBC of loliolide.

21 Page S-9: **Figure S8.** HRESI-MS of epiloliolide.

22 Page S-10: **Figure S9.** ¹H NMR of epiloliolide.

23 Page S-11: **Figure S10.** ¹³C NMR of epiloliolide.


24 Page S-12: **Figure S11.** NOSEY of epiloliolide.

25 Page S-13: **Figure S12.** COSY of epiloliolide.

26 Page S-14: **Figure S13.** HMBC of epiloliolide.

27


28 **Figure S1**



29


30

31 Figure S2



32


33 **Figure S3**



34

35


36 **Figure S4**



37

38


39 **Figure S5**



40

41


42 **Figure S6**



43

44


45 **Figure S7**



46

47


48 Figure S8



49

50


51 **Figure S9**




52


53

54 **Figure S10**




55





60 **Figure S13**



61