Electronic Supplementary Material (ESI) for RSC Chemical Biology. This journal is © The Royal Society of Chemistry 2021



Preliminary Full wwPDB X-ray Structure Validation Report (i)

Apr 13, 2021 – 11:57 PM EDT

Deposition ID : $D_{1000253297}$

This is a Preliminary Full wwPDB X-ray Structure Validation Report.

This report is produced by the wwPDB Deposition System during initial deposition but before annotation of the structure.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

	/	
MolProbity /:	:	4.02b-467
Mogul :	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) :	:	1.13
EDS :	:	2.18
buster-report :	:	1.1.7 (2018)
Percentile statistics :	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac :	:	5.8.0158
CCP4 :	:	7.0.044 (Gargrove)
Ideal geometry (proteins) :	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA) :	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) :	:	2.18

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: *X-RAY DIFFRACTION*

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	(# Entries, resolution range(Å))
R_{free}	130704	2092 (3.00-3.00)
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)
RSRZ outliers	127900	1990 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain Length	Quality of chain		
1 /	A 258	77%	23%	_
1	B 258	80%	20%	_
1	C 258	74%	25%	•

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:



	Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density	
Ļ	2	LIG	D	401	-	-	-	X	/
	2	LIG	F,	301	-	-	-		
					/		Y /		
					171				
			/						
			/ .						
			<u> </u>	Y					
			N						
		•							
/									
		\sim							
		\mathbf{Y}							
	X								
	Z'		7		Y	WORLDWIDE			
					1	PROTEIN DATA BANK			

2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 5636 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
1	Δ	257	Total C N O S	0	0	Ο
1	Л	201	1796 1148 $292/342$ 14	0	0	0
1	В	257	Total C N O S		0	Ο
T		201	1826 1150 302 358 16	• 0	0	U
1	С	255	Total C N O S	Q	0	Ο
	200	1773 1114 298 348 13	0	0	0	

• Molecule 2 is a ligand with the chemical component id LIG but its atom names do not match the existing wwPDB Chemical Component Dictionary definition for LIG. Consequently no firm identification of ligand chemistry can be made. Once the structure is annotated then an identification and diagram will be given here.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	Е	1	Total C N O S 96 60 16 17 3	0	0
2	D	1	Total C N O S 63 39 10 11 3	0	0
2	F	1	Total C N O S 82 52 12 15 3	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	70.96Å 84.47Å 134.21Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Bosolution(A)	48.76 - 3.00	Depositor
Resolution (A)	48.76 - 3.00	EDS
% Data completeness	99.7 (48.76-3.00)	Depositor
(in resolution range)	99.7 (48.76-3.00)	EDS
R_{merge}	0.18	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.09 (at 3.01 Å)	Xtriage
Refinement program	PHENIX 1.18.2 3874	Depositor
B B.	0.260 , 0.275	Depositor
$\mathbf{n}, \mathbf{n}_{free}$	0.260 , 0.276	DCC
R_{free} test set	859 reflections $(5.15%)$	wwPDB-VP
Wilson B-factor (Å ²)	102.6	Xtriage
Anisotropy	0.366	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.27 , 70.7	EDS
L-test for twinning ²	$ < L >=0.48, < L^2>=0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	5636	wwPDB-VP
Average B, all atoms $(Å^2)$	106.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.64% of the height of the origin peak. No significant pseudotranslation is detected.

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: LIG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bond	lengths	Bond angles		
	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.26	0/1822	0.54	0/2485	
1	В	0.28	0/1852	0.54	0/2523	
1	С	0.27	0/1799	0.54	0/2454	
All	All	0.27	0/5473	0.54	0/7462	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (1)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1796	0	1678	40	0
1	B	1826	Ø	1687	33	0
1	С	1773	0	1564	44	0
2	D	63	0	0	2	0
2	Е	96	0	0	3	0
2	F	82	0	0	2	0
All	All	5636	0	4929	123	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:147:ILE:HG23	1:A:180:ILE:HD11	1.58	0.85
1:C:137:VAL:HG22	1:C:197:ILE:HG12	1.60	0.81
1:C:9:GLY:HA3	1:C:88:ILE:HD13	1.66	0.76
1:C:147:ILE:HG21	1:C:169:PHE:HD2	1.51	0.74
1:C:38:GLN:NE2	1:C:126:LEU:O	2.22	0.73
1:B:134:SER:H	1:B:230:SER:HB3	1.54	0.72
2:E:301:LIG:N70	2:E:301:LIG:O68	2.18	0.72
1:A:159:VAL:HG22	1:A:206:THR:HG22	1.72	0.71
1:A:143:GLU:O	1:A:147:ILE:HG12	1.93	0.69
1:B:159:VAL:HG22	1:B:206:THR:HG22	1.76	0.68
1:B:19:LEU:HD11	1:B:247:LEU:HD11	1.76	0.67
1:C:1:MET:HB3	1:C:61:ARG:HH12	1.59	0.66
1:A:50:LEU:HD13	1:A:247:LEU:HD12	1.77	0.65
1:C:29:ASP:OD2	1:C:125:GLN:NE2	2.31	0.64
2:D:401:LIG:N29	2:D:401:LIG:O27	2.30	0.64
1:A:23:ILE:HB	1:A:72:LEU:HD12	1.80	0.63
1:C:87:ILE:HB	1:C:104:GLU:HB2	1.81	0.63
1:A:184:GLN:HA	1:A:195:VAL:HG13	1.81	0.62
1:A:5:ARG:HB3	1:A:59:THR:HB	1.81	0.62
1:A:185:THR:HG23	1:A:195:VAL:HG12	1.82	0.62
1:C:5:ARG:HA	1:C:89:THR:HG22	1.81	0.62
1:B:9:GLY:HA3	1:B:88:ILE:HD13	1.83	0.61
1:C:141:SER:HB3	1:C:219:THR:HG23	1.83	0.60
1:B:5:ARG:HG3	1:B:89:THR:HG22	1.85	0.59
1:C:147:ILE:HG21	1:C:169:PHE:CD2	2.35	0.59
1:B:162:CYS:HB2	1:B:203:VAL:HG12	1.84	0.58
1:B:14:LYS:HD3	1:B:220:PRO:HB2	1.85	0.57
2:E:301:LIG:O27	2:E:301:LIG:N32	2.37	0.57
1:A:226:THR:OG1	1:A:238:GLU:HB3	2.04	0.57
1:B:152:SER:HA	1:B:209:LEU:HD13	1.86	0.57
1:B:89:/THR:OG1	I:B:102:VAL:HB	2.05	0.57
1:C:9:GLY:HA3	1:C:88:ILE:CD1	2.35	0.56
1:A:24:ASN:O	1:A:72:LEU:N	2.39	0.56
1:A:68:MET:HB3	1:A:118:LEU:HD11	1.87	0.55
1:C:1:MET:HB3	1:C:61:ARG:NH1	2.21	0.55
1:B:7:VAL:HA	1:B:87:ILE:HG12	1.88	0.54
1:C:61:ARG:NH1	1:C:63:ASP:OD2	2.41	0.54
2:D:401:LIG:O40	2:D:401:LIG:O36	2.25	0.53
1:A:206:THR:OG1	1:A:254:LYS:O	2.27	0.53
1:C:53:ARG:HG2	1:C:244:MET:O	2.08	0.53
	¥ F		ed on next page

All (123) close contacts within the same asymmetric unit are listed below, sorted by their clash/ magnitude.



~ /

Atom-1	Atom-2	Interatomic	Clash
		distance (A)	overlap (A)
1:B:226:THR:OG1	1:B:238:GLU:HB3	2.09	0.53
1:B:37:LEU:HD23	1:B:50:LEU:HB3	1.92	0.52
1:C:14:LYS:NZ	1:C:220:PRO:O	2.36	0.52
1:C:158:VAL:HB	1:C:209:LEU:HD11	1.91	0.52
1:C:38:GLN:HA	1:C:48:VAL:O	2.10	0.51
1:C:74:SER:O	1:C:78:ILE:HG12	2.09	0.51
1:C:89:THR:OG1	1:C:102:VAL:HB	2.11	0.51
1:C:6:LEU:HD11	1:C:244:MET:HE3	1.93	0.51
1:C:226:THR:OG1	1:C:238:GLU:HB3	2,10	0.51
1:A:175:LEU:H	1:A:175:LEU:HD23	1.76 人	0.51
1:C:105:ALA:HB2	1:C:110:LYS:HB3	1.92	0.51
1:A:39:SER:O	1:A:47:LEU:HD12	2.11	0,50
1:B:11:ILE:O	1:B:15:VAL:HG23	2.12	0.50
1:A:82:ALA:HB2	1:A:103:PHE:CD2	2.46	0.50
1:A:228:SER:OG	1:A:236:VAL:HB	2.12	0.50
1:A:99:LEU:HB3	1:A:118:LEU:HD21	1.93	0.49
1:B:47:LEU:HD23	1:B:250:TYR:CD2	2.46	0.49
1:B:158:VAL:HB	1:B:209:LEU;/HD21	1.95	0.49
1:B:159:VAL:HB	1:B:170:SER:OG	2.13	0.49
1:A:47:LEU:HB3	1:A:250:TYR:HB2	1.94	0.49
1:A:205:LEU:HD21	1:A:231:ALA:HA	1.95	0.49
1:C:57:PHE:HE1	1:C:244:MET:HE3	1,77	0.49
1:B:137:VAL:HG22	1:B:197:ILE:HG12	1.94	0.48
1:B:9:GLY:HA3	1:B:88:ILE:CD1	2.44	0.48
1:B:30:ILE:HG23	1:B:35:VAL:HG12	1.96	0.48
1:B:175:LEU:H	1:B:175:LEU:HD23	1.79	0.48
1:A:89:THR:OG1	1:A:102:VAL:HB	2.14	0.47
1:A:1:MET:HB3	1:A:63:ASP:OD2	2.14	0.47
1:C:57:PHE:CD2	1:C:60:TYR:HD1	2.33	0.47
1:C:236:VAL;HG13	1:C:250:TYR:CE1	2.49	0.47
1:A:147:ILÉ:HD12	1:A:180:ILE:CD1	2.44	0.47
1:B:189:ASP:O 📥	1:B:193:GLU:HA	2.15	0.47
1:A:227:LEU:HD13	1:A:237:VAL:HG22	1.96	0.47
1:C:7:VAL:HG23	1:C:8:GLN:H	1.80	0.47
1:B:138:LYS:HB3	1:B:196:THR:OG1	2.16	0.46
1:C:37:LEU:O	1:C:49:GLN:HA	2.15	0.46
1:B:38:GLN:HA	/1:B:48:VAL:O	2.16	0.46
1:A:66:LEU:HD23	1:A:67:ALA:N	2.31	0.46
1:C:88:ILE:HG13	1:C:103:PHE:CD1	2.51	0.46
1:A:156:ASP:OD1	1:A:156:ASP:N	2.49	0.46
1.C.129.PBO.HD2	2:F:301:LIG:C92	2.46	0.46

..... α tia d fa



	ious puye	International	Clash	
Atom-1	Atom-2	distance $(Å)$	overlap (Å)	
1.C.88.II F.HC13	1.C.103.PHF.HD1			
2·E·301·LIG·N56	2·E·301·LIC·O40	2.49	0.40	
1.A.66.LEU.HD11	1.A.96.ALA.HB2	1.98	0.45	- V
$\frac{1 \cdot A \cdot 4 \cdot A \downarrow A \cdot \Omega}{1 \cdot A \cdot 4 \cdot A \downarrow A \cdot \Omega}$		2.16	0.45	-
1.C.28.TRP.O	1.C.67.ALA.HA	2.10	0.45	- /
1.C.20.110	1.C.94.ASP.OD1	2.10	0.45	
1.0.1.WEF.R	1.0.94.MOI .0D1	1.98	0.44	
1.C.215.PHE.CE2	1.C.249.TVR.HB3	2 52	0.11	1
1.C.16.LEU.HD13	1.C.19.LEU.HD11	2.02	0.44	-
1.C.6.LEU.HD13	1.C.12.LEU.HD23	1 99	0.44	_
1:C:36:ASN:OD1	1.C.37.LEU.N	2.51	0.11	-
1.0.30.ADA.HB2	1.0.07.1110.11 1.A.103.PHE:CE2	2.54	0.43	-
$\frac{1.1.02.11 \pm 1.11D2}{1.4.184.GLN.OE1}$	1.A.184.GLN·N	2.01	0.43	-
1.R.143.GLU.O	1.B.147.ILE.HG13	2.40	0.43	-
1.B.119.010.0	1.B.44.HIS.N	2.51	0.19	-
1.D.41.MS1.OD1 1.C.7.VAL.HG23	1.D.44.III5.N	2.01	0.43	-
1.0.70.VAL:HG11	1.0.0.0EIV.IV	2.39	0.49	-
1.C.33.SEB.O	1:C:54:SEB:N	2.15	0.12	-
1.0.55.5ER.0	1.C.171.ALA.HB3	2.00	0.42	-
1.0.101.110.110222	1.A.205.LEU.HD22	2.01	0.12	-
1.A.147.ILE.HD12	1:A·180·ILE·HD12	2.10	0.12	-
1.C·242·ALA·O	$1 \cdot C \cdot 244 \cdot MET \cdot N$	2.50	0.12	-
1.C.50.LEU.HD13	$1 \cdot C \cdot 247 \cdot LEU \cdot HD12$	2.00	0.12	-
1:A:112:SEB:HB3	1.A.114.TYB.HE1	1.85	0.42	_
1·B·5·ARG·HG3	1·B·89·THB·CG2	2.50	0.42	_
1:B:10:SEB:0	1:B:14:LYS:HG3	2.20	0.42	-
1:A:40:MET:HE2	1:A:44:HIS:HA	2.02	0.41	-
1:A:235:LEU:O	1:A:250:TYB:HA	2.20	0.41	-
1:B:50:LEU:HD13	1:B:247:LEU:HD12	2.02	0.41	-
1:B:236:VAL/HG22	1:B:250:TYR:CD1	2.55	0.41	-
1:A:158:VAL:HB	1:A:209:LEU:HD21	2.02	0.41	-
1:B:70:VAL:HG11	1:B:75:MET:SD	2.61	0.41	-
1:C:103:PHE:HB2	1:C:112:SER:HB2	2.02	0.41	-
1:C:182:LEU:HD23	1:C:182:LEU:HA	1.95	0.41	_
1:A:5:ARG:HA	1:A:89:THR:HG22	2.01	0.41	-
1:B:37:LEU:CD2	1:B:50:LEU:HB3	2.50	0.41	-
1:B:169:PHE:HE1	1;B:182:LEU:HD13	1.86	0.41	-
1:C:185:THR:H	/1:C:194:ALA:HB1	1.87	0.40	1
2:F:301:LIG:O46	2:F:301:LIG:N18	2.54	0.40	-
1:A:242:ALA:O	1:A:244:MET:N	2.52	0.40	1
1:A:12:LEU:HD23	1:A:88:ILE:HG21	2.01	0.40	1

Continued from previous page...



· · · · ·	
tinued from previous p	age
inaca from previous p	uy

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)	
1:B:219:THR:N	1:B:220:PRO:HD2	2.36	0.40	

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percer	ntiles
1	А	255/258~(99%)	242 (95%)	12 (5%)	1(0%)	34	72
1	В	255/258~(99%)	241 (94%)	14 (6%)	0	100	100
1	С	253/258 (98%)/	242 (96%)	9 (4%)	2(1%)	19	57
All	All	763/774 (99%)	725 (95%)	35 (5%)	3~(0%)	34	72

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	А	189	ASP
1	С	184	GLN
1	C	194	ALA

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol Chain Analysed		Rotameric	Outliers	Percentiles		
1	A	/169/226 (75%)	169 (100%)	0	100	100
	/			Continuel		



f									
Mol	Chain	Analysed	sed Rotameric C		Perce	entiles			
1	В	179/226~(79%)	179 (100%)	0	100	100			
1	С	158/226 (70%)	158 (100%)	0	100	100			
All	All	506/678~(75%)	506 (100%)	0	100	100			

Continued from previous page...

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 3 ligands modelled in this entry, 3 could not be matched to an existing wwPDB Chemical Component Dictionary definition at this stage - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	$\#RSRZ{>}2$	$OWAB(Å^2)$	Q<0.9
1	А	257/258~(99%)	-0.15	3 (1%) 79 54	71, 95, 159, 204	0
1	В	257/258~(99%)	-0.19	2(0%) 86 65	71, 95, 146, 195	0
1	С	255/258~(98%)	-0.07	6 (2%) 59 30	87, 110, 157, 198	0
All	All	769/774~(99%)	-0.14	11 (1%) 75 49	71, 101, 154, 204	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	С	107	ASN	3.8
1	С	22	LEU	3.0
1	А	124	GLU	2.9
1	В	28	TRÝ	2.8
1	С	185	THR	2.6
1	С	207	PHE	2.5
1	С	23 /	ILE	2.3
1	А	28	TRP	2.3
1	А	123	VAL	2.2
1	С	28	TRP	2.1
1	B	68	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.



6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(A^2)$	Q<0.9
2	LIG	D	401	63/?	0.65	0.44	115,153,173,174	0 /
2	LIG	F	301	82/?	0.73	0.42	109,129,140,143	0⁄
2	LIG	Е	301	96/?	0.77	0.29	$92,\!122,\!139,\!147$	/0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







6.5 Other polymers (i)

There are no such residues in this entry.

