

Supporting Information

Molecular adduct of amantadine ferulate presents a pathway for slowing *in vitro/vivo* releases and raising synergistic antiviral effects via dual optimization salification strategy

Yuan-Yuan Niu^{‡a}, Ling-Yang Wang^{‡a}, Yue-Ming Yu,^a Yan-Tuan Li^{a,b,*}, Zhi-Yong Wu^{a,*}, Cui-Wei Yan^{a,*}

^a*School of Medicine and Pharmacy and College of Marine Life Science, Ocean University of China, Qingdao, Shandong 266003, PR China*

^b*Laboratory for Marine Drugs and Bioproducts, Qingdao National Laboratory for Marine Science and Technology, 266003, PR China*

*Correspondence to: Yan-Tuan Li, Zhi-Yong Wu and Cui-Wei Yan.

E-mail addresses: yantuanli@ouc.edu.cn (Yan-Tuan Li), wuzy@ouc.edu.cn (Zhi-Yong Wu), cuiweiyan@ouc.edu.cn (Cui-Wei Yan).

This ESI for *CrystEngComm*, 2021, **23**, 4389-4401, DOI: 10.1039/D1CE00382H, originally published on 4th June 2021, was updated on 8th June 2023.

Contents

Figure S1. FT-IR spectra of FD, AIN, and the molecular salt.

Figure S2. XRPD comparison of simulated pattern for the molecular salt and the residual solids after the powder dissolution measurements.




Figure S1. FT-IR spectra of FD, AIN, and the molecular salt.




Figure S2. XRPD comparison of simulated pattern for the molecular salt and the residual solids after the powder dissolution measurements.