## **Electronic Supplementary Information (ESI)**

## Zinc delivery system constructed by food-borne nanoparticles derived from *Undaria pinnatifida*

Fengzhi Qiao,<sup>a,b,c</sup> Xiaoting Yu,<sup>a,b,c</sup> Shanshan Tie,<sup>a,b,c</sup> Yannan Chen, <sup>a,b,c</sup> Shuai Hou,<sup>\*a,b,c</sup> and

Mingqian Tan,\*a,b,c

<sup>a</sup> Academy of Food Interdisciplinary Science, Food Science and Technology, Dalian Polytechnic University, Qinggongyuan1, Ganjingzi District, Dalian 116034, Liaoning, China

<sup>b</sup> National Engineering Research Center of Seafood, Dalian Polytechnic University, Dalian 116034, Liaoning, China

<sup>c</sup> Collaborative Innovation Center of Seafood Deep Processing, Dalian Polytechnic University Dalian, 116034, China

\*To whom correspondence should be addressed.

Shuai Hou, E-mail: houshuai009@126.com; ORCID: 0000-0001-5274-5264.

Mingqian Tan, E-mail: mqtan@dlpu.edu.cn; ORCID: 0000-0002-7535-0035.




Figure. S1 MALDI-TOF-MS spectrum of UPFNs from Undaria pinnatifida.

	C (%)	N (%)	O (%)	S (%)	Zn (%)
UPFNs	69.04	4.28	25.98	0.71	
UPFN-Zn <sup>2+</sup>	37.05	1.79	49.18	6.33	5.65

Table S1 Elemental composition of UPFNs and UPFN- $Zn^{2+}$ 

"---" The element is not present in the sample