

**Phycocyanin Ameliorates Mouse Colitis via Phycocyanobilin-Dependent
Antioxidant and Anti-Inflammatory Protection of Intestinal Epithelial Barrier**

Wei Guo,^{a, b} Mingyong Zeng,^b Suqin Zhu,^c Shiyang Li,^b Yilin Qian,^b and Haohao
Wu^{*b}

^a School of Pharmacy, Binzhou Medical University, 346 Guanhai Road, Yantai,
Shandong, 264003, China

^b College of Food Science and Engineering, Ocean University of China, 5 Yushan
Road, Qingdao, Shandong Province, 266003, China

^c Institute of Nutrition and Health, School of Public Health, Qingdao University,
Qingdao 266021, China

* The author to whom correspondence should be addressed; E-mail:

wuhaohao@ouc.edu.cn; Tel. & Fax: +86-532-8203-2400.

Table S1 The criteria for DAI

Score	Trait of stool	Rectal bleeding	Weight loss
0	Normal stools	Negative	None
1	Soft stools	Negative	1-5%
2	Soft stools	Positive	6-10%
3	Very soft	Visible in stool	11-15%
4	Watery stool	Gross bleeding	> 15%

Figure S1 (a) Primary mass spectra of PCB (3Z); (e) secondary mass spectra of PCB (3Z).

