

Table S1. Primer sequences used for real-time RT-PCR

Gene	Forward primer (5' → 3')	Reverse primer (5' → 3')
<i>P16</i>	CGCAGGTTCTTGGTCACTGT	TGTTACGAAAGCCAGAGCG
<i>Foxl2</i>	ACAACACCGGAGAAACCAGAC	CGTAGAACGGGAACCTGGCTA
<i>Gdf9</i>	CAAGTACAGCCCCCTGAGTG	CACTCTAAGGCCACAGCAT
<i>Bmp15</i>	TGCTTTGGAGATTAGGAGAGAGTG	CCCTCAGGGATTAAAGTAGAGC
<i>Cyp11a1</i>	AGGTCCTTCAATGAGATCCCTT	TCCCTGTAAATGGGGCCATAC
<i>Hsd3b1</i>	AGCTCTGGACAAAGTATTCCGA	GCCTCCAATAGGTTCTGGGT
<i>Star</i>	TGTTAAGGACTGCCACCACATCT	TGTCCTTGGCTGAAGGTGAACAGA
<i>Tnfa</i>	CCTGTAGCCCACGTCGTAG	GGGAGTAGACAAGGTACAACCC
<i>Il6ra</i>	GCCACCGTTACCCTGATTTG	TCCTGTGGTAGTCCATTCTCTG
<i>Il10rb</i>	GCCACCGTTACCCTGATTTG	TCCTGTGGTAGTCCATTCTCTG
<i>Il2</i>	TGAGCAGGATGGAGAATTACAGG	GTCCAAGTTCATCTTCTAGGCAC
<i>Tgfb1</i>	CCACCTGCAAGACCATCGAC	CTGGCGAGCCTTAGTTTGGAC
<i>Lc3</i>	CATGAGCGAGTTGGTCAAGA	TTGACTCAGAAGCCGAAGGT
<i>Gapdh</i>	AACTGAGGACCAGGTTGTCTC	TACTCCTTGGAGGCCATGTAG