Supplementary Information

Elucidating the mechanisms of enhanced lignin bioconversion by an alkali sterilization strategy

Zhi-Min Zhao,^{a,b} Shuyang Zhang,^b Xianzhi Meng,^b Yunqiao Pu,^c Zhi-Hua Liu,^d William K. Ledford,^e S. Michael Kilbey II,^{b,e} Bing-Zhi Li^d and Arthur J. Ragauskas*^{b,c,f}

^a School of Ecology and Environment, Inner Mongolia Key Laboratory of Environmental Pollution Control & Wastes Reuse, Inner Mongolia University, Hohhot, 010021, China

^b Department of Chemical & Biomolecular Engineering, University of Tennessee Knoxville, Knoxville, TN 37996, USA

^c Joint Institute of Biological Science, Biosciences Division, Oak Ridge National Laboratory (ORNL), Oak Ridge, TN 37831, USA

^d Frontiers Science Center for Synthetic Biology and Key Laboratory of Systems Bioengineering (Ministry of Education), School of Chemical Engineering and Technology, Tianjin University, Tianjin, 300072, China

^e Department of Chemistry, University of Tennessee Knoxville, Knoxville, TN 37996, USA

^f Department of Forestry, Wildlife, and Fisheries, Center for Renewable Carbon, University of Tennessee Institute of Agriculture, Knoxville, TN 37996, USA

* Corresponding author: Arthur J. Ragauskas, E-mail address: <u>aragausk@utk.edu</u>




Fig. S1 Examination of the aseptic effect of alkaline sterilized lignin (ASL) and nonsterilized lignin (NSL) mediums by spread plate method. These experiments were conducted with two replicates.




Fig. S2 Comparison of lignin medium color under natural light (a) and light transmittance in front of a 600-W fluorescent lamp strip (b). ASL: alkaline sterilized lignin, ATSL: alkaline and thermally sterilized lignin, NSL: non-sterilized lignin, TSL: thermally sterilized lignin.