Electronic Supplementary Material (ESI) for Materials Advances. This journal is © The Royal Society of Chemistry 2021

Electronic Supplementary Information (ESI)

Ruthenium Inserted Hydrotalcite (Ru-HT) Heterogeneous Catalyst for the Kinetic Studies of Selective Hydrogenation of Carbon Dioxide to Formic Acid

Minaxi S. Maru^{a,c,d*}, Sanwala Ram^{a,b}, Noor-ul H. Khan^{a,b}, Ram S. Shukla^{a,b*}

^aInorganic Materials and Catalysis Division, Council of Scientific and Industrial Research (CSIR), Central Salt and Marine Chemicals Research Institute, G. B. Marg, Bhavnagar, 364002, Gujarat, India

^bAcademy of Scientific and Innovative Research (AcSIR), Council of Scientific and Industrial Research (CSIR), Central Salt and Marine Chemicals Research Institute, G. B. Marg, Bhavnagar, 364002, Gujarat, India

^cInorganic Chemistry Division, Department of Chemistry, Saurashtra University, Rajkot, 360005, Gujarat, India

^dDepartment of Chemistry, Indian Institute of Teachers Education (IITE), Gandhinagar, 382016, Gujarat, India




Figure S1. (a) HPLC graph of standard formic acid solution. **(b)** HPLC graph of formic acid obtained from the hydrogenation of CO₂.




Figure S2. Field Emission Scanning Electron Microscope (FE-SEM) images of Ru-HT (**a-d**) and RRu-HT (**e-h**).





Figure S3. Scanning Electron Microscope-EDX graph of Ru-HT showing the presence of Ru^{II} ion.






Figure S5. XRD spectra of HT, Ru-HT and RRu-HT (Recycled Ru-HT).




Figure S6. The X-ray photoelectron spectroscopy (XPS) of Ru^{II} 3d species observed in Ru-HT and RRu-HT.