Electronic Supplementary Material (ESI) for RSC Medicinal Chemistry. This journal is © The Royal Society of Chemistry 2021

Supplementary Information

SF₅ and SCF₃-substituted tetrahydroquinoline compounds as potent bactericidal agents against multidrug resistant persister Gram-positive bacteria

Kenneth I. Onyedibe^{1,3}, Neetu Dayal¹, Herman O. Sintim^{* 1,2,3},


Department of Chemistry, Purdue University, 560 Oval Drive, West Lafayette, Indiana 47907, United States


² Center for Drug Discovery, Purdue University, 720 Clinic Drive, West Lafayette, Indiana 47907, United States


³Purdue Institute of Inflammation, Immunology, and Infectious Disease, West Lafayette, IN, 47907, USA


^{*}Corresponding author. Email: hsintim@purdue.edu


¹H Spectra for Compound 1

















-10 -20 -30 -40 -50 -60 -70 -80 -90 -100 -110 -120 -130 -140 -150 -160 -170 -180 -190 -200 -210 -220 f1 (ppm)


¹H Spectra for Compound 4 ndayal-SF5pych.1.fid

,ΝH





12.0 11.5 11.0 10.5 10.0 9.5 9.0 8.5 8.0 7.5 7.0 6.5 6.0 5.5 5.0 4.5 4.0 3.5 3.0 2.5 2.0 1.5 1.0 0.5 0.0 -0.5 -1.0 -1. f1 (ppm)






12.0 11.5 11.0 10.5 10.0 9.5 9.0 8.5 8.0 7.5 7.0 6.5 6.0 5.5 5.0 4.5 4.0 3.5 3.0 2.5 2.0 1.5 1.0 0.5 0.0 -0.5 -1.0 -1. f1 (ppm)







¹³H Spectra for Compound 8

