

Supplementary Materials

Synthesis of a light-harvesting ruthenium porphyrin complex substituted with BODIPY units. Implications for visible light-promoted catalytic oxidations

Jonathan Malone, Seth Klaine, Christian Alcantar, Fox Bratcher, and Rui Zhang*

*Department of Chemistry, Western Kentucky University, 1906 College Heights Blvd
#11079, Bowling Green, KY 42101-1079*

Contents

Characterization spectra S1-S7 for compounds 1-3	Pages 2-5
---	-----------





Figure S1. (A) UV-Vis spectrum of the BODIPY-appended benzaldehyde (**2**) in CH_2Cl_2 and (B) ¹H-NMR spectrum of **2** in CDCl_3 .




Figure S2. ^1H -NMR spectrum of the light-harvesting porphyrin (**3**) in CDCl_3




Figure S3. ^{13}C -NMR spectrum of the light-harvesting porphyrin (**3**) in CDCl_3 .




Figure S4. ESI MS of the light-harvesting porphyrin (**3**).




Figure S5. ¹H-NMR spectrum of [Ru^{II}(L-Por)(CO)] (**1**) in CDCl₃




Figure S6. ¹³C-NMR spectrum of [Ru^{II}(L-Por)(CO)] (**1**).




Figure S7. IR spectrum of Ru^{II}[(L-Por)](CO) (**1**).