

Hematin-conjugated gelatin performing an effectively catalytic function for preparing biological hydrogel

Tan Phuoc Ton,^{a,b,†} Van Toan Nguyen,^{a,c,†} Phuong Doan,^b Dinh Trung Nguyen,^{a,b} Thi Phuong Nguyen,^d Chan Khon Huynh,^e Thi Cam Quyen Ngo,^{a,c} Le Hang Dang,^{*a,b} and Ngoc Quyen Tran^{*ab}

^{a.} Graduate University of Science and Technology, Vietnam Academy of Science and Technology, HCMC, Vietnam;

^{b.} Institute of Applied Materials Science Vietnam Academy of Science and Technology, HCMC, Vietnam;

^{c.} Institute of Fundamental and Applied Sciences, Duy Tan University, HCMC 700000, Vietnam;

^{d.} Faculty of Chemical Technology, HCMC University of Food Industry, HCMC, Vietnam;

^{e.} Department of Biomedical Engineering, International University, Vietnam National University - HCMC, HCMC, Vietnam;

^{f.} NTT Hi-Tech Institute, Nguyen Tat Thanh University, HCMC, Vietnam.

[†] First two authors contributed equally to this study.

Supporting materials




Figure S1: UV/Vis spectra of hematin (He) and Ge-He in Tris buffer (pH 8.5), PBS buffer (pH 7.4) and DMSO, [He] = 10ppm, [Ge-He] = 25ppm.