

Supplementary Information

A Disposable Electrochemical Sensor Based on Iron Molybdate for Analysis of Dopamine in Biological Samples

Elumalai Ashok Kumar^{a,b}, Tse-Wei Chen^{c,d,e}, Shen-Ming Chen^{c*}, Tzyy-Jiann Wang^a, Allen Joseph Anthuvan^f, Suliman Yousef AlOmar^g, Naushad Ahmad^h, Yu-Hsu Chang^b,

^aDepartment of Electro-Optical Engineering, National Taipei University of Technology, Taipei 106, Taiwan.

^bDepartment of Materials and Mineral Resources Engineering, Institute of Mineral Resources Engineering, National Taipei University of Technology, Taipei 106, Taiwan.

^cDepartment of Chemical Engineering and Biotechnology, National Taipei University of Technology, Taipei 106, Taiwan.

^dResearch and Development Center for Smart Textile Technology, National Taipei University of Technology, Taipei 106, Taiwan.

^eDepartment of Materials, Imperial College London, London, SW7 2AZ, United Kingdom.

^fDRDO-Centre for Life Sciences, Bharathiar University, Coimbatore 641 046, India.

^gDoping Research Chair, Zoology Department, College of Science, King Saud University, Kingdom of Saudi Arabia.

^hChemistry Department, College of Science, King Saud University, Kingdom of Saudi Arabia.

Corresponding author

*E-mail: smchen78@ms15.hinet.net

Tel: +886 2270 17147, Fax: +886 2270 25238.




Fig. S1 CV response for concentration (mg) of the $\text{Fe}_2(\text{MoO}_4)_3/\text{SPCE}$.




Fig. S2 CV response for loading amount (μL) of the $\text{Fe}_2(\text{MoO}_4)_3/\text{SPCE}$.