Electronic Supplementary Material (ESI) for New Journal of Chemistry. This journal is © The Royal Society of Chemistry and the Centre National de la Recherche Scientifique 2021

Aggregation-induced emissions in mesogenic BF₂ complexes

Yi-Hong Cai^a, Hsin-Yun Chih^a, Gene-Hsiang Lee^b, Chung K. Lai^a*

Fax: (+8862)-03-4277664; (+886)-03-4259207 ext. 65904; e-mail: cklai@cc.ncu.edu.tw

^a Department of Chemistry, National Central University, Chung-Li 32011, Taiwan, ROC

^b Instrumentation Center, National Taiwan University, Taipei 10660, Taiwan, ROC

Table of Contents

| 1. Т | The ¹ H, | ^{13}C ar | nd ¹⁹ F I | NMR | spectra | of c | ompounds | 1a, | 1b, | 1a-BF ₂ | and | 1b-BF | 22 |
|------|---------------------|-------------|----------------------|-----|---------|------|----------|-----|-----|--------------------|-----|-------|----|
|------|---------------------|-------------|----------------------|-----|---------|------|----------|-----|-----|--------------------|-----|-------|----|

2. The DSC thermograph of complex $1a-BF_2$ (n = 8)19





Fig. S1 The ¹H spectrum of compound 1a (n = 8).



Fig. S2 The ¹³C NMR spectrum of compound 1a (n = 8).



Fig. S3 The ¹H spectrum of compound 1b (n = 8).



Fig. S4 The ¹³C NMR spectrum of compound 1b (n = 8).



Fig. S5 The ¹H spectrum of compound 1a-BF₂ (n = 8).



Fig. S6 The ¹³C NMR spectrum of compound 1a-BF₂ (n = 8).



Fig. S7 The ¹⁹F-NMR spectrum of compound 1a-BF₂ (n = 8).



Fig. S8 The ¹H spectrum of compound 1a-BF₂ (n = 10).



Fig. S9 The ¹³C NMR spectrum of compound 1a-BF₂ (n = 10).



Fig. S10 The ¹⁹F-NMR spectrum of compound $1a-BF_2$ (n = 10).



Fig. S11 The ¹H spectrum of compound 1a-BF₂ (n = 12)



Fig. S12 The ¹³C NMR spectrum of compound 1a-BF₂ (n = 12).



Fig. S13 The ¹⁹F-NMR spectrum of compound 1a-BF₂ (n = 12).



Fig. S14 The ¹H spectrum of compound 1a-BF₂ (n = 14).



Fig. S15 The ¹³C NMR spectrum of compound 1a-BF₂ (n = 14).



Fig. S16 The ¹⁹F-NMR spectrum of compound $1a-BF_2$ (n = 14).



Fig. S17 The ¹H spectrum of compound 1a-BF₂ (n = 16).







Fig. S19 The ¹⁹F-NMR spectrum of compound 1a-BF₂ (n = 16).





Fig. S21 The ¹³C NMR spectrum of compound 1b-BF₂ (n = 8).







Fig. S23 The ¹H spectrum of compound 1b-BF₂ (n = 10).







Fig. S25 The ¹⁹F-NMR spectrum of compound $1b-BF_2$ (n = 10).





Fig. S27 The ¹³C NMR spectrum of compound 1b-BF₂ (n = 12).



Fig. S28 The ¹⁹F-NMR spectrum of compound $1b-BF_2$ (n = 12).



Fig. S29 The ¹H spectrum of compound 1b-BF₂ (n = 14).



Fig. S31 The ¹⁹F-NMR spectrum of compound 1b-BF₂ (n = 14).





Fig. S33 The ¹³C NMR spectrum of compound 1b-BF₂ (n = 16).



Fig. S34 The ¹⁹F-NMR spectrum of compound $1b-BF_2$ (n = 16).

2. The DSC Thermographs of Compounds $1a-BF_2$ (n = 8)



Fig. S35 DSC thermograph of compound 1a-BF₂ (n = 8).