

Electronic Supplementary Material for:

Tunable Cu-M bimetal catalysts enable syngas electrosynthesis from carbon dioxide

Bo Xiong, Jing Liu*, Yingju Yang*, Junyan Ding, Zhixuan Hua

State Key Laboratory of Coal Combustion, School of Energy and Power Engineering, Huazhong University of Science and Technology, Wuhan 430074, China

Corresponding Author:

* Tel: +86 27 87545526; fax: +86 27 87545526;

E-mail address: liujing27@mail.hust.edu.cn;

yangyingju@hust.edu.cn.

Content includes: one.

Fig. S1 The dissolution potential of Cu-M catalysts at 0 V.

Fig. S2 (a) Schematic illustration of the structural model. a denotes Cu sites. b denotes M (M = Fe, Ru, Os, Rh, Ir, Ni, Pd, Pt, Cr, Mn, Mo, and W) sites. (b) The atomic structure of Cu-M catalysts.

Fig. S3 Optimized structures of CO* over the Cu-M catalysts.

Fig. S4 Optimized structures of H* over the Cu-M catalysts.




Fig. S1 The dissolution potential of Cu-M catalysts at 0 V.




Fig. S2 (a) Schematic illustration of the structural model. a denote Cu sites. b denote M ($M = \text{Fe, Ru, Os, Rh, Ir, Ni, Pd, Pt, Cr, Mn, Mo, and W}$) sites. (b) The atomic structure of Cu-M catalysts.




Fig. S3 Optimized structures of CO* over the Cu-M catalysts.




Fig. S4 Optimized structures of H^* over the Cu-M catalysts.