

Dual Quinoline-Hybrid Compounds with Antimalarial Activity Against *Plasmodium falciparum* Parasites

Geraldo José da Silva Neto,¹ Leandro Rocha Silva,² Rafael Jorge Melo de Omena,¹ Anna Caroline Campos Aguiar,³ Yasmin Annunziato,³ Bárbara Rosseto,³ Marcos Gazarine,³ Luana Heimfarth,⁴ Lucindo José Quintans-Júnior,⁴ Edeildo Ferreira da Silva-Júnior,^{2,*} and Mario Roberto Meneghetti^{1,*}

¹ Group of Catalysis and Chemical Reactivity, Institute of Chemistry and Biotechnology, Federal University of Alagoas, Av. Lourival Melo Mota, 57072-900, Maceió, Brazil.

² Institute of Chemistry and Biotechnology, Federal University of Alagoas, Av. Lourival Melo Mota, 57072-900, Maceió, Brazil.

³ Department of Bioscience, Federal University of São Paulo, Central Building - Silva Jardim Street, 136 - Vila Matias, 11015-020, Santos, Brazil.

⁴ Department of Physiology, Federal University of Sergipe, São Cristóvão, Sergipe, Brazil


*Corresponding authors: Meneghetti, MR. Tel.: +55-82-3214-1744; email: mrm@qui.ufal.br; and da Silva-Júnior, EF. Tel.: +55-87-99610-8311; email: edeildo.junior@iqb.ufal.br.

Supplementary Material

Summary

Figure 1S. ^1H NMR (δ , $\text{CD}_3\text{OD}-d_4$, 400 MHz) of GIQ	2
Figure 2S. ^{13}C NMR (δ , $\text{CH}_3\text{OD}-d_4$, 100 MHz) of GIQ	3
Figure 3S. FTIR of compound GIQ	3
Figure 4S. ^1H NMR (δ , $\text{DMSO}-d_6$, 600 MHz) and expansion of TOQ	4
Figure 5S. ^{13}C NMR (δ , $\text{DMSO}-d_6$, 100 MHz) of TOQ	5
Figure 6S. FTIR of TOQ	5
Figure 7S. ^1H NMR (δ , $\text{DMSO}-d_6$, 600 MHz) and expansion of DEQ	6
Figure 8S. ^{13}C NMR (δ , $\text{DMSO}-d_6$, 100 MHz) and expansion of DEQ	7
Figure 9S. FTIR of DEQ	9
Figure 10S. ^1H NMR (δ , $\text{CD}_3\text{OD}-d_4$, 400 MHz) of CEQ	10
Figure 11S. ^{13}C NMR (δ , $\text{CD}_3\text{OD}-d_4$, 100 MHz) of CEQ	11
Figure 12S. FTIR of CEQ	11
Figure 13S. ^1H NMR (δ , CDCl_3 , 400 MHz) and expansion of IBUCEQ	12
Figure 14S. ^{13}C NMR (δ , CDCl_3 , 100 MHz) of IBUCEQ	13
Figure 15S. FTIR of IBUCEQ	14

GIQ



Chemical Formula: C₁₁H₉ClN₂O₂
Exact Mass: 236,04

Figure 1S. ¹H NMR (δ , CD₃OD-*d*4, 400 MHz) of **GIQ**.




Figure 2S. ^{13}C NMR (δ , $\text{CH}_3\text{OD-}d_4$, 100 MHz) of **GIQ**.






Figure 3S. FTIR of compound **GIQ**.



TOQ



Chemical Formula: C₂₀H₁₆ClN₃O₂

Exact Mass: 365,09

Figure 4S. ¹H NMR (δ , DMSO-*d*6, 600 MHz) and expansion of TOQ.





Figure 5S. ^{13}C NMR (δ , DMSO-*d*6, 100 MHz) of TOQ.







Figure 6S. FTIR of TOQ.





Chemical Formula: C₂₁H₂₉ClN₂O₂
Exact Mass: 376,19

Figure 7S. ^1H NMR (δ , DMSO-*d*6, 600 MHz) and expansion of **DEQ**.







Figure 8S. ^{13}C NMR (δ , DMSO-*d*6, 100 MHz) and expansion of **DEQ**.









Figure 9S. FTIR of **DEQ**.



CEQ



Chemical Formula: C₁₁H₁₂ClN₃
Exact Mass: 221,07

Figure 10S. ¹H NMR (δ , CD₃OD-*d*4, 400 MHz) of **CEQ**.




Figure 11S. ^{13}C NMR (δ , $\text{CD}_3\text{OD-}d_4$, 100 MHz) of **CEQ**.






Figure 12S. FTIR of **CEQ**.






Chemical Formula: C₂₄H₂₈ClN₃O

Exact Mass: 409,19

Figure 13S. ^1H NMR (δ , CDCl_3 , 400 MHz) and expansion of **IBUCEQ**.






Figure 14S. ^{13}C NMR (δ , CDCl_3 , 100 MHz) of **IBUCEQ**.




Figure 15S. FTIR of IBUCEQ.

