Supporting Information for

Specific capture and intact release of breast cancer cells using twin-layer vein-shaped microchip with self-assembled surface

Yixing Gou^{a,b}, Zhuyuan Chen^c, Changku Sun^a, Peng Wang^a, Zheng You^b, Yaxiaer Yalikun^{d,e}, Yo Tanaka^d, and Dahai Ren^{*b}

^a State Key Laboratory of Precision Measurement Technology and Instruments, Tianjin University, Tianjin 300072, China

^b State Key Laboratory of Precision Measurement Technology and Instruments, Department of Precision Instrument, Tsinghua University, Beijing 100084, China

^c Department of Basic Sciences, Tsinghua University School of Medicine, Beijing 100084, China

^d Center for Biosystems Dynamics Research (BDR), RIKEN, 1-3 Yamadaoka, Suita, Osaka 565-0871, Japan

^e Division of Materials Science, Nara Institute of Science and Technology, 8916-5 Takayamacho, Ikoma, Nara 630-0192, Japan

Corresponding author: *rendh@tsinghua.edu.cn

Table of Contents

Figure S1: The 3D topography of the chip in a single unit, the colors represent the height in
Z-axis
Figure S2: The geometric size of vein-shaped microchip
Figure S3: SEM image of AuNPs
Figure S4: UV - Visible absorption spectrum of AuNPs. The maximum absorption
wavelength was 514-520 nm, and the maximum absorbance was 0.88-1
Figure S5: Distribution of the flow field in Y-Z plane of the single-layer microchip S4
Figure S6: The distributions of net flow field, flow field in Y-axis and Z-axis different cross
sections of the microchip under different cross sections of microchip in the same flow
throughput (a) the heights of bottom and top layer were equal; (b) the height of bottom layer
was twice of the top layer; (c) the height of bottom layer was half of the top layer S5
Figure S7: Normalized fluorescence intensity reflecting the FITC-labeled Goat anti-Mouse
IgG antibodies immobilization on surface without modification, AuNP-functionalized surface,
and self-assembled surface
Figure S8: Characterization of the aggregation of AuNPs
Figure S9: Different state of cells captured on the self-assembled surface. (a) single cell; (b)
multicellular cluster two cells; (c) multicellular cluster three cells
Figure S10: The proportion of apoptotic/live cells after being cultured 0 min, 20 min, 40 min,
and 60 min under the incubate time of 8min, 12min, 16min, and 20min



Figure S1: The 3D topography of the chip in a single unit, the colors represent the height in Z-axis.



Figure S2: The geometric size of vein-shaped microchip.



Figure S3: SEM image of AuNPs. Scale bar: 20nm.



Figure S4: UV - Visible absorption spectrum of AuNPs. The maximum absorption wavelength was 514-520 nm, and the maximum absorbance was 0.88-1.



Figure S5: Distribution of the flow field in Y-Z plane of the single-layer microchip.



Figure S6: The distributions of net flow field, flow field in Y-axis and Z-axis different cross sections of the microchip under different cross sections of microchip in the same flow throughput (a) the heights of bottom and top layer were equal; (b) the height of bottom layer was twice of the top layer; (c) the height of bottom layer was half of the top layer.



Figure S7: Normalized fluorescence intensity reflecting the FITC-labeled Goat anti-Mouse IgG antibodies immobilization on surface without modification, AuNP-functionalized surface, and self-assembled surface.



Figure S8: Characterization of the aggregation of AuNPs.



Figure S9: Different state of cells captured on the self-assembled surface. (a) single cell; (b)

multicellular cluster -- two cells; (c) multicellular cluster -- three cells.



Figure S10: The proportion of apoptotic/live cells after being cultured 0 min, 20 min, 40 min, and 60 min under the incubate time of 8min, 12min, 16min, and 20min.