

**ELECTRONIC SUPPLEMENTARY INFORMATION (ESI)**

**Ceria doping boosts methylene blue photodegradation in titania nanostructures**

**Chiara Alberoni<sup>1</sup>, Isabel Barroso-Martín<sup>2</sup>, Antonia Infantes-Molina<sup>2</sup>, Enrique Rodríguez-Castellón<sup>2</sup>, Aldo Talon<sup>1</sup>, Haiguang Zhao<sup>3</sup>, Shujie You<sup>4</sup>, Alberto Vomiero<sup>1,4\*</sup>, Elisa Moretti<sup>1\*</sup>**

<sup>1</sup>*Department of Molecular Sciences and Nanosystems, Ca' Foscari University of Venice, Via Torino 155, 30172, Venezia Mestre, Italy.*

<sup>2</sup>*Departamento de Química Inorgánica, Cristalografía y Mineralogía (Unidad Asociada al ICP-CSIC), Facultad de Ciencias, Universidad de Málaga, Campus de Teatinos, 29071 Málaga, Spain.*


<sup>3</sup>*Qingdao University - College of Physics & State Key Laboratory of Bio-Fibers and Eco-Textiles, 308 Ningxia Road, Qingdao 266071, P.R. China.*

<sup>4</sup>*Division of Material Science, Department of Engineering Sciences and Mathematics, Luleå University of Technology, 971 87, Luleå, Sweden.*


\* Corresponding authors: Elisa Moretti ([elisa.moretti@unive.it](mailto:elisa.moretti@unive.it)); Alberto Vomiero ([alberto.vomiero@unive.it](mailto:alberto.vomiero@unive.it)) & [alberto.vomiero@ltu.se](mailto:alberto.vomiero@ltu.se)). Tel. +39 0412346745 (E.M.); +39 0412346762 (A.V.).

**Table S1.** XPS binding energy values of O 1s and Ti 2p<sub>3/2</sub> core level signals of ceria-titania nanotubes and nanoparticles samples.


Sample	Binding Energy values (eV)			
	O 1s	Ti 2p		
<b>TNT</b>	530.1 (90.9)	531.6 (9.10)	-	458.6
<b>CeTNT0.25</b>	529.7 (87.0)	531.0 (13.0)	-	458.2
<b>CeTNT0.5</b>	529.8 (86.9)	531.2 (13.1)	-	458.5
<b>CeTNT1</b>	529.8 (92.1)	531.2 (7.90)	-	458.3
<b>CeTNT2.5</b>	529.9 (89.3)	531.3 (10.7)	-	458.3
<b>CeTNT5</b>	530.0 (89.1)	531.5 (10.9)	-	458.4
<b>TNP</b>	529.7 (95.7)	531.5 (4.30)	-	458.0
<b>CeTNP0.25</b>	529.7 (69.4)	531.4 (18.4)	533.5 (12.2)	458.2
<b>CeTNP0.5</b>	529.7 (87.9)	531.3 (9.80)	534.1 (2.30)	458.2
<b>CeTNP1</b>	529.7 (88.9)	531.5 (9.00)	534.0 (2.10)	458.2
<b>CeTNP2.5</b>	529.8 (84.2)	531.7 (12.5)	534.0 (3.30)	458.3
<b>CeTNP5</b>	529.8 (91.5)	531.4 (7.40)	533.7 (1.41)	458.1




**Figure S1.** Raman spectra of (a) Ce-TNT25 recorded at two different positions and (b) TNP, Ce-TNP25 and Ce-TNP50. Red circles mark anatase peaks. Asterisks in (b) mark weak peaks coming from reagents. Inset in (a): expanded view of the Ce-TNP5 sample, highlighting the presence of Rutile phase (black arrow).




**Figure S2.** Representative HRTEM of nanotube systems: a, b) CeTNT2.5 and c, d) CeTNT5 sample.




**Figure S3.** HRTEM micrographs of pure titania: (a)-(b) TNP sample; (c)-(d) TNT sample.



**Figure S4.** EDX spectrum of CeTNT5.



**Figure S5.** Representative HRTEM images of nanoparticle systems: a, b) CeTNP2.5 and c, d) CeTNP5.



**Figure S6.** X-ray diffraction patterns of the best performing samples, CeTNT025 and CeTNP025, before (fresh) and after (used) 3-cycle catalytic activity under UV light irradiation.