

# Stochastic Biosensors Based on N- and S-Doped Graphene for the Enantioanalysis of Aspartic Acid in Biological Samples

Raluca-Ioana Stefan-van Staden<sup>1,2\*</sup>, Damaris-Cristina Gheorghe<sup>1,2</sup>, Ruxandra-Maria Ilie-Mihai<sup>1\*</sup>, Lucian-Barbu Tudoran<sup>3</sup>, and Stela Maria Pruneanu<sup>3</sup>

- 1 Laboratory of Electrochemistry and PATLAB, National Institute of Research for Electrochemistry and Condensed Matter, 202 Splaiul Independentei Str., 060021, Bucharest-6, Romania; [ralucavanstaden@gmail.com](mailto:ralucavanstaden@gmail.com)
  - 2 Faculty of Applied Chemistry and Material Science, Politehnica University of Bucharest, Bucharest, Romania
  - 3 National Institute for Research and Development of Isotopic and Molecular Technologies, 67-103, Donat Street, Cluj-Napoca, Romania; [Stela.Pruneanu@itim-cj.ro](mailto:Stela.Pruneanu@itim-cj.ro)
- \* Correspondence: [ralucavanstaden@gmail.com](mailto:ralucavanstaden@gmail.com) (RI SvS); [i.ruxandra04@yahoo.com](mailto:i.ruxandra04@yahoo.com) (RMIM)





Figure S1. The UV-Vis spectrum of the E-NSGr sample



**Figure S2.** The FTIR spectrum of E-NSGr sample embedded in the KBr pellet (a,b)