## **Electronic Supplementary Material**

## Photosensitive metal–organic framework having a flower-like structure for effective visible light–driven photodegradation of rhodamine B

Lu Qin, Shicheng Zhao<sup>\*</sup> (🖾), Chenran Fan, Qian Ye

Shanghai Key Laboratory of Multiphase Materials Chemical Engineering, School of

Chemical Engineering, East China University of Science and Technology, 130

Meilong Road, Shanghai 200237, China

## This file includes

Figure S1. UV-vis spectra of TCPP and flower-like M-TCPP (M=Cu, Zn, Co).

- Figure S2. EDS point scan of flower-like Cu-TCPP.
- Figure S3. Thermogravimetric curve of flower-like Cu-TCPP.
- Figure S4. XRD spectra (a) and infrared spectra (b) of flower-like Cu-TCPP before and after the photocatalytic experiments.
- Figure S5. SEM images of flower-like Cu-TCPP before and after the photocatalytic reaction.
- Figure S6. Structure and standard curve of Rhodamine B.
- Figure S7. UV-vis DRS spectra of flower-like Cu-TCPP; (b)Tauc plot of the UV-vis DRS spectrum of flower-like Cu-TCPP.
- Figure S8. VB XPS diagram of flower-like Cu-TCPP.



Figure S1. UV–vis spectra of TCPP and flower-like M-TCPP (M=Cu Zn Co). (Added 10  $\mu L$  of TFA) .



Figure S2. EDS point scan of flower-like Cu-TCPP.



Figure S3. Thermogravimetric curve of flower-like Cu-TCPP.



Figure S4. XRD spectra (a) and infrared spectra (b) of Cu-TCPP nanoflowers before and after the photocatalytic experiments. Experimental conditions: RhB, 10 mg  $L^{-1}$ ; flower-like Cu-TCPP, 0.1 g  $L^{-1}$ ; initial pH 11.



Figure S5. SEM images of flower-like Cu-TCPP before and after the photocatalytic reaction. Experimental conditions: RhB, 10 mg  $L^{-1}$ ; flower-like Cu-TCPP, 0.1 g  $L^{-1}$ ; initial pH 11.



Figure S6. Structure and standard curve of Rhodamine B.



Figure S7. (a) UV-vis DRS spectra of flower-like Cu-TCPP; (b)Tauc plot of the UV-

vis DRS spectrum of flower-like Cu-TCPP.



Figure S8. VB XPS diagram of flower-like Cu-TCPP.