



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 17, 2020 – 03:19 PM GMT

PDB ID : 7BAU
Title : A de novo pentameric coiled-coil assembly: CC-Type2-(TgIaId)4-W19BrPhe.
Deposited on : 2020-12-16
Resolution : 1.42 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report.

This report is produced by the wwPDB biocuration pipeline after annotation of the structure.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.15.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.15.1

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.42 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.








Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2579 (1.44-1.40)
Clashscore	141614	2696 (1.44-1.40)
Ramachandran outliers	138981	2632 (1.44-1.40)
Sidechain outliers	138945	2631 (1.44-1.40)
RSRZ outliers	127900	2528 (1.44-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	32	<div> <div>91%</div> <div>6%</div> </div>
1	B	32	<div> <div>3%</div> <div>81%</div> <div>9%</div> <div>9%</div> </div>
1	C	32	<div> <div>94%</div> <div>6%</div> </div>
1	D	32	<div> <div>91%</div> <div>6%</div> </div>
1	E	32	<div> <div>97%</div> <div>.</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	32	
1	G	32	
1	H	32	
1	I	32	
1	J	32	

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 2460 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CC-Type2-(TgLaId)4-W19BrPhe..

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	30	Total	Br	C	N	O	0	0	0
			220	1	143	32	44			
1	B	29	Total	Br	C	N	O	0	0	0
			225	1	146	35	43			
1	C	30	Total	Br	C	N	O	0	0	1
			222	1	144	36	41			
1	D	31	Total	Br	C	N	O	0	0	1
			230	1	148	37	44			
1	E	32	Total	Br	C	N	O	0	0	1
			233	1	150	37	45			
1	F	30	Total	Br	C	N	O	0	1	1
			229	1	149	36	43			
1	G	30	Total	Br	C	N	O	0	0	0
			231	1	149	37	44			
1	H	31	Total	Br	C	N	O	0	0	1
			227	1	147	36	43			
1	I	31	Total	Br	C	N	O	0	0	1
			228	1	147	38	42			
1	J	31	Total	Br	C	N	O	0	0	0
			231	1	149	36	45			

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	14	Total	O	0	0
			14	14		
2	B	13	Total	O	0	0
			13	13		
2	C	21	Total	O	0	0
			21	21		
2	D	16	Total	O	0	0
			16	16		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	E	24	Total 24	O 24	0	0
2	F	21	Total 21	O 21	0	0
2	G	14	Total 14	O 14	0	0
2	H	15	Total 15	O 15	0	0
2	I	17	Total 17	O 17	0	0
2	J	29	Total 29	O 29	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: CC-Type2-(TgLaId)4-W19BrPhe.

Chain A:  91% 6%




- Molecule 1: CC-Type2-(TgLaId)4-W19BrPhe.

Chain B:  3% 81% 9% 9%



- Molecule 1: CC-Type2-(TgLaId)4-W19BrPhe.

Chain C:  94% 6%




- Molecule 1: CC-Type2-(TgLaId)4-W19BrPhe.

Chain D:  91% 6%




- Molecule 1: CC-Type2-(TgLaId)4-W19BrPhe.



Chain E:  97%





- Molecule 1: CC-Type2-(TgLaId)4-W19BrPhe.




- Molecule 1: CC-Type2-(TgLaId)4-W19BrPhe.




- Molecule 1: CC-Type2-(TgLaId)4-W19BrPhe.



- Molecule 1: CC-Type2-(TgLaId)4-W19BrPhe.



- Molecule 1: CC-Type2-(TgLaId)4-W19BrPhe.



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	40.60 Å 53.45 Å 56.16 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.60 – 1.42 40.60 – 1.42	Depositor EDS
% Data completeness (in resolution range)	98.8 (40.60-1.42) 98.8 (40.60-1.42)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.11 (at 1.42 Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.177 , 0.226 0.176 , 0.227	Depositor DCC
R_{free} test set	2341 reflections (5.23%)	wwPDB-VP
Wilson B-factor (Å ²)	13.3	Xtriage
Anisotropy	0.776	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 34.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.013 for -h,l,k 0.018 for -h,-l,-k 0.478 for h,-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2460	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 25.60 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.0745e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NH2, ACE, 4BF

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.87	0/204	0.66	0/274
1	B	0.78	0/211	0.66	0/280
1	C	0.82	0/207	0.79	0/275
1	D	0.84	0/215	0.84	0/285
1	E	0.83	0/216	0.68	0/287
1	F	0.77	0/217	0.80	0/289
1	G	0.80	0/217	0.70	0/287
1	H	0.84	0/211	0.68	0/281
1	I	0.82	0/213	0.82	0/282
1	J	0.83	0/215	0.70	0/286
All	All	0.82	0/2126	0.74	0/2826

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	220	0	223	1	0
1	B	225	0	244	2	0
1	C	222	0	240	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	230	0	250	1	0
1	E	233	0	251	0	0
1	F	229	0	251	3	0
1	G	231	0	257	3	0
1	H	227	0	249	4	0
1	I	228	0	253	0	0
1	J	231	0	249	2	0
2	A	14	0	0	0	0
2	B	13	0	0	0	0
2	C	21	0	0	0	0
2	D	16	0	0	0	0
2	E	24	0	0	0	0
2	F	21	0	0	0	0
2	G	14	0	0	0	0
2	H	15	0	0	1	0
2	I	17	0	0	0	0
2	J	29	0	0	0	0
All	All	2460	0	2467	10	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (10) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:27:THR:HG21	1:H:28:ILE:HG22	1.87	0.55
1:A:28:ILE:HG22	1:B:27:THR:HG21	1.91	0.53
1:B:2:GLU:HG2	1:B:5:GLN:H	1.74	0.52
1:F:14[B]:ILE:HD11	1:J:10:ILE:HA	1.92	0.51
1:G:16:GLU:OE2	1:H:22:LYS:NZ	2.27	0.49
1:H:16:GLU:HG3	2:H:110:HOH:O	2.15	0.46
1:F:14[A]:ILE:HD13	1:J:14:ILE:HD11	1.99	0.45
1:D:24:ILE:O	1:D:27:THR:HG22	2.19	0.42
1:G:6:THR:HG21	1:H:7:ILE:HG22	2.02	0.40
1:F:24:ILE:O	1:F:27:THR:HG22	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	27/32 (84%)	27 (100%)	0	0	100	100
1	B	26/32 (81%)	26 (100%)	0	0	100	100
1	C	27/32 (84%)	27 (100%)	0	0	100	100
1	D	28/32 (88%)	28 (100%)	0	0	100	100
1	E	29/32 (91%)	28 (97%)	1 (3%)	0	100	100
1	F	28/32 (88%)	28 (100%)	0	0	100	100
1	G	27/32 (84%)	27 (100%)	0	0	100	100
1	H	28/32 (88%)	28 (100%)	0	0	100	100
1	I	28/32 (88%)	28 (100%)	0	0	100	100
1	J	28/32 (88%)	27 (96%)	1 (4%)	0	100	100
All	All	276/320 (86%)	274 (99%)	2 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	19/23 (83%)	19 (100%)	0	100	100
1	B	22/23 (96%)	22 (100%)	0	100	100
1	C	21/23 (91%)	21 (100%)	0	100	100
1	D	22/23 (96%)	22 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	22/23 (96%)	22 (100%)	0	100	100
1	F	23/23 (100%)	23 (100%)	0	100	100
1	G	23/23 (100%)	23 (100%)	0	100	100
1	H	22/23 (96%)	22 (100%)	0	100	100
1	I	22/23 (96%)	22 (100%)	0	100	100
1	J	22/23 (96%)	22 (100%)	0	100	100
All	All	218/230 (95%)	218 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

10 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	4BF	A	19	1	11,12,13	0.53	0	12,15,17	0.33	0
1	4BF	B	19	1	11,12,13	0.50	0	12,15,17	0.58	0
1	4BF	C	19	1	11,12,13	0.51	0	12,15,17	0.76	0
1	4BF	D	19	1	11,12,13	0.67	0	12,15,17	0.85	0
1	4BF	E	19	1	11,12,13	0.39	0	12,15,17	0.79	1 (8%)
1	4BF	F	19	1	11,12,13	0.60	0	12,15,17	0.80	0
1	4BF	G	19	1	11,12,13	0.46	0	12,15,17	0.56	0
1	4BF	H	19	1	11,12,13	0.50	0	12,15,17	0.45	0
1	4BF	I	19	1	11,12,13	0.70	1 (9%)	12,15,17	0.83	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	4BF	J	19	1	11,12,13	0.52	0	12,15,17	0.76	1 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	4BF	A	19	1	-	0/5/6/8	0/1/1/1
1	4BF	B	19	1	-	0/5/6/8	0/1/1/1
1	4BF	C	19	1	-	0/5/6/8	0/1/1/1
1	4BF	D	19	1	-	0/5/6/8	0/1/1/1
1	4BF	E	19	1	-	0/5/6/8	0/1/1/1
1	4BF	F	19	1	-	0/5/6/8	0/1/1/1
1	4BF	G	19	1	-	0/5/6/8	0/1/1/1
1	4BF	H	19	1	-	0/5/6/8	0/1/1/1
1	4BF	I	19	1	-	0/5/6/8	0/1/1/1
1	4BF	J	19	1	-	0/5/6/8	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	I	19	4BF	O-C	2.04	1.28	1.19

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	19	4BF	CG-CB-CA	-2.26	109.52	114.10
1	J	19	4BF	CG-CB-CA	-2.08	109.89	114.10

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

CONFIDENTIAL VALIDATION REPORT

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	28/32 (87%)	-0.22	0 100 100	14, 17, 30, 35	0
1	B	28/32 (87%)	-0.01	1 (3%) 42 42	13, 20, 39, 44	0
1	C	28/32 (87%)	-0.12	0 100 100	12, 15, 24, 29	0
1	D	29/32 (90%)	-0.13	0 100 100	13, 17, 27, 32	0
1	E	29/32 (90%)	-0.14	0 100 100	12, 16, 23, 33	0
1	F	28/32 (87%)	-0.14	1 (3%) 42 42	13, 16, 25, 36	0
1	G	29/32 (90%)	0.19	3 (10%) 6 5	12, 21, 37, 44	0
1	H	29/32 (90%)	-0.07	1 (3%) 45 44	13, 17, 29, 40	0
1	I	29/32 (90%)	-0.14	1 (3%) 45 44	13, 17, 28, 33	0
1	J	29/32 (90%)	-0.14	0 100 100	12, 16, 23, 38	0
All	All	286/320 (89%)	-0.09	7 (2%) 59 58	12, 17, 34, 44	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	30	GLY	4.0
1	G	3	ILE	3.4
1	I	30	GLY	2.4
1	G	30	GLY	2.3
1	G	1	GLY	2.3
1	B	3	ILE	2.3
1	F	30	GLY	2.2

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column

labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
1	4BF	H	19	12/13	0.65	0.14	15,21,30,48	0
1	4BF	A	19	12/13	0.67	0.15	15,22,33,48	0
1	4BF	I	19	12/13	0.82	0.16	12,18,27,30	0
1	4BF	F	19	12/13	0.83	0.16	12,21,32,34	0
1	4BF	D	19	12/13	0.84	0.15	12,19,26,30	0
1	4BF	C	19	12/13	0.84	0.16	12,20,30,31	0
1	4BF	E	19	12/13	0.85	0.15	11,15,20,26	0
1	4BF	J	19	12/13	0.88	0.14	11,15,19,25	0
1	4BF	G	19	12/13	0.89	0.12	11,15,21,27	0
1	4BF	B	19	12/13	0.91	0.13	12,17,20,26	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.