

Electronic Supporting Information (ESI)

Gadofullerene based liver-specific MRI contrast agent for early diagnosis of orthotopic hepatocellular carcinoma

Zhongpu Zhao,^{ab} Mingming Zhen*,^{ab} Chen Zhou,^{ab} Lei Li,^{ab} Wang Jia,^{ab} Shuai Liu,^{ab}
Xue Li,^{ab} Xiaodan Liao^{ab} and Chunru Wang*^{ab}

^aBeijing National Laboratory for Molecular Sciences, Key Laboratory of Molecular
Nanostructure and Nanotechnology, CAS Research/Education Center for Excellence
in Molecular Sciences, Institute of Chemistry, Chinese Academy of Sciences, Beijing,
100190, China

^b University of Chinese Academy of Sciences, Beijing, 100049, China

Corresponding Author

*E-mail: zhenmm@iccas.ac.cn; crwang@iccas.ac.cn

Table of Contents

Fig. S1. the FT-IR spectrum of GF-NH₂ and GF-Ala.

Fig. S2. the AFM of GF-NH₂ and GF-Ala.

Fig. S3 XPS results of GF-OH (a), GF-Ala (b), GF-NH₂ (c and d).




Fig. S1 the FT-IR spectrum of GF-NH₂ (a) and GF-Ala (b).




Fig. S2 the AFM of GF-NH₂ (a) and GF-Ala (b).




Fig. S3 XPS results of GF-OH (a), GF-Ala (b), GF-NH₂ (c and d).