

Chondrocytes secretome enriched microparticles encapsulated with chondrocyte membrane to facilitate the chondrogenesis of BMSCs and reduce hypertrophy

Zizhao Tang^{1,2}, Yan Lu^{1,2}, Shixin Zhang^{1,2}, Jing Wang³, Qiguang Wang^{1,2}, Yumei Xiao^{1,2}, and Xingdong Zhang^{1,2}*

1 National Engineering Research Center for Biomaterials, Sichuan University, 29 Wangjiang Road, Chengdu, Sichuan, 610065, China

2 College of Biomedical Engineering, Sichuan University, 29 Wangjiang Road, Chengdu, Sichuan, 610065, China

3 College of Materials and Chemistry & Chemical Engineering, Chengdu University of Technology, Chengdu, Sichuan, 610059, China

Corresponding Author

*Email: wqgwang@126.com

Table. S1. Primers were used to amplify mRNAs encoding rabbit GAPDH.

Gene	Forward primer	Reverse primer
<i>GAPDH</i>	TCGGAGTGAACGGATTTGGC	TTCCCGTTCTCAGCCTTGAC
<i>COL2A1</i>	TGATAAGGATGTGTGGAAGCCG	CAGGCAGTCCTTGGTGTCTTC
<i>AGG</i>	GGCCACTGTTACCGTCACTT	GTCCTGAGCGTTGTTGTTGAC
<i>SOX9</i>	TCTGGAGACTGCTGAACGAG	CTGCCCATTCCTCACCGACTT
<i>COL1A1</i>	AGAGGACCACGTGGAGAAAG	CCATCAAACCTGAGCAGCAAA
<i>COLX</i>	TCCCAGA ACCCAGAATCCATC	GGTTGTGGGCCT TTTATGCC