Supplementary Information

Multi-Functional Switch Effect in Interlocking Molecular Rotators-on-Graphene Systems Using Electric Fields

Xiaobo Li^{1,2*}, Fei Qi^{2,3}, Rundong Zhao^{2,4}, Zijie Qiu⁵, Yun Li⁶, Mengqiu Long⁶ and Guanghui Zhou⁷

- ¹ Department of Applied Physics, School of Microelectronics and Physics, Hunan University of Technology and Business, Changsha 410205, China
- ² Institute of Computational and Theoretical Studies & Department of Physics, Hong Kong Baptist University, Hong Kong SAR, China
- ³ CAS Key Laboratory of Quantitative Engineering Biology, Shenzhen Institute of Synthetic Biology, Shenzhen Institutes of Advanced Technology, Chinese Academy of Sciences, Shenzhen 518055, China
- ⁴School of Physics, Beihang University, Beijing 100191, China
- ⁵Max Planck Institute for Polymer Research, Ackermannweg 10, 55128 Mainz, Germany
- ⁶ Hunan Key Laboratory of Super-Microstructure and Ultrafast Process, School of Physics and Electronics, Central South University, Changsha 410083, China
- ⁷ Department of Physics and Key Laboratory for Low-Dimensional Structures and Quantum Manipulation (Ministry of Education), Hunan Normal University, Changsha 410081, China

^{*} Corresponding author; E-mail: xiaoboli2010@hutb.edu.cn; xiaoboli2010@gmail.com(X. L.)



Figure S1: Density of states (DOS) as a function of energy. (a) The DOS of the rotators-on-graphene system without leads; (b) The projected DOS of the rotators and the rotating pivot of Chromium atoms.