

Supporting information

A mitochondria targetable dimethylphosphorothionate-based far-red and colorimetric fluorescent probe with large Stokes shift for monitoring peroxynitrite in living cells

Chunxiang Zhang, Xiangyang Zhang, Youming Shen, Zile Zhou*

Hunan Provincial Key Laboratory of Water Treatment Functional Materials, College of Chemistry and Materials Engineering, Hunan University of Arts and Science, Changde 415000, PR China

*Corresponding author. Fax: 0746-7186115

E-mail address: Zile_Zhou@163.com (Z. Zhou)




Fig. S1 ^1H NMR spectrum of PCPA (400 MHz, $\text{DMSO-}d_6$).




Fig. S2 ^{13}C NMR spectrum of PCPA (100 MHz, $\text{DMSO-}d_6$).




Fig. S3 TOF-MS spectrum of PCPA.




Fig. S4 Fluorescence intensity of PCPA in the presence of ONOO⁻ containing other analytes in PBS (20 mM, pH 7.4, 10% DMSO). (1) H₂O₂, (2) ClO⁻, (3) ONOO⁻, (4) NO, (5) ¹O₂, (6) ·OH, (7) NO₃⁻, (8) TBHP, (9) O₂⁻, (10) HNO, (11) F⁻, (12) Cl⁻, (13) Br⁻, (14) I⁻, (15) Ac⁻, (16) SO₃²⁻, (17) S₂O₃²⁻, (18) S²⁻, (19) Cys, (20) Hcy, (21) GSH⁻, (22) K⁺, (23) Mg²⁺, (24) Zn²⁺, (25) Al³⁺. λ_{ex} = 445 nm, λ_{em} = 645 nm.




Fig. S5 ^1H NMR spectrum of PCPA + ONOO^- .




Fig. S6 TOF-MS spectrum of PCPA + ONOO^- .




Fig. S7 Cytotoxicity assays of PCPA with different concentrations (2, 4, 6, 8, 10 and 12 µM) in HeLa cells.