

Supplementary material for manuscript:

Synthesis, structural characterization and DFT study of N-(pyrimidyl)- ω -amino acids/peptide: β -alanine, γ -aminobutyric, 5-aminovaleric, 6-aminohexanoic and glycylglycine

Angel García-Raso^{a,*}, Angel Terrón^a, Juan J. Fiol^a, Adela López-Zafra^a, Gemma Picó, Antonio Frontera^{a,*}, Ezequiel M. Vázquez-López^b, Miquel Barceló-Oliver^a

^aDepartament de Química, Universitat de les Illes Balears, Crta. de Valldemossa km 7.5, 07122 Palma (Balears), Spain. E-mail: jfa950@uib.es; toni.frontera@uib.es.

^bUniversidade de Vigo, Departamento de Química Inorgánica, Facultade de Química, Edificio Ciencias Experimentais, E-36310 Vigo, Galicia, Spain. E-mail: ezequiel@uvigo.es.




Figure S1. Asymmetric unit of compound 2




Figure S2. Asymmetric unit of compound **2a**. H-atoms omitted for clarity




Figure S3. Asymmetric unit of compound **3a**

ORTEP PLOTS of compounds **1-5** at 50% probability level.




Figure S4. ORTEP plot of compound 1




Figure S5. ORTEP plot of compound 2




Figure S6. ORTEP plot of compound 2a




Figure S7. ORTEP plot of compound 3




Figure S8. ORTEP plot of compound **3a**




Figure S9. ORTEP plot of compound **4**.




Figure S9. ORTEP plot of compound 4a




Figure S10. ORTEP plot of compound 5