

Supplementary Material

Theoretical study of the CO₂–O₂ van der Waals complex: potential energy surface and applications

Yosra Ajili,¹ Ernesto Quintas-Sánchez,² Bilel Mehnen,³ Piotr S. Żuchowski,³ Filip Brzęk,³ Nayla El-Kork,⁴ Marko Gacesa,⁴ Richard Dawes² and Majdi Hochlaf^{5,*}

¹ Université de Tunis El Manar, Faculté des Sciences de Tunis, LR01ES09 Spectroscopie Atomique et Moléculaire et Applications, 1060, Tunis, Tunisia.

² Department of Chemistry, Missouri University of Science and Technology, Rolla, MO 65409-0010, USA.

³ Institute of Physics, Faculty of Physics, Astronomy and Informatics, Nicolaus Copernicus University, Grudziadzka 5, 87-100 Torun, Poland.

⁴ Space and Planetary Sciences Center, and Department of Physics, Khalifa University, P.O. Box 127788, Abu Dhabi, United Arab Emirates.

⁵ Université Gustave Eiffel, COSYS/IMSE, 5 Bd Descartes 77454, Champs sur Marne, France.




Figure S1: The SAPT/aug-cc-pVTZ contributions are plotted for radial cuts through the 4D-PES passing through each of the 6 structures given in Figure 1.