

Ruthenium-based assemblies incorporating tetrapyrridylporphyrin panels: A photosensitizers delivery strategy for the treatment of rheumatoid arthritis by photodynamic therapy

Manuel Gallardo-Villagr  n ^{a,b}, Lucie Paulus ^b, Jean-Louis Charissoux ^c, David Yannick Leger ^b, Pascale Vergne-Salle ^d, Bruno Therrien ^{a,†,*}, Bertrand Liagre ^{b,†,*}

^a Institut de Chimie, Universit   de Neuch  tel, Avenue de Bellevaux 51, CH-2000 Neuch  tel, Switzerland;

^b Universit   de Limoges, Laboratoire PEIRENE UR 22722, Facult   de Pharmacie, F-87025 Limoges, France;

^c Service d'Orthop  die-Traumatologie, CHRU Dupuytren, 2 avenue Martin Luther King, 87042 Limoges Cedex, France;

^d Service de Rhumatologie, CHRU Dupuytren 2, 16 rue Bernard Descottes, 87042 Limoges Cedex, France;

* Correspondence: bruno.therrien@unine.ch; bertrand.liagre@unilim.fr

† These authors contributed equally to the work.

Supporting information

Table of Contents

I) UV-vis spectra	2
C1	2
Zn-C1	2
C2	3
Zn-C2	3
Co-C2	4
II) Fluorescence spectra	4
C1	4
Zn-C1	5
C2	5
Zn-C2	6
Co-C2	6

I) UV-vis spectra:




Fig S1. UV-vis absorbance spectrum of C1 in DMSO (10 nM).




Fig S2. UV-vis absorbance spectrum of Zn-C1 in DMSO (10 nM).




Fig S3. UV-vis absorbance spectrum of **C2** in DMSO (10 nM).




Fig S4. UV-vis absorbance spectrum of **Zn-C2** in DMSO (10 nM).




Fig S5. UV-vis absorbance spectrum of **Co-C2** in DMSO (10 nM).

II) Fluorescence spectra:




Fig S6. Fluorescence emission spectrum of **C1** in DMSO (10 nM).




Fig S7. Fluorescence emission spectrum of **Zn-C1** in DMSO (10 nM).




Fig S8. Fluorescence emission spectrum of **C2** in DMSO (10 nM).




Fig S9. Fluorescence emission spectrum of **Zn-C2** in DMSO (10 nM).




Figure S10. Fluorescence emission spectrum of **Co-C2** in DMSO (10 nM).