

C3 compounds

For C3 oxidation, considered chemical pathways represent at least 75% of the estimated reactivity.

Reactions		k_{298} ($M^{-n+1} s^{-1}$)	Ea/R (K)	References	Notes
Oxidation of 1-propanol					
Pathway 1: $CH_3CH_2CH_2(OH) + HO^\bullet \rightarrow C^\bullet H_2CH_2CH_2(OH) + H_2O$		$4.3 \cdot 10^8$			BR: 15% - 1
$C^\bullet H_2CH_2CH_2(OH) + O_2 \rightarrow CH_2(OH)CH_2CH_2(OO^\bullet)$		$4.7 \cdot 10^9$		Adams and Willson, 1969	
Pathway 2: $CH_3CH_2CH_2(OH) + HO^\bullet \rightarrow CH_3C^\bullet HCH_2(OH) + H_2O$		$4.3 \cdot 10^8$			BR: 15% - 1
$CH_3C^\bullet HCH_2(OH) + O_2 \rightarrow CH_3CH(OO^\bullet)CH_2(OH)$		$4.7 \cdot 10^9$		Adams and Willson, 1969	
Pathway 3: $CH_3CH_2CH_2(OH) + HO^\bullet \rightarrow CH_3CH_2C^\bullet H(OH) + H_2O$		$1.9 \cdot 10^9$			BR: 70% - 1
$CH_3CH_2C^\bullet H(OH) + O_2 \rightarrow CH_3CH_2CH(OH)(OO^\bullet)$		$4.7 \cdot 10^9$		Adams and Willson, 1969	
$CH_3CH_2CH_2(OH) + HO^\bullet \rightarrow 0.15 CH_2(OH)CH_2CH_2(OO^\bullet) + 0.15 CH_3CH(OO^\bullet)CH_2(OH) + 0.70 CH_3CH_2CH(OH)(OO^\bullet) + H_2O - O_2$	R(389)	$2.7 \cdot 10^9$	782	Monod et al., 2005	
Pathway 1: $CH_3CH_2CH_2(OH) + NO_3^\bullet \rightarrow C^\bullet H_2CH_2CH_2(OH) + NO_3^- + H^+$		$5.1 \cdot 10^5$			BR: 15%
$C^\bullet H_2CH_2CH_2(OH) + O_2 \rightarrow CH_2(OH)CH_2CH_2(OO^\bullet)$		$4.7 \cdot 10^9$		Adams and Willson, 1969	
Pathway 2: $CH_3CH_2CH_2(OH) + NO_3^\bullet \rightarrow CH_3C^\bullet HCH_2(OH) + NO_3^- + H^+$		$5.1 \cdot 10^5$			BR: 15%
$CH_3C^\bullet HCH_2(OH) + O_2 \rightarrow CH_3CH(OO^\bullet)CH_2(OH)$		$4.7 \cdot 10^9$		Adams and Willson, 1969	
Pathway 3: $CH_3CH_2CH_2(OH) + NO_3^\bullet \rightarrow CH_3CH_2C^\bullet H(OH) + NO_3^- + H^+$		$2.2 \cdot 10^6$			BR: 70%
$CH_3CH_2C^\bullet H(OH) + O_2 \rightarrow CH_3CH_2CH(OH)(OO^\bullet)$		$4.7 \cdot 10^9$		Adams and Willson, 1969	
$CH_3CH_2CH_2(OH) + NO_3^\bullet \rightarrow 0.15 CH_2(OH)CH_2CH_2(OO^\bullet) + 0.15 CH_3CH(OO^\bullet)CH_2(OH) + 0.70 CH_3CH_2CH(OH)(OO^\bullet) + NO_3^- + H^+ - O_2$	R(390)	$3.2 \cdot 10^6$		Herrmann et al., 1994	2
Pathway 1: $2 CH_2(OH)CH_2CH_2(OO^\bullet) \rightarrow 2 CH_2(OH)CH_2CHO + H_2O_2$		$3.0 \cdot 10^7$			BR: 20% - 3
Pathway 2: $2 CH_2(OH)CH_2CH_2(OO^\bullet) \rightarrow 2 CH_2(OH)CH_2CH_2(O^\bullet) + O_2$		$1.3 \cdot 10^8$			BR: 80% - 3
$CH_2(OH)CH_2CH_2(O^\bullet) \rightarrow CH_2(OH)CH_2C^\bullet H(OH)$					4
$CH_2(OH)CH_2C^\bullet H(OH) + O_2 \rightarrow CH_2(OH)CH_2CH(OH)(OO^\bullet)$		$2.0 \cdot 10^9$			5
$2 CH_2(OH)CH_2CH_2(OO^\bullet) \rightarrow 0.40 CH_2(OH)CH_2CHO + 1.60 CH_2(OH)CH_2CH(OH)(OO^\bullet) + 0.20 H_2O_2 - 0.80 O_2$	R(391)	$1.6 \cdot 10^8$	-1600		= $k(2 CH_3CH_2(OO^\bullet)) - 8$
Pathway 1: $2 CH_3CH(OO^\bullet)CH_2(OH) \rightarrow 2 CH_3COCH_2(OH) + H_2O_2$		$5.0 \cdot 10^7$			BR: 50%
Pathway 2: $2 CH_3CH(OO^\bullet)CH_2(OH) \rightarrow CH_3COCH_2(OH) + CH_3CH(OH)CH_2(OH) + O_2$		$3.3 \cdot 10^7$			BR: 33%
Pathway 3: $2 CH_3CH(OO^\bullet)CH_2(OH) \rightarrow 2 CH_3CH(O^\bullet)CH_2(OH) + O_2$		$1.7 \cdot 10^7$			BR: 17%
$CH_3CH(O^\bullet)CH_2(OH) \rightarrow CH_3CHO + C^\bullet H_2(OH)$					6 - 7
$C^\bullet H_2(OH) + O_2 \rightarrow CH_2(OH)(OO^\bullet)$		$2.0 \cdot 10^9$			5
$2 CH_3CH(OO^\bullet)CH_2(OH) \rightarrow 1.33 CH_3COCH_2(OH) + 0.33 CH_3CH(OH)CH_2(OH) + 0.34 CH_2(OH)(OO^\bullet) + 0.34 CH_3CHO + 0.50 H_2O_2 + 0.16 O_2$	R(392)	$1.0 \cdot 10^8$			= $k(2 CH_2(OH)CH_2(OO^\bullet)) - 8$
$CH_3CH_2CH(OH)(OO^\bullet) + OH^- \rightarrow CH_3CH_2CH(O^\bullet)(OO^\bullet) + H_2O$		$4.0 \cdot 10^9$			
$CH_3CH_2CH(O^\bullet)(OO^\bullet) \rightarrow CH_3CH_2CHO + O_2^\bullet$					9
$CH_3CH_2CH(OH)(OO^\bullet) + OH^- \rightarrow CH_3CH_2CHO + O_2^{\bullet -} + H_2O$	R(393)	$4.0 \cdot 10^9$			= $k(CH_3CH(OH)(OO^\bullet) + OH^-)$
$CH_3CH_2CH(OH)(OO^\bullet) \rightarrow CH_3CH_2CHO + HO_2^\bullet$	R(394)	52			10
$CH_2(OH)CH_2CH(OH)(OO^\bullet) + OH^- \rightarrow CH_2(OH)CH_2CH(O^\bullet)(OO^\bullet) + H_2O$		$4.0 \cdot 10^9$			
$CH_2(OH)CH_2CH(O^\bullet)(OO^\bullet) \rightarrow CH_2(OH)CH_2CHO + O_2^\bullet$					9
$CH_2(OH)CH_2CH(OH)(OO^\bullet) + OH^- \rightarrow CH_2(OH)CH_2CHO + O_2^{\bullet -} + H_2O$	R(395)	$4.0 \cdot 10^9$			= $k(CH_3CH(OH)(OO^\bullet) + OH^-)$
$CH_2(OH)CH_2CH(OH)(OO^\bullet) \rightarrow CH_2(OH)CH_2CHO + HO_2^\bullet$	R(396)	52			10
Oxidation of 3-hydroxypropionaldehyde					11
Pathway 1: $CH_2(OH)CH_2CHO + HO^\bullet \rightarrow CH_2(OH)CH_2C^\bullet O + H_2O$		$2.6 \cdot 10^9$			BR: 62% - 12
$CH_2(OH)CH_2C^\bullet O + O_2 \rightarrow CH_2(OH)CH_2CO(OO^\bullet)$		$2.0 \cdot 10^9$			5
Pathway 2: $CH_2(OH)CH_2CHO + HO^\bullet \rightarrow C^\bullet H(OH)CH_2CHO + H_2O$		$1.5 \cdot 10^9$			BR: 38% - 12
$C^\bullet H(OH)CH_2CHO + O_2 \rightarrow CHOCH_2CH(OH)(OO^\bullet)$		$2.0 \cdot 10^9$			5

Reactions		k ₂₉₈ (M ⁻ⁿ⁺¹ s ⁻¹)	Ea/R (K)	References	Notes
CH ₂ (OH)CH ₂ CHO + HO• → 0.62 CH ₂ (OH)CH ₂ CO(OO•) + 0.38 CHOCH ₂ CH(OH)(OO•) + H ₂ O - O ₂	R(397)	4.1 10 ⁹			13
Pathway 1: CH ₂ (OH)CH ₂ CHO + NO ₃ • → CH ₂ (OH)CH ₂ C•O + NO ₃ ⁻ + H ⁺		1.9·10 ⁶			BR: 62%
CH ₂ (OH)CH ₂ C•O + O ₂ → CH ₂ (OH)CH ₂ CO(OO•)		2.0·10 ⁹			5
Pathway 2: CH ₂ (OH)CH ₂ CHO + NO ₃ • → C•H(OH)CH ₂ CHO + NO ₃ ⁻ + H ⁺		1.2·10 ⁶			BR: 38%
C•H(OH)CH ₂ CHO + O ₂ → CHOCH ₂ CH(OH)(OO•)		2.0·10 ⁹			5
CH ₂ (OH)CH ₂ CHO + NO ₃ • → 0.62 CH ₂ (OH)CH ₂ CO(OO•) + 0.38 CHOCH ₂ CH(OH)(OO•) + NO ₃ ⁻ + H ⁺ - O ₂	R(398)	3.1 10 ⁶			= k(CH ₂ (OH)CHO + NO ₃ •) - 2
Pathway 1: CH ₂ (OH)CH ₂ CH(OH)(OH) + HO• → CH ₂ (OH)CH ₂ C•(OH)(OH) + H ₂ O		1.1·10 ⁹			BR: 33% - 14
CH ₂ (OH)CH ₂ C•(OH)(OH) + O ₂ → CH ₂ (OH)CH ₂ C(OH)(OH)(OO•)		2.0·10 ⁹			5
Pathway 2: CH ₂ (OH)CH ₂ CH(OH)(OH) + HO• → C•H(OH)CH ₂ CH(OH)(OH) + H ₂ O		2.1·10 ⁹			BR: 67% - 14
C•H(OH)CH ₂ CH(OH)(OH) + O ₂ → CH(OH)(OH)CH ₂ CH(OH)(OO•)		2.0·10 ⁹			5
CH ₂ (OH)CH ₂ CH(OH)(OH) + HO• → 0.33 CH ₂ (OH)CH ₂ C(OH)(OH)(OO•) + 0.67 CH(OH)(OH)CH ₂ CH(OH)(OO•) + H ₂ O - O ₂	R(399)	3.2 10 ⁹			13
Pathway 1: CH ₂ (OH)CH ₂ CH(OH)(OH) + NO ₃ • → CH ₂ (OH)CH ₂ C•(OH)(OH) + NO ₃ ⁻ + H ⁺		3.6·10 ⁵			BR: 33%
CH ₂ (OH)CH ₂ C•(OH)(OH) + O ₂ → CH ₂ (OH)CH ₂ C(OH)(OH)(OO•)		2.0·10 ⁹			5
Pathway 2: CH ₂ (OH)CH ₂ CH(OH)(OH) + NO ₃ • → C•H(OH)CH ₂ CH(OH)(OH) + NO ₃ ⁻ + H ⁺		7.4·10 ⁵			BR: 67%
C•H(OH)CH ₂ CH(OH)(OH) + O ₂ → CH(OH)(OH)CH ₂ CH(OH)(OO•)		2.0·10 ⁹			5
CH ₂ (OH)CH ₂ CH(OH)(OH) + NO ₃ • → 0.33 CH ₂ (OH)CH ₂ C(OH)(OH)(OO•) + 0.67 CH(OH)(OH)CH ₂ CH(OH)(OO•) + NO ₃ ⁻ + H ⁺ - O ₂	R(400)	1.1 10 ⁶			= k(CH ₂ (OH)CH(OH)(OH) + NO ₃ •) - 2
2 CH ₂ (OH)CH ₂ CO(OO•) → 2 CH ₂ (OH)CH ₂ CO(O•) + O ₂		1.6·10 ⁸			
CH ₂ (OH)CH ₂ CO(O•) → CH ₂ (OH)C•H ₂ + CO ₂					6 - 7
CH ₂ (OH)C•H ₂ + O ₂ → CH ₂ (OH)CH ₂ (OO•)		2.0·10 ⁹			5
2 CH ₂ (OH)CH ₂ CO(OO•) → 2 CH ₂ (OH)CH ₂ (OO•) + 2 CO ₂ - O ₂	R(401)	1.6·10 ⁸	-1600		= k(2 CH ₃ CH ₂ (OO•))
CHOCH ₂ CH(OH)(OO•) + OH• → CHOCH ₂ CH(O•)(OO•) + H ₂ O		4.0 10 ⁹			
CHOCH ₂ CH(O•)(OO•) → CHOCH ₂ CHO + O ₂ •					9
CHOCH ₂ CH(OH)(OO•) + OH• → CHOCH ₂ CHO + O ₂ • + H ₂ O	R(402)	4.0 10 ⁹			= k(CH ₃ CH(OH)(OO•) + OH•)
CHOCH ₂ CH(OH)(OO•) → CHOCH ₂ CHO + HO ₂ •	R(403)	52			10
CH ₂ (OH)CH ₂ C(OH)(OH)(OO•) + OH• → CH ₂ (OH)CH ₂ C(OH)(O•)(OO•) + H ₂ O		4.0 10 ⁹			
CH ₂ (OH)CH ₂ C(OH)(O•)(OO•) → CH ₂ (OH)CH ₂ CO(OH) + O ₂ •					9
CH ₂ (OH)CH ₂ C(OH)(OH)(OO•) + OH• → CH ₂ (OH)CH ₂ CO(OH) + O ₂ • + H ₂ O	R(404)	4.0 10 ⁹			= k(CH ₃ CH(OH)(OO•) + OH•)
CH ₂ (OH)CH ₂ C(OH)(OH)(OO•) → CH ₂ (OH)CH ₂ CO(OH) + HO ₂ •	R(405)	1.0 10 ⁶			15
CH(OH)(OH)CH ₂ CH(OH)(OO•) + OH• → CH(OH)(OH)CH ₂ CH(O•)(OO•) + H ₂ O		4.0 10 ⁹			
CH(OH)(OH)CH ₂ CH(O•)(OO•) → CHOCH ₂ CH(OH)(OH) + O ₂ •					9
CH(OH)(OH)CH ₂ CH(OH)(OO•) + OH• → CHOCH ₂ CH(OH)(OH) + O ₂ • + H ₂ O	R(406)	4.0 10 ⁹			= k(CH ₃ CH(OH)(OO•) + OH•)
CH(OH)(OH)CH ₂ CH(OH)(OO•) → CHOCH ₂ CH(OH)(OH) + HO ₂ •	R(407)	52			10
Oxidation of propane-1,2-diol					
Pathway 1: CH ₃ CH(OH)CH ₂ (OH) + HO• → CH ₃ CH(OH)C•H(OH) + H ₂ O		1.0·10 ⁹			BR: 60% - 16
CH ₃ CH(OH)C•H(OH) + O ₂ → CH ₃ CH(OH)CH(OH)(OO•)		2.0·10 ⁹			5
Pathway 2: CH ₃ CH(OH)CH ₂ (OH) + HO• → CH ₃ C•(OH)CH ₂ (OH) + H ₂ O		7.0·10 ⁸			BR: 40% - 16
CH ₃ C•(OH)CH ₂ (OH) + O ₂ → CH ₃ C(OH)(OO•)CH ₂ (OH)		2.0·10 ⁹			5
CH ₃ CH(OH)CH ₂ (OH) + HO• → 0.60 CH ₃ CH(OH)CH(OH)(OO•) + 0.40 CH ₃ C(OH)(OO•)CH ₂ (OH) + H ₂ O - O ₂	R(408)	1.7 10 ⁹		Hoffmann et al., 2009	
Pathway 1: CH ₃ CH(OH)CH ₂ (OH) + NO ₃ • → CH ₃ CH(OH)C•H(OH) + NO ₃ ⁻ + H ⁺		5.9·10 ⁶			BR: 60%
CH ₃ CH(OH)C•H(OH) + O ₂ → CH ₃ CH(OH)CH(OH)(OO•)		2.0·10 ⁹			5
Pathway 2: CH ₃ CH(OH)CH ₂ (OH) + NO ₃ • → CH ₃ C•(OH)CH ₂ (OH) + NO ₃ ⁻ + H ⁺		4.0·10 ⁶			BR: 40%
CH ₃ C•(OH)CH ₂ (OH) + O ₂ → CH ₃ C(OH)(OO•)CH ₂ (OH)		2.0·10 ⁹			5
CH ₃ CH(OH)CH ₂ (OH) + NO ₃ • → 0.60 CH ₃ CH(OH)CH(OH)(OO•) + 0.40 CH ₃ C(OH)(OO•)CH ₂ (OH) + NO ₃ ⁻ + H ⁺ - O ₂	R(409)	9.9 10 ⁶		Hoffmann et al., 2009	2
CH ₃ CH(OH)CH(OH)(OO•) + OH• → CH ₃ CH(OH)CH(O•)(OO•) + H ₂ O		4.0 10 ⁹			

Reactions		k ₂₉₈ (M ⁻ⁿ⁺¹ s ⁻¹)	Ea/R (K)	References	Notes
CH ₃ CH(OH)CH(O ⁻)(OO [•]) → CH ₃ CH(OH)CHO + O ₂ ^{•-}					9
CH ₃ CH(OH)CH(OH)(OO [•]) + OH ⁻ → CH ₃ CH(OH)CHO + O ₂ ^{•-} + H ₂ O	R(410)	4.0 10 ⁹			= k(CH ₃ CH(OH)(OO [•]) + OH ⁻)
CH ₃ CH(OH)CH(OH)(OO [•]) → CH ₃ CH(OH)CHO + HO ₂ [•]	R(411)	1.9 10 ²			10
CH ₃ C(OH)(OO [•])CH ₂ (OH) + OH ⁻ → CH ₃ C(O ⁻)(OO [•])CH ₂ (OH) + H ₂ O		4.0 10 ⁹			
CH ₃ C(O ⁻)(OO [•])CH ₂ (OH) → CH ₃ COCH ₂ (OH) + O ₂ ^{•-}					9
CH ₃ C(OH)(OO [•])CH ₂ (OH) + OH ⁻ → CH ₃ COCH ₂ (OH) + O ₂ ^{•-} + H ₂ O	R(412)	4.0 10 ⁹			= k(CH ₃ CH(OH)(OO [•]) + OH ⁻)
CH ₃ C(OH)(OO [•])CH ₂ (OH) → CH ₃ COCH ₂ (OH) + HO ₂ [•]	R(413)	1.9 10 ²			10
Oxidation of 2-hydroxypropionaldehyde					17
Pathway 1: CH ₃ CH(OH)CH(OH)(OH) + HO [•] → CH ₃ CH(OH)CH(OH)(O [•]) + H ₂ O		4.8 10 ⁸			BR: 37% - 18
CH ₃ CH(OH)CH(OH)(O [•]) → CH ₃ C [•] H(OH) + CHO(OH)					6 - 7
CH ₃ C [•] H(OH) + O ₂ → CH ₃ CH(OH)(OO [•])		2.0·10 ⁹			5
Pathway 2: CH ₃ CH(OH)CH(OH)(OH) + HO [•] → CH ₃ CH(OH)C [•] (OH)(OH) + H ₂ O		4.6 10 ⁸			BR: 35% - 18
CH ₃ CH(OH)C [•] (OH)(OH) + O ₂ → CH ₃ CH(OH)C(OH)(OH)(OO [•])		2.0·10 ⁹			5
Pathway 3: CH ₃ CH(OH)CH(OH)(OH) + HO [•] → CH ₃ C [•] (OH)CH(OH)(OH) + H ₂ O		3.6 10 ⁸			BR: 28% - 18
CH ₃ C [•] (OH)CH(OH)(OH) + O ₂ → CH ₃ C(OH)(OO [•])CH(OH)(OH)		2.0·10 ⁹			5
CH ₃ CH(OH)CH(OH)(OH) + HO [•] → 0.37 CH ₃ CH(OH)(OO [•]) + 0.37 CHO(OH) + 0.35 CH ₃ CH(OH)C(OH)(OH)(OO [•]) + 0.28 CH ₃ C(OH)(OO [•])CH(OH)(OH) + H ₂ O - O ₂	R(414)	1.3 10 ⁹			13
Pathway 1: CH ₃ CH(OH)CH(OH)(OH) + NO ₃ [•] → CH ₃ CH(OH)C [•] (OH)(OH) + NO ₃ ⁻ + H ⁺		6.2 10 ⁵			BR: 56%
CH ₃ CH(OH)C [•] (OH)(OH) + O ₂ → CH ₃ CH(OH)C(OH)(OH)(OO [•])		2.0·10 ⁹			5
Pathway 2: CH ₃ CH(OH)CH(OH)(OH) + NO ₃ [•] → CH ₃ C [•] (OH)CH(OH)(OH) + NO ₃ ⁻ + H ⁺		4.8 10 ⁵			BR: 44%
CH ₃ C [•] (OH)CH(OH)(OH) + O ₂ → CH ₃ C(OH)(OO [•])CH(OH)(OH)		2.0·10 ⁹			5
CH ₃ CH(OH)CH(OH)(OH) + NO ₃ [•] → 0.56 CH ₃ CH(OH)C(OH)(OH)(OO [•]) + 0.44 CH ₃ C(OH)(OO [•])CH(OH)(OH) + NO ₃ ⁻ + H ⁺ - O ₂	R(415)	1.1 10 ⁶			= k(CH ₂ (OH)CH(OH)(OH) + NO ₃ [•]) - 2
CH ₃ CH(OH)C(OH)(OH)(OO [•]) + OH ⁻ → CH ₃ CH(OH)C(OH)(O ⁻)(OO [•]) + H ₂ O		4.0 10 ⁹			
CH ₃ CH(OH)C(OH)(O ⁻)(OO [•]) → CH ₃ CH(OH)CO(OH) + O ₂ ^{•-}					9
CH ₃ CH(OH)C(OH)(OH)(OO [•]) + OH ⁻ → CH ₃ CH(OH)CO(OH) + O ₂ ^{•-} + H ₂ O	R(416)	4.0 10 ⁹			= k(CH ₃ CH(OH)(OO [•]) + OH ⁻)
CH ₃ CH(OH)C(OH)(OH)(OO [•]) → CH ₃ CH(OH)CO(OH) + HO ₂ [•]	R(417)	1.0 10 ⁶			15
CH ₃ C(OH)(OO [•])CH(OH)(OH) + OH ⁻ → CH ₃ C(O ⁻)(OO [•])CH(OH)(OH) + H ₂ O		4.0 10 ⁹			
CH ₃ C(O ⁻)(OO [•])CH(OH)(OH) → CH ₃ COCH(OH)(OH) + O ₂ ^{•-}					9
CH ₃ C(OH)(OO [•])CH(OH)(OH) + OH ⁻ → CH ₃ COCH(OH)(OH) + O ₂ ^{•-} + H ₂ O	R(418)	4.0 10 ⁹			= k(CH ₃ CH(OH)(OO [•]) + OH ⁻)
CH ₃ C(OH)(OO [•])CH(OH)(OH) → CH ₃ COCH(OH)(OH) + HO ₂ [•]	R(419)	1.9 10 ²			10
Oxidation of propanedial					19
Pathway 1: CH(OH)(OH)CH ₂ CH(OH)(OH) + HO [•] → CH(OH)(OH)CH ₂ C [•] (OH)(OH) + H ₂ O		1.7 10 ⁹			BR: 67% - 20
CH(OH)(OH)CH ₂ C [•] (OH)(OH) + O ₂ → CH(OH)(OH)CH ₂ C(OH)(OH)(OO [•])		2.0·10 ⁹			5
Pathway 2: CH(OH)(OH)CH ₂ CH(OH)(OH) + HO [•] → CH(OH)(OH)CH ₂ CH(OH)(O [•]) + H ₂ O		8.0 10 ⁸			BR: 33% - 20
CH(OH)(OH)CH ₂ CH(OH)(O [•]) → CH(OH)(OH)CH ₂ C [•] (OH)(OH)					4
CH(OH)(OH)CH ₂ C [•] (OH)(OH) + O ₂ → CH(OH)(OH)CH ₂ C(OH)(OH)(OO [•])		2.0·10 ⁹			5
CH(OH)(OH)CH ₂ CH(OH)(OH) + HO [•] → CH(OH)(OH)CH ₂ C(OH)(OH)(OO [•]) + H ₂ O - O ₂	R(420)	2.5 10 ⁹			13
CH(OH)(OH)CH ₂ CH(OH)(OH) + NO ₃ [•] → CH(OH)(OH)CH ₂ C [•] (OH)(OH) + NO ₃ ⁻ + H ⁺		1.0 10 ⁶			BR: 100%
CH(OH)(OH)CH ₂ C [•] (OH)(OH) + O ₂ → CH(OH)(OH)CH ₂ C(OH)(OH)(OO [•])		2.0·10 ⁹			5
CH(OH)(OH)CH ₂ CH(OH)(OH) + NO ₃ [•] → CH(OH)(OH)CH ₂ C(OH)(OH)(OO [•]) + NO ₃ ⁻ + H ⁺	R(421)	1.0 10 ⁶			= k(CH(OH)(OH)CH(OH)(OH) + NO ₃ [•]) - 2
CH(OH)(OH)CH ₂ C(OH)(OH)(OO [•]) + OH ⁻ → CH(OH)(OH)CH ₂ C(OH)(O ⁻)(OO [•]) + H ₂ O		4.0 10 ⁹			
CH(OH)(OH)CH ₂ C(OH)(O ⁻)(OO [•]) → CO(OH)CH ₂ CH(OH)(OH) + O ₂ ^{•-}					9
CH(OH)(OH)CH ₂ C(OH)(OH)(OO [•]) + OH ⁻ → CO(OH)CH ₂ CH(OH)(OH) + O ₂ ^{•-} + H ₂ O	R(422)	4.0 10 ⁹			= k(CH ₃ CH(OH)(OO [•]) + OH ⁻)

Reactions		k ₂₉₈ (M ⁻ⁿ⁺¹ s ⁻¹)	Ea/R (K)	References	Notes
CH(OH)(OH)CH ₂ C(OH)(OH)(OO•) → CO(OH)CH ₂ CH(OH)(OH) + HO ₂ •	R(423)	1.0 10 ⁶			15
Oxidation of 3-hydroxypropionic acid					
Pathway 1: CH ₂ (OH)CH ₂ CO(OH) + HO• → C•H(OH)CH ₂ CO(OH) + H ₂ O C•H(OH)CH ₂ CO(OH) + O ₂ → CH(OH)(OO•)CH ₂ CO(OH)		1.3 10 ⁹ 2.0·10 ⁹			BR: 100% - 21 5
CH ₂ (OH)CH ₂ CO(OH) + HO• → CH(OH)(OO•)CH ₂ CO(OH) + H ₂ O - O ₂ Pathway 1: CH ₂ (OH)CH ₂ CO(OH) + NO ₃ • → C•H(OH)CH ₂ CO(OH) + NO ₃ ⁻ + H ⁺ C•H(OH)CH ₂ CO(OH) + O ₂ → CH(OH)(OO•)CH ₂ CO(OH)	R(424)	1.3 10 ⁹ 2.1 10 ⁶ 2.0·10 ⁹			13 BR: 100% 5
CH ₂ (OH)CH ₂ CO(OH) + NO ₃ • → CH(OH)(OO•)CH ₂ CO(OH) + NO ₃ ⁻ + H ⁺ - O ₂	R(425)	2.1·10 ⁶	3248		= k(CH ₃ CH(OH)CO(OH) + NO ₃ •) - 2 BR: 100% - 22 5
Pathway 1: CH ₂ (OH)CH ₂ CO(O•) + HO• → C•H(OH)CH ₂ CO(O•) + H ₂ O C•H(OH)CH ₂ CO(O•) + O ₂ → CH(OH)(OO•)CH ₂ CO(O•)		2.5 10 ⁹ 2.0·10 ⁹			13
CH ₂ (OH)CH ₂ CO(O•) + HO• → CH(OH)(OO•)CH ₂ CO(O•) + H ₂ O - O ₂ Pathway 1: CH ₂ (OH)CH ₂ CO(O•) + NO ₃ • → C•H(OH)CH ₂ CO(O•) + NO ₃ ⁻ + H ⁺ C•H(OH)CH ₂ CO(O•) + O ₂ → CH(OH)(OO•)CH ₂ CO(O•)	R(426)	2.5 10 ⁹ 1.0 10 ⁷ 2.0·10 ⁹			BR: 100% 5
CH ₂ (OH)CH ₂ CO(O•) + NO ₃ • → CH(OH)(OO•)CH ₂ CO(O•) + NO ₃ ⁻ + H ⁺ - O ₂	R(427)	1.0·10 ⁷	2646		= k(CH ₃ CH(OH)CO(O•) + NO ₃ •) - 2
CH(OH)(OO•)CH ₂ CO(OH) + OH• → CH(O•)(OO•)CH ₂ CO(OH) + H ₂ O CH(O•)(OO•)CH ₂ CO(OH) → CO(OH)CH ₂ CHO + O ₂ •		4.0 10 ⁹			9
CH(OH)(OO•)CH ₂ CO(OH) + OH• → CO(OH)CH ₂ CHO + O ₂ •• + H ₂ O	R(428)	4.0 10 ⁹			= k(CH ₃ CH(OH)(OO•) + OH•)
CH(OH)(OO•)CH ₂ CO(OH) → CO(OH)CH ₂ CHO + HO ₂ •	R(429)	52 4.0 10 ⁹			10
CH(OH)(OO•)CH ₂ CO(O•) + OH• → CH(O•)(OO•)CH ₂ CO(O•) + H ₂ O CH(O•)(OO•)CH ₂ CO(O•) → CO(O•)CH ₂ CHO + O ₂ •					9
CH(OH)(OO•)CH ₂ CO(O•) + OH• → CO(O•)CH ₂ CHO + O ₂ •• + H ₂ O	R(430)	4.0 10 ⁹			= k(CH ₃ CH(OH)(OO•) + OH•)
CH(OH)(OO•)CH ₂ CO(O•) → CO(O•)CH ₂ CHO + HO ₂ •	R(431)	52			10
Oxidation of 2-propanol					
Pathway 1: CH ₃ CH(OH)CH ₃ + HO• → CH ₃ CH(OH)C•H ₂ + H ₂ O CH ₃ CH(OH)C•H ₂ + O ₂ → CH ₃ CH(OH)CH ₂ (OO•)		2.0 10 ⁸ 4.2·10 ⁹		Padmaja et al., 1993 Schaefer et al., 2014	BR: 13% - 23
Pathway 2: CH ₃ CH(OH)CH ₃ + HO• → CH ₃ C•(OH)CH ₃ + H ₂ O CH ₃ C•(OH)CH ₃ + O ₂ → CH ₃ C(OH)(OO•)CH ₃		1.7 10 ⁹ 4.2·10 ⁹			BR: 87% - 23
CH ₃ CH(OH)CH ₃ + HO• → 0.87 CH ₃ C(OH)(OO•)CH ₃ + 0.13 CH ₃ CH(OH)CH ₂ (OO•) + H ₂ O - O ₂ Pathway 1: CH ₃ CH(OH)CH ₃ + NO ₃ • → CH ₃ CH(OH)C•H ₂ + NO ₃ ⁻ + H ⁺ CH ₃ CH(OH)C•H ₂ + O ₂ → CH ₃ CH(OH)CH ₂ (OO•)	R(432)	1.9·10 ⁹ 5.0 10 ⁵ 4.2·10 ⁹		Schaefer et al., 2014 Monod et al., 2005	BR: 13%
Pathway 2: CH ₃ CH(OH)CH ₃ + NO ₃ • → CH ₃ C•(OH)CH ₃ + NO ₃ ⁻ + H ⁺ CH ₃ C•(OH)CH ₃ + O ₂ → CH ₃ C(OH)(OO•)CH ₃		3.2 10 ⁶ 4.2·10 ⁹		Schaefer et al., 2014	BR: 87%
CH ₃ CH(OH)CH ₃ + NO ₃ • → 0.87 CH ₃ C(OH)(OO•)CH ₃ + 0.13 CH ₃ CH(OH)CH ₂ (OO•) + NO ₃ ⁻ + H ⁺ - O ₂ CH ₃ C(OH)(OO•)CH ₃ + OH• → CH ₃ C(O•)(OO•)CH ₃ + H ₂ O CH ₃ C(O•)(OO•)CH ₃ → CH ₃ COCH ₃ + O ₂ •	R(433)	3.7·10 ⁶ 5.2 10 ⁹		Herrmann et al., 1994	2 9
CH ₃ C(OH)(OO•)CH ₃ + OH• → CH ₃ COCH ₃ + O ₂ •• + H ₂ O CH ₃ C(OH)(OO•)CH ₃ → CH ₃ COCH ₃ + HO ₂ •	R(434) R(435)	5.2 10 ⁹ 7.0 10 ²		Ilán et al., 1976 Ilán et al., 1976	
Pathway 1: 2 CH ₃ CH(OH)CH ₂ (OO•) → 2 CH ₃ CH(OH)CHO + H ₂ O ₂ Pathway 2: 2 CH ₃ CH(OH)CH ₂ (OO•) → CH ₃ CH(OH)CH ₂ (OH) + CH ₃ CH(OH)CHO + O ₂ Pathway 3: 2 CH ₃ CH(OH)CH ₂ (OO•) → 2 CH ₃ CH(OH)CH ₂ (O•) + O ₂ CH ₃ CH(OH)CH ₂ (O•) → CH ₂ O + CH ₃ C•H(OH) CH ₃ C•H(OH) + O ₂ → CH ₃ CH(OH)(OO•)		5.0 10 ⁷ 3.3 10 ⁷ 1.7 10 ⁷			BR: 50% BR: 33% BR: 17% 6 - 7 5
2 CH ₃ CH(OH)CH ₂ (OO•) → 1.33 CH ₃ CH(OH)CHO + 0.33 CH ₃ CH(OH)CH ₂ (OH) + 0.34 CH ₃ CH(OH)(OO•) + 0.34 CH ₂ O + 0.50 H ₂ O ₂ + 0.16 O ₂	R(436)	2.0·10 ⁹ 1.0 10 ⁸			= k(2 CH ₂ (OH)CH ₂ (OO•)) - 8

Reactions	k ₂₉₈ (M ⁻ⁿ⁺¹ s ⁻¹)	Ea/R (K)	References	Notes
Oxidation of propionaldehyde - ethylperoxide formation				24
Pathway 1: CH ₃ CH ₂ CHO + HO• → CH ₃ CH ₂ C•O + H ₂ O CH ₃ CH ₂ C•O + O ₂ → CH ₃ CH ₂ CO(OO•)	2.6 10 ⁹ 2.0·10 ⁹			BR: 87% - 25 5
Pathway 2: CH ₃ CH ₂ CHO + HO• → C•H ₂ CH ₂ CHO + H ₂ O C•H ₂ CH ₂ CHO + O ₂ → CHOCH ₂ CH ₂ (OO•)	4.0 10 ⁸ 2.0·10 ⁹			BR: 13% - 25 5
CH ₃ CH ₂ CHO + HO• → 0.87 CH ₃ CH ₂ CO(OO•) + 0.13 CHOCH ₂ CH ₂ (OO•) + H ₂ O - O ₂	R(437) 3.2·10 ⁹			13
Pathway 1: CH ₃ CH ₂ CHO + NO ₃ • → CH ₃ CH ₂ C•O + NO ₃ ⁻ + H ⁺ CH ₃ CH ₂ C•O + O ₂ → CH ₃ CH ₂ CO(OO•)	5.0 10 ⁷ 2.0·10 ⁹			BR: 87% 5
Pathway 2: CH ₃ CH ₂ CHO + NO ₃ • → C•H ₂ CH ₂ CHO + NO ₃ ⁻ + H ⁺ C•H ₂ CH ₂ CHO + O ₂ → CHOCH ₂ CH ₂ (OO•)	8.0 10 ⁶ 2.0·10 ⁹			BR: 13% 5
CH ₃ CH ₂ CHO + NO ₃ • → 0.87 CH ₃ CH ₂ CO(OO•) + 0.13 CHOCH ₂ CH ₂ (OO•) + NO ₃ ⁻ + H ⁺ - O ₂	R(438) 5.8·10 ⁷	2646	De Semainville et al., 2007	2
Pathway 1: CH ₃ CH ₂ CH(OH)(OH) + HO• → CH ₃ CH ₂ C•(OH)(OH) + H ₂ O CH ₃ CH ₂ C•(OH)(OH) + O ₂ → CH ₃ CH ₂ C(OH)(OH)(OO•)	9.9 10 ⁸ 2.0·10 ⁹			BR: 52% - 26 5
Pathway 2: CH ₃ CH ₂ CH(OH)(OH) + HO• → C•H ₂ CH ₂ CH(OH)(OH) + H ₂ O C•H ₂ CH ₂ CH(OH)(OH) + O ₂ → CH(OH)(OH)CH ₂ CH ₂ (OO•)	4.8 10 ⁸ 2.0·10 ⁹			BR: 25% - 26 5
Pathway 3: CH ₃ CH ₂ CH(OH)(OH) + HO• → CH ₃ CH ₂ CH(OH)(O•) + H ₂ O CH ₃ CH ₂ CH(OH)(O•) → CH ₃ CH ₂ C•(OH)(OH) CH ₃ CH ₂ C•(OH)(OH) + O ₂ → CH ₃ CH ₂ C(OH)(OH)(OO•)	4.3 10 ⁸ 2.0·10 ⁹			BR: 23% - 26 4 5
CH ₃ CH ₂ CH(OH)(OH) + HO• → 0.75 CH ₃ CH ₂ C(OH)(OH)(OO•) + 0.25 CH(OH)(OH)CH ₂ CH ₂ (OO•) + H ₂ O - O ₂	R(439) 1.9·10 ⁹			13
Pathway 1: CH ₃ CH ₂ CH(OH)(OH) + NO ₃ • → CH ₃ CH ₂ C•(OH)(OH) + NO ₃ ⁻ + H ⁺ CH ₃ CH ₂ C•(OH)(OH) + O ₂ → CH ₃ CH ₂ C(OH)(OH)(OO•)	3.9 10 ⁷ 2.0·10 ⁹			BR: 67% 5
Pathway 2: CH ₃ CH ₂ CH(OH)(OH) + NO ₃ • → C•H ₂ CH ₂ CH(OH)(OH) + NO ₃ ⁻ + H ⁺ C•H ₂ CH ₂ CH(OH)(OH) + O ₂ → CH(OH)(OH)CH ₂ CH ₂ (OO•)	1.9 10 ⁷ 2.0·10 ⁹			BR: 33% 5
CH ₃ CH ₂ CH(OH)(OH) + NO ₃ • → 0.67 CH ₃ CH ₂ C(OH)(OH)(OO•) + 0.33 CH(OH)(OH)CH ₂ CH ₂ (OO•) + NO ₃ ⁻ + H ⁺ - O ₂	R(440) 5.8·10 ⁷	2646	De Semainville et al., 2007	2
2 CH ₃ CH ₂ CO(OO•) → 2 CH ₃ CH ₂ CO(O•)· + O ₂ CH ₃ CH ₂ CO(O•)· → CH ₃ CH ₂ C• + CO ₂ CH ₃ CH ₂ C• + O ₂ → CH ₃ CH ₂ (OO•)	1.6·10 ⁸ 2.0·10 ⁹			6 - 7 5
2 CH ₃ CH ₂ CO(OO•) → 2 CH ₃ CH ₂ (OO•) + 2 CO ₂ - O ₂ Pathway 1: 2 CHOCH ₂ CH ₂ (OO•) → 2 CHOCH ₂ CHO + H ₂ O ₂ Pathway 2: 2 CHOCH ₂ CH ₂ (OO•) → 2 CHOCH ₂ CH ₂ (O•) + O ₂ CHOCH ₂ CH ₂ (O•) → CHOCH ₂ C•H(OH) CHOCH ₂ C•H(OH) + O ₂ → CHOCH ₂ CH(OH)(OO•)	R(441) 1.6·10 ⁸ 3.0 10 ⁷ 1.3 10 ⁸	-1600		= k(2 CH ₃ CH ₂ (OO•)) - 8 BR: 20% - 3 BR: 80% - 3 4 5
2 CHOCH ₂ CH ₂ (OO•) → 0.40 CHOCH ₂ CHO + 1.60 CHOCH ₂ CH(OH)(OO•) + 0.20 H ₂ O ₂ - 0.80 O ₂ CH ₃ CH ₂ C(OH)(OH)(OO•) + OH ⁻ → CH ₃ CH ₂ C(OH)(O ⁻)(OO•) + H ₂ O CH ₃ CH ₂ C(OH)(O ⁻)(OO•) → CH ₃ CH ₂ CO(OH) + O ₂ ⁻	R(442) 1.6 10 ⁸ 4.0 10 ⁹	-1600		= k(2 CH ₃ CH ₂ (OO•)) - 8 9
CH ₃ CH ₂ C(OH)(OH)(OO•) + OH ⁻ → CH ₃ CH ₂ CO(OH) + O ₂ ⁻ + H ₂ O	R(443) 4.0 10 ⁹			= k(CH ₃ CH(OH)(OO•) + OH ⁻)
CH ₃ CH ₂ C(OH)(OH)(OO•) → CH ₃ CH ₂ CO(OH) + HO ₂ •	R(444) 1.0 10 ⁶			15
Pathway 1: 2 CH(OH)(OH)CH ₂ CH ₂ (OO•) → 2 CHOCH ₂ CH(OH)(OH) + H ₂ O ₂ Pathway 2: 2 CH(OH)(OH)CH ₂ CH ₂ (OO•) → 2 CH(OH)(OH)CH ₂ CH ₂ (O•) + O ₂ CH(OH)(OH)CH ₂ CH ₂ (O•) → CH(OH)(OH)CH ₂ C•H(OH) CH(OH)(OH)CH ₂ C•H(OH) + O ₂ → CH(OH)(OH)CH ₂ CH(OH)(OO•)	3.0 10 ⁷ 1.3 10 ⁸			BR: 20% - 3 BR: 80% - 3 4 5
2 CH(OH)(OH)CH ₂ CH ₂ (OO•) → 0.40 CHOCH ₂ CH(OH)(OH) + 1.60 CH(OH)(OH)CH ₂ CH(OH)(OO•) + 0.20 H ₂ O ₂ - 0.80 O ₂	R(445) 1.6 10 ⁸	-1600		= k(2 CH ₃ CH ₂ (OO•)) - 8
Oxidation of methylglyoxal				27
Pathway 1: CH ₃ COCH(OH)(OH) + HO• → C•H ₂ COCH(OH)(OH) + H ₂ O	1.2 10 ⁸			BR: 14% - 28

Reactions		k_{298} ($M^{-n+1} s^{-1}$)	Ea/R (K)	References	Notes
$C^*H_2COCH(OH)(OH) + O_2 \rightarrow CH(OH)(OH)COCH_2(OO^*)$		$2.0 \cdot 10^9$			5
Pathway 2: $CH_3COCH(OH)(OH) + HO^* \rightarrow CH_3COC^*(OH)(OH) + H_2O$		$2.7 \cdot 10^8$			BR: 29% - 28
$CH_3COC^*(OH)(OH) + O_2 \rightarrow CH_3COC(OH)(OH)(OO^*)$		$2.0 \cdot 10^9$			5
Pathway 3: $CH_3COCH(OH)(OH) + HO^* \rightarrow CH_3COCH(OH)(O^*) + H_2O$		$5.3 \cdot 10^8$			BR: 57% - 28
$CH_3COCH(OH)(O^*) \rightarrow CH_3C^*O + CHO(OH)$					6 - 7
$CH_3C^*O + O_2 \rightarrow CH_3CO(OO^*)$		$2.0 \cdot 10^9$			5
$CH_3COCH(OH)(OH) + HO^* \rightarrow 0.14 CH(OH)(OH)COCH_2(OO^*) + 0.29 CH_3COC(OH)(OH)(OO^*) + 0.57 CHO(OH) + 0.57 CH_3CO(OO^*) + H_2O - O_2$	R(446)	$9.2 \cdot 10^8$	1235	Schaefer et al., 2014	
Pathway 1: $CH_3COCH(OH)(OH) + NO_3^* \rightarrow C^*H_2COCH(OH)(OH) + NO_3^- + H^+$		$1.5 \cdot 10^6$			BR: 33%
$C^*H_2COCH(OH)(OH) + O_2 \rightarrow CH(OH)(OH)COCH_2(OO^*)$		$2.0 \cdot 10^9$			5
Pathway 2: $CH_3COCH(OH)(OH) + NO_3^* \rightarrow CH_3COC^*(OH)(OH) + NO_3^- + H^+$		$3.0 \cdot 10^6$			BR: 67%
$CH_3COC^*(OH)(OH) + O_2 \rightarrow CH_3COC(OH)(OH)(OO^*)$		$2.0 \cdot 10^9$			5
$CH_3COCH(OH)(OH) + NO_3^* \rightarrow 0.33 CH(OH)(OH)COCH_2(OO^*) + 0.67 CH_3COC(OH)(OH)(OO^*) + NO_3^- + H^+ - O_2$	R(447)	$4.5 \cdot 10^6$	4213	Schaefer et al., 2014	2
Pathway 1: $CH_3C(OH)(OH)CH(OH)(OH) + HO^* \rightarrow CH_3C(OH)(O^*)CH(OH)(OH) + H_2O$		$4.9 \cdot 10^8$			BR: 53% - 29
$CH_3C(OH)(O^*)CH(OH)(OH) \rightarrow CH_3CO(OH) + C^*H(OH)(OH)$					6 - 7
$C^*H(OH)(OH) + O_2 \rightarrow CH(OH)(OH)(OO^*)$		$2.0 \cdot 10^9$			5
Pathway 2: $CH_3C(OH)(OH)CH(OH)(OH) + HO^* \rightarrow CH_3C(OH)(OH)CH(OH)(O^*) + H_2O$		$4.3 \cdot 10^8$			BR: 47% - 29
$CH_3C(OH)(OH)CH(OH)(O^*) \rightarrow CH_3C^*(OH)(OH) + CHO(OH)$					6 - 7
$CH_3C^*(OH)(OH) + O_2 \rightarrow CH_3C(OH)(OH)(OO^*)$		$2.0 \cdot 10^9$			5
$CH_3C(OH)(OH)CH(OH)(OH) + HO^* \rightarrow 0.53 CH_3CO(OH) + 0.53 CH(OH)(OH)(OO^*) + 0.47 CHO(OH) + 0.47 CH_3C(OH)(OH)(OO^*) + H_2O - O_2$	R(448)	$9.2 \cdot 10^8$	1235	Schaefer et al., 2014	
Pathway 1: $CH_3C(OH)(OH)CH(OH)(OH) + NO_3^* \rightarrow CH_3C(OH)(OH)C^*(OH)(OH) + NO_3^- + H^+$		$3.0 \cdot 10^6$			BR: 67%
$CH_3C(OH)(OH)C^*(OH)(OH) + O_2 \rightarrow CH_3C(OH)(OH)C(OH)(OH)(OO^*)$		$2.0 \cdot 10^9$			5
Pathway 2: $CH_3C(OH)(OH)CH(OH)(OH) + NO_3^* \rightarrow C^*H_2C(OH)(OH)CH(OH)(OH) + NO_3^- + H^+$		$1.5 \cdot 10^6$			BR: 33%
$C^*H_2C(OH)(OH)CH(OH)(OH) + O_2 \rightarrow CH(OH)(OH)C(OH)(OH)CH_2(OO^*)$		$2.0 \cdot 10^9$			5
$CH_3C(OH)(OH)CH(OH)(OH) + NO_3^* \rightarrow 0.67 CH_3C(OH)(OH)C(OH)(OH)(OO^*) + 0.33 CH(OH)(OH)C(OH)(OH)CH_2(OO^*) + NO_3^- + H^+ - O_2$	R(449)	$4.5 \cdot 10^6$	4213	Schaefer et al., 2014	2
$CH_3COC(OH)(OH)(OO^*) + OH^- \rightarrow CH_3COC(OH)(O^-)(OO^*) + H_2O$		$4.0 \cdot 10^9$			
$CH_3COC(OH)(O^-)(OO^*) \rightarrow CH_3COCO(OH) + O_2^{\bullet-}$	R(450)	$4.0 \cdot 10^9$			9
$CH_3COC(OH)(OH)(OO^*) + OH^- \rightarrow CH_3COCO(OH) + O_2^{\bullet-} + H_2O$	R(451)	$1.0 \cdot 10^6$			= $k(CH_3CH(OH)(OO^*) + OH^-)$
$CH_3COC(OH)(OH)(OO^*) \rightarrow CH_3COCO(OH) + HO_2^*$		$4.0 \cdot 10^9$			15
$CH_3C(OH)(OH)C(OH)(OH)(OO^*) + OH^- \rightarrow CH_3C(OH)(OH)C(OH)(O^-)(OO^*) + H_2O$					
$CH_3C(OH)(OH)C(OH)(O^-)(OO^*) \rightarrow CH_3C(OH)(OH)CO(OH) + O_2^{\bullet-}$					9
$CH_3C(OH)(OH)C(OH)(OH)(OO^*) + OH^- \rightarrow CH_3C(OH)(OH)CO(OH) + O_2^{\bullet-} + H_2O$	R(452)	$4.0 \cdot 10^9$			
$CH_3C(OH)(OH)C(OH)(OH)(OO^*) \rightarrow CH_3C(OH)(OH)CO(OH) + HO_2^*$	R(453)	$1.0 \cdot 10^6$			15
Pathway 1: $2 CH(OH)(OH)COCH_2(OO^*) \rightarrow 2 CHOCOCH(OH)(OH) + H_2O_2$		$1.8 \cdot 10^8$			BR: 45%
Pathway 2: $2 CH(OH)(OH)COCH_2(OO^*) \rightarrow CH_2(OH)COCH(OH)(OH) + CHOCOCH(OH)(OH) + O_2$		$8.0 \cdot 10^7$			BR: 20%
Pathway 3: $2 CH(OH)(OH)COCH_2(OO^*) \rightarrow 2 CH_2(O^*)COCH(OH)(OH) + O_2$		$1.4 \cdot 10^8$			BR: 35%
$CH_2(O^*)COCH(OH)(OH) \rightarrow C^*H(OH)COCH(OH)(OH)$					4
$C^*H(OH)COCH(OH)(OH) + O_2 \rightarrow CH(OH)(OH)COCH(OH)(OO^*)$		$2.0 \cdot 10^9$			5
$2 CH(OH)(OH)COCH_2(OO^*) \rightarrow 1.10 CHOCOCH(OH)(OH) + 0.20 CH_2(OH)COCH(OH)(OH) + 0.70 CH(OH)(OH)COCH(OH)(OO^*) + 0.45 H_2O_2 - 0.15 O_2$	R(454)	$4.0 \cdot 10^8$			= $k(2 CH_3COCH_2(OO^*)) - 8$
Pathway 1: $2 CH(OH)(OH)C(OH)(OH)CH_2(OO^*) \rightarrow 2 CHOC(OH)(OH)CH(OH)(OH) + H_2O_2$		$1.8 \cdot 10^8$			BR: 45%
Pathway 2: $2 CH(OH)(OH)C(OH)(OH)CH_2(OO^*) \rightarrow CH_2(OH)C(OH)(OH)CH(OH)(OH) + CHOC(OH)(OH)CH(OH)(OH) + O_2$		$8.0 \cdot 10^7$			BR: 20%
Pathway 3: $2 CH(OH)(OH)C(OH)(OH)CH_2(OO^*) \rightarrow 2 CH_2(O^*)C(OH)(OH)CH(OH)(OH) + O_2$		$1.4 \cdot 10^8$			BR: 35%
$CH_2(O^*)C(OH)(OH)CH(OH)(OH) \rightarrow C^*H(OH)C(OH)(OH)CH(OH)(OH)$					4
$C^*H(OH)C(OH)(OH)CH(OH)(OH) + O_2 \rightarrow CH(OH)(OH)C(OH)(OH)CH(OH)(OO^*)$		$2.0 \cdot 10^9$			5

Reactions		k_{298} ($M^{-n+1} s^{-1}$)	Ea/R (K)	References	Notes
2 CH(OH)(OH)C(OH)(OH)CH ₂ (OO•) → 1.10 CHOC(OH)(OH)CH(OH)(OH) + 0.20 CH ₂ (OH)C(OH)(OH)CH(OH)(OH) + 0.70 CH(OH)(OH)C(OH)(OH)CH(OH)(OO•) + 0.45 H ₂ O ₂ - 0.15 O ₂	R(455)	4.0 10 ⁸			= k(2 CH ₃ COCH ₂ (OO•)) - 8
Oxidation of 2-oxo, 3-hydroxypropanal					30
Pathway 1: CH ₂ (OH)COCH(OH)(OH) + HO• → CH ₂ (OH)COCH(OH)(O•) + H ₂ O CH ₂ (OH)COCH(OH)(O•) → CH ₂ (OH)C•O + CHO(OH) CH ₂ (OH)C•O + O ₂ → CH ₂ (OH)CO(OO•)		3.6 10 ⁸ 2.0·10 ⁹			BR: 41% - 31 6 - 7 5
Pathway 2: CH ₂ (OH)COCH(OH)(OH) + HO• → C•H(OH)COCH(OH)(OH) + H ₂ O C•H(OH)COCH(OH)(OH) + O ₂ → CH(OH)(OH)COCH(OH)(OO•)		3.5 10 ⁸ 2.0·10 ⁹			BR: 40% - 31 5
Pathway 3: CH ₂ (OH)COCH(OH)(OH) + HO• → CH ₂ (OH)COC•(OH)(OH) + H ₂ O CH ₂ (OH)COC•(OH)(OH) + O ₂ → CH ₂ (OH)COC(OH)(OH)(OO•)		1.6 10 ⁸ 2.0·10 ⁹			BR: 19% - 31 5
CH ₂ (OH)COCH(OH)(OH) + HO• → 0.41 CHO(OH) + 0.41 CH ₂ (OH)CO(OO•) + 0.40 CH(OH)(OH)COCH(OH)(OO•) + 0.19 CH ₂ (OH)COC(OH)(OH)(OO•) + H ₂ O - O ₂	R(456)	8.7 10 ⁸			13
Pathway 1: CH ₂ (OH)COCH(OH)(OH) + NO ₃ • → C•H(OH)COCH(OH)(OH) + NO ₃ ⁻ + H ⁺ C•H(OH)COCH(OH)(OH) + O ₂ → CH(OH)(OH)COCH(OH)(OO•)		7.4 10 ⁵ 2.0·10 ⁹			BR: 67% 5
Pathway 2: CH ₂ (OH)COCH(OH)(OH) + NO ₃ • → CH ₂ (OH)COC•(OH)(OH) + NO ₃ ⁻ + H ⁺ CH ₂ (OH)COC•(OH)(OH) + O ₂ → CH ₂ (OH)COC(OH)(OH)(OO•)		3.6 10 ⁵ 2.0·10 ⁹			BR: 33% 5
CH ₂ (OH)COCH(OH)(OH) + NO ₃ • → 0.67 CH(OH)(OH)COCH(OH)(OO•) + 0.33 CH ₂ (OH)COC(OH)(OH)(OO•) + NO ₃ ⁻ + H ⁺ - O ₂	R(457)	1.1 10 ⁶			= k(CH ₂ (OH)CH(OH)(OH) + NO ₃ •) - 2
Pathway 1: CH ₂ (OH)C(OH)(OH)CH(OH)(OH) + HO• → CH ₂ (OH)C(OH)(O•)CH(OH)(OH) + H ₂ O CH ₂ (OH)C(OH)(O•)CH(OH)(OH) → CH ₂ (OH)CO(OH) + C•H(OH)(OH) C•H(OH)(OH) + O ₂ → CH(OH)(OH)(OO•)		4.6 10 ⁸ 2.0·10 ⁹			BR: 38% - 32 6 - 7 5
Pathway 2: CH ₂ (OH)C(OH)(OH)CH(OH)(OH) + HO• → CH ₂ (OH)C(OH)(OH)CH(OH)(O•) + H ₂ O CH ₂ (OH)C(OH)(OH)CH(OH)(O•) → CH ₂ (OH)C•(OH)(OH) + CHO(OH) CH ₂ (OH)C•(OH)(OH) + O ₂ → CH ₂ (OH)C(OH)(OH)(OO•)		4.2 10 ⁸ 2.0·10 ⁹			BR: 35% - 32 6 - 7 5
Pathway 3: CH ₂ (OH)C(OH)(OH)CH(OH)(OH) + HO• → C•H(OH)C(OH)(OH)CH(OH)(OH) + H ₂ O C•H(OH)C(OH)(OH)CH(OH)(OH) + O ₂ → CH(OH)(OH)C(OH)(OH)CH(OH)(OO•)		3.2 10 ⁸ 2.0·10 ⁹			BR: 27% - 32 5
CH ₂ (OH)C(OH)(OH)CH(OH)(OH) + HO• → 0.38 CH ₂ (OH)CO(OH) + 0.38 CH(OH)(OH)(OO•) + 0.35 CHO(OH) + 0.35 CH ₂ (OH)C(OH)(OH)(OO•) + 0.27 CH(OH)(OH)C(OH)(OH)CH(OH)(OO•) + H ₂ O - O ₂	R(458)	1.2 10 ⁹			13
Pathway 1: CH ₂ (OH)C(OH)(OH)CH(OH)(OH) + NO ₃ • → C•H(OH)C(OH)(OH)CH(OH)(OH) + NO ₃ ⁻ + H ⁺ C•H(OH)C(OH)(OH)CH(OH)(OH) + O ₂ → CH(OH)(OH)C(OH)(OH)CH(OH)(OO•)		1.0 10 ⁶ 2.0·10 ⁹			BR: 100% 5
CH ₂ (OH)C(OH)(OH)CH(OH)(OH) + NO ₃ • → CH(OH)(OH)C(OH)(OH)CH(OH)(OO•) + NO ₃ ⁻ + H ⁺ - O ₂	R(459)	1.0 10 ⁶			= k(CH(OH)(OH)CH(OH)(OH) + NO ₃ •) - 2
CH ₂ (OH)COC(OH)(OH)(OO•) + OH• → CH ₂ (OH)COC(OH)(O•)(OO•) + H ₂ O CH ₂ (OH)COC(OH)(O•)(OO•) → CH ₂ (OH)COCO(OH) + O ₂ ⁻		4.0 10 ⁹			9
CH ₂ (OH)COC(OH)(OH)(OO•) + OH• → CH ₂ (OH)COCO(OH) + O ₂ ⁻ + H ₂ O	R(460)	4.0 10 ⁹			= k(CH ₃ CH(OH)(OO•) + OH•)
CH ₂ (OH)COC(OH)(OH)(OO•) → CH ₂ (OH)COCO(OH) + HO ₂ •	R(461)	1.0 10 ⁶			15
CH(OH)(OH)COCH(OH)(OO•) + OH• → CH(OH)(OH)COCH(O•)(OO•) + H ₂ O CH(OH)(OH)COCH(O•)(OO•) → CHOCOCH(OH)(OH) + O ₂ ⁻		4.0 10 ⁹			9
CH(OH)(OH)COCH(OH)(OO•) + OH• → CHOCOCH(OH)(OH) + O ₂ ⁻ + H ₂ O	R(462)	4.0 10 ⁹			= k(CH ₃ CH(OH)(OO•) + OH•)
CH(OH)(OH)COCH(OH)(OO•) → CHOCOCH(OH)(OH) + HO ₂ •	R(463)	1.9 10 ²			10
CH(OH)(OH)C(OH)(OH)CH(OH)(OO•) + OH• → CH(OH)(OH)C(OH)(OH)CH(O•)(OO•) + H ₂ O CH(OH)(OH)C(OH)(OH)CH(O•)(OO•) → CHOC(OH)(OH)CH(OH)(OH) + O ₂ ⁻		4.0 10 ⁹			9
CH(OH)(OH)C(OH)(OH)CH(OH)(OO•) + OH• → CHOC(OH)(OH)CH(OH)(OH) + O ₂ ⁻ + H ₂ O	R(464)	4.0 10 ⁹			
CH(OH)(OH)C(OH)(OH)CH(OH)(OO•) → CHOC(OH)(OH)CH(OH)(OH) + HO ₂ •	R(465)	1.0 10 ⁶			15
CH ₂ (OH)C(OH)(OH)C(OH)(OH)(OO•) + OH• → CH ₂ (OH)C(OH)(OH)C(OH)(O•)(OO•) + H ₂ O CH ₂ (OH)C(OH)(OH)C(OH)(O•)(OO•) → CH ₂ (OH)C(OH)(OH)CO(OH) + O ₂ ⁻		4.0 10 ⁹			9

Reactions		k ₂₉₈ (M ⁻ⁿ⁺¹ s ⁻¹)	Ea/R (K)	References	Notes
CH ₂ (OH)C(OH)(OH)C(OH)(OH)(OO•) + OH ⁻ → CH ₂ (OH)C(OH)(OH)CO(OH) + O ₂ • ⁻ + H ₂ O	R(466)	4.0 10 ⁹			= k(CH ₃ CH(OH)(OO•) + OH ⁻)
CH ₂ (OH)C(OH)(OH)C(OH)(OH)(OO•) → CH ₂ (OH)C(OH)(OH)CO(OH) + HO ₂ •	R(467)	1.0 10 ⁶			15
Oxidation of oxopropanedial					33
Pathway 1: CH(OH)(OH)C(OH)(OH)CH(OH)(OH) + HO• → CH(OH)(OH)C(OH)(OH)CH(OH)(O•) + H ₂ O		8.7 10 ⁸			BR: 67% - 34
CH(OH)(OH)C(OH)(OH)CH(OH)(O•) → CH(OH)(OH)C•(OH)(OH) + CHO(OH)					6 - 7
CH(OH)(OH)C•(OH)(OH) + O ₂ → CH(OH)(OH)C(OH)(OH)(OO•)		2.0·10 ⁹			5
Pathway 2: CH(OH)(OH)C(OH)(OH)CH(OH)(OH) + HO• → CH(OH)(OH)C(OH)(O•)CH(OH)(OH) + H ₂ O		4.3 10 ⁸			BR: 33% - 34
CH(OH)(OH)C(OH)(O•)CH(OH)(OH) → C•H(OH)(OH) + CH(OH)(OH)CO(OH)					6 - 7
C•H(OH)(OH) + O ₂ → CH(OH)(OH)(OO•)		2.0·10 ⁹			5
CH(OH)(OH)C(OH)(OH)CH(OH)(OH) + HO• → 0.67 CH(OH)(OH)C(OH)(OH)(OO•) + 0.67 CHO(OH) + 0.33	R(468)	1.3 10 ⁹			13
CH(OH)(OH)CO(OH) + 0.33 CH(OH)(OH)(OO•) + H ₂ O - O ₂					
CH(OH)(OH)C(OH)(OH)C(OH)(OH)(OO•) + OH ⁻ → CH(OH)(OH)C(OH)(OH)C(OH)(O•)(OO•) + H ₂ O		4.0 10 ⁹			
CH(OH)(OH)C(OH)(OH)C(OH)(O•)(OO•) → CO(OH)C(OH)(OH)CH(OH)(OH) + O ₂ • ⁻					9
CH(OH)(OH)C(OH)(OH)C(OH)(OH)(OO•) + OH ⁻ → CO(OH)C(OH)(OH)CH(OH)(OH) + O ₂ • ⁻ + H ₂ O	R(469)	4.0 10 ⁹			= k(CH ₃ CH(OH)(OO•) + OH ⁻)
CH(OH)(OH)C(OH)(OH)C(OH)(OH)(OO•) → CO(OH)C(OH)(OH)CH(OH)(OH) + HO ₂ •	R(470)	1.0 10 ⁶			15
Oxidation of acetone					35
CH ₃ COCH ₃ + HO• → CH ₃ COC•H ₂ + H ₂ O		1.8·10 ⁸			
CH ₃ COC•H ₂ + O ₂ → CH ₃ COCH ₂ (OO•)		3.1 10 ⁹		Zegota et al., 1986a	
CH ₃ COCH ₃ + HO• → CH ₃ COCH ₂ (OO•) + H ₂ O - O ₂	R(471)	1.8·10 ⁸		Gligorovski et al., 2009	
CH ₃ COCH ₃ + NO ₃ • → CH ₃ COC•H ₂ + NO ₃ ⁻ + H ⁺		3.7 10 ³	4332		
CH ₃ COC•H ₂ + O ₂ → CH ₃ COCH ₂ (OO•)		3.1 10 ⁹		Zegota et al., 1986a	5
CH ₃ COCH ₃ + NO ₃ • → CH ₃ COCH ₂ (OO•) + NO ₃ ⁻ + H ⁺ - O ₂	R(472)	3.7·10 ³	4332	Herrmann and Zellner, 1998	
Pathway 1: 2 CH ₃ COCH ₂ (OO•) → 2 CH ₃ COCHO + H ₂ O ₂		1.8 10 ⁸		Zegota et al., 1986a	BR: 45%
Pathway 2: 2 CH ₃ COCH ₂ (OO•) → CH ₃ COCHO + CH ₃ COCH ₂ (OH) + O ₂		8.0 10 ⁷		Zegota et al., 1986a	BR: 20%
Pathway 3: 2 CH ₃ COCH ₂ (OO•) → 2 CH ₃ COCH ₂ (O•) + O ₂		1.4 10 ⁸		Zegota et al., 1986a	BR: 35%
CH ₃ COCH ₂ (O•) → CH ₃ COC•H(OH)					4
CH ₃ COC•H(OH) + O ₂ → CH ₃ COCH(OH)(OO•)		2.0 10 ⁹			5
2 CH ₃ COCH ₂ (OO•) → 1.10 CH ₃ COCHO + 0.20 CH ₃ COCH ₂ (OH) + 0.70 CH ₃ COCH(OH)(OO•) + 0.45 H ₂ O ₂ - 0.15 O ₂	R(473)	4.0 10 ⁸		Zegota et al., 1986a	
Oxidation of hydroxyacetone					36
Pathway 1: CH ₃ COCH ₂ (OH) + HO• → C•H ₂ COCH ₂ (OH) + H ₂ O		8.5 10 ⁷			BR: 16% - 37
C•H ₂ COCH ₂ (OH) + O ₂ → CH ₂ (OH)COCH ₂ (OO•)		2.0 10 ⁹			5
Pathway 2: CH ₃ COCH ₂ (OH) + HO• → CH ₃ COC•H(OH) + H ₂ O		3.5 10 ⁸			BR: 69% - 37
CH ₃ COC•H(OH) + O ₂ → CH ₃ COCH(OH)(OO•)		2.0 10 ⁹			5
Pathway 3: CH ₃ COCH ₂ (OH) + HO• → CH ₃ COCH ₂ (O•) + H ₂ O		7.5 10 ⁷			BR: 15% - 37
CH ₃ COCH ₂ (O•) → CH ₃ C•O + CH ₂ O					6 - 7
CH ₃ C•O + O ₂ → CH ₃ CO(OO•)		2.0 10 ⁹			5
CH ₃ COCH ₂ (OH) + HO• → 0.69 CH ₃ COCH(OH)(OO•) + 0.16 CH ₂ (OH)COCH ₂ (OO•) + 0.15 CH ₂ O + 0.15	R(474)	5.1 10 ⁸			13
CH ₃ CO(OO•) + H ₂ O - O ₂					
CH ₃ COCH ₂ (OH) + NO ₃ • → CH ₃ COC•H(OH) + NO ₃ ⁻ + H ⁺		1.8 10 ⁷			BR: 100%
CH ₃ COC•H(OH) + O ₂ → CH ₃ COCH(OH)(OO•)		2.0 10 ⁹			5
CH ₃ COCH ₂ (OH) + NO ₃ • → CH ₃ COCH(OH)(OO•) + NO ₃ ⁻ + H ⁺ - O ₂	R(475)	1.8·10 ⁷	1564	De Semainville et al., 2007	2
CH ₃ COCH(OH)(OO•) + OH ⁻ → CH ₃ COCH(O•)(OO•) + H ₂ O		4.0 10 ⁹			
CH ₃ COCH(O•)(OO•) → CH ₃ COCHO + O ₂ • ⁻					9

Reactions		k ₂₉₈ (M ⁻ⁿ⁺¹ s ⁻¹)	Ea/R (K)	References	Notes
CH ₃ COCH(OH)(OO•) + OH ⁻ → CH ₃ COCHO + O ₂ ^{•-} + H ₂ O	R(476)	4.0 10 ⁹			= k(CH ₃ CH(OH)(OO•) + OH ⁻)
CH ₃ COCH(OH)(OO•) → CH ₃ COCHO + HO ₂ [•]	R(477)	1.9 10 ²			10
Pathway 1: 2 CH ₂ (OH)COCH ₂ (OO•) → 2 CH ₂ (OH)COCHO + H ₂ O ₂		1.8 10 ⁸			BR: 45%
Pathway 2: 2 CH ₂ (OH)COCH ₂ (OO•) → CH ₂ (OH)COCH ₂ (OH) + CH ₂ (OH)COCHO + O ₂		8.0 10 ⁷			BR: 20%
Pathway 3: 2 CH ₂ (OH)COCH ₂ (OO•) → 2 CH ₂ (OH)COCH ₂ (O•) + O ₂		1.4 10 ⁸			BR: 35%
CH ₂ (OH)COCH ₂ (O•) → CH ₂ (OH)COC•H(OH)					4
CH ₂ (OH)COC•H(OH) + O ₂ → CH ₂ (OH)COCH(OH)(OO•)		2.0 10 ⁹			5
2 CH ₂ (OH)COCH ₂ (OO•) → 1.10 CH ₂ (OH)COCHO + 0.20 CH ₂ (OH)COCH ₂ (OH) + 0.70 CH ₂ (OH)COCH(OH)(OO•) + 0.45 H ₂ O ₂ - 0.15 O ₂	R(478)	4.0 10 ⁸			= k(2 CH ₃ COCH ₂ (OO•)) - 8
Oxidation of dihydroxyacetone					38
CH ₂ (OH)COCH ₂ (OH) + HO• → CH ₂ (OH)COC•H(OH) + H ₂ O		8.1 10 ⁶			BR: 100% - 39
CH ₂ (OH)COC•H(OH) + O ₂ → CH ₂ (OH)COCH(OH)(OO•)		2.0 10 ⁹			5
CH ₂ (OH)COCH ₂ (OH) + HO• → CH ₂ (OH)COCH(OH)(OO•) + H ₂ O - O ₂	R(479)	8.1 10 ⁸			13
CH ₂ (OH)COCH ₂ (OH) + NO ₃ [•] → CH ₂ (OH)COC•H(OH) + NO ₃ ⁻ + H ⁺		6.6·10 ⁶			BR: 100%
CH ₂ (OH)COC•H(OH) + O ₂ → CH ₂ (OH)COCH(OH)(OO•)		2.0 10 ⁹			5
CH ₂ (OH)COCH ₂ (OH) + NO ₃ [•] → CH ₂ (OH)COCH(OH)(OO•) + NO ₃ ⁻ + H ⁺ - O ₂	R(480)	6.6·10 ⁶	2117		= k(CH ₂ (OH)CH ₂ (OH) + NO ₃ [•]) - 2
CH ₂ (OH)COCH(OH)(OO•) + OH ⁻ → CH ₂ (OH)COCH(O ⁻)(OO•) + H ₂ O		4.0 10 ⁹			
CH ₂ (OH)COCH(O ⁻)(OO•) → CH ₂ (OH)COCHO + O ₂ ^{•-}					9
CH ₂ (OH)COCH(OH)(OO•) + OH ⁻ → CH ₂ (OH)COCHO + O ₂ ^{•-} + H ₂ O	R(481)	4.0 10 ⁹			= k(CH ₃ CH(OH)(OO•) + OH ⁻)
CH ₂ (OH)COCH(OH)(OO•) → CH ₂ (OH)COCHO + HO ₂ [•]	R(482)	1.9 10 ²			10
Oxidation of propionic acid					
Pathway 1: CH ₃ CH ₂ CO(OH) + HO• → CH ₃ C•HCO(OH) + H ₂ O		5.0 10 ⁷			BR: 17% - 40
CH ₃ C•HCO(OH) + O ₂ → CH ₃ CH(OO•)CO(OH)		2.0 10 ⁹			5
Pathway 2: CH ₃ CH ₂ CO(OH) + HO• → C•H ₂ CH ₂ CO(OH) + H ₂ O		2.7 10 ⁸			BR: 83% - 40
C•H ₂ CH ₂ CO(OH) + O ₂ → CH ₂ (OO•)CH ₂ CO(OH)		2.0 10 ⁹			5
CH ₃ CH ₂ CO(OH) + HO• → 0.17 CH ₃ CH(OO•)CO(OH) + 0.83 CH ₂ (OO•)CH ₂ CO(OH) + H ₂ O - O ₂	R(483)	3.2·10 ⁸	2300	Ervens et al., 2003	
Pathway 1: CH ₃ CH ₂ CO(OH) + NO ₃ [•] → CH ₃ C•HCO(OH) + NO ₃ ⁻ + H ⁺		1.3 10 ⁴			BR: 17%
CH ₃ C•HCO(OH) + O ₂ → CH ₃ CH(OO•)CO(OH)		2.0 10 ⁹			5
Pathway 2: CH ₃ CH ₂ CO(OH) + NO ₃ [•] → C•H ₂ CH ₂ CO(OH) + NO ₃ ⁻ + H ⁺		6.1 10 ⁴			BR: 83%
C•H ₂ CH ₂ CO(OH) + O ₂ → CH ₂ (OO•)CH ₂ CO(OH)		2.0 10 ⁹			5
CH ₃ CH ₂ CO(OH) + NO ₃ [•] → 0.17 CH ₃ CH(OO•)CO(OH) + 0.83 CH ₂ (OO•)CH ₂ CO(OH) + NO ₃ ⁻ + H ⁺ - O ₂	R(484)	7.4·10 ⁴		Rousse and George, 2004	2
Pathway 1: CH ₃ CH ₂ CO(O ⁻) + HO• → CH ₃ C•HCO(O ⁻) + H ₂ O		2.0 10 ⁸			BR: 28% - 41
CH ₃ C•HCO(O ⁻) + O ₂ → CH ₃ CH(OO•)CO(O ⁻)		2.0 10 ⁹			5
Pathway 2: CH ₃ CH ₂ CO(O ⁻) + HO• → C•H ₂ CH ₂ CO(O ⁻) + H ₂ O		5.3 10 ⁸			BR: 72% - 41
C•H ₂ CH ₂ CO(O ⁻) + O ₂ → CH ₂ (OO•)CH ₂ CO(O ⁻)		2.0 10 ⁹			5
CH ₃ CH ₂ CO(O ⁻) + HO• → 0.28 CH ₃ CH(OO•)CO(O ⁻) + 0.72 CH ₂ (OO•)CH ₂ CO(O ⁻) + H ₂ O - O ₂	R(485)	7.3·10 ⁸	1800	Ervens et al., 2003	
Pathway 1: CH ₃ CH ₂ CO(O ⁻) + NO ₃ [•] → CH ₃ C•HCO(O ⁻) + NO ₃ ⁻ + H ⁺		2.1 10 ⁴			BR: 28%
CH ₃ C•HCO(O ⁻) + O ₂ → CH ₃ CH(OO•)CO(O ⁻)		2.0 10 ⁹			5
Pathway 2: CH ₃ CH ₂ CO(O ⁻) + NO ₃ [•] → C•H ₂ CH ₂ CO(O ⁻) + NO ₃ ⁻ + H ⁺		5.3 10 ⁴			BR: 72%
C•H ₂ CH ₂ CO(O ⁻) + O ₂ → CH ₂ (OO•)CH ₂ CO(O ⁻)		2.0 10 ⁹			5
CH ₃ CH ₂ CO(O ⁻) + NO ₃ [•] → 0.28 CH ₃ CH(OO•)CO(O ⁻) + 0.72 CH ₂ (OO•)CH ₂ CO(O ⁻) + NO ₃ ⁻ + H ⁺ - O ₂	R(486)	7.4·10 ⁴			= k(CH ₃ CH ₂ CO(OH) + NO ₃ [•]) - 2
Pathway 1: 2 CH ₃ CH(OO•)CO(OH) → 2 CH ₃ COCO(OH) + H ₂ O ₂		2.2 10 ⁷			BR: 30%
Pathway 2: 2 CH ₃ CH(OO•)CO(OH) (+ 2 H ₂ O) → 2 CH ₃ CHO + 2 CO ₂ + H ₂ O ₂ + 2 H ₂ O		2.2 10 ⁷			BR: 30%
Pathway 3: 2 CH ₃ CH(OO•)CO(OH) → CH ₃ COCO(OH) + CH ₃ CH(OH)CO(OH) + O ₂		2.2 10 ⁷			BR: 30%
Pathway 4: 2 CH ₃ CH(OO•)CO(OH) → 2 CH ₃ CH(O•)CO(OH) + O ₂		9.0 10 ⁶			BR: 10%

Reactions		k_{298} ($M^{-n+1} s^{-1}$)	Ea/R (K)	References	Notes
$CH_3CH(O^{\bullet})CO(OH) \rightarrow CH_3C^{\bullet}(OH)CO(OH)$					4
$CH_3C^{\bullet}(OH)CO(OH) + O_2 \rightarrow CH_3C(OH)(OO^{\bullet})CO(OH)$		$2.0 \cdot 10^9$			5
$2 CH_3CH(OO^{\bullet})CO(OH) \rightarrow 0.90 CH_3COCO(OH) + 0.30 CH_3CH(OH)CO(OH) + 0.20 CH_3C(OH)(OO^{\bullet})CO(OH) + 0.60 CH_3CHO + 0.60 CO_2 + 0.60 H_2O_2 + 0.20 O_2$	R(487)	$7.5 \cdot 10^7$			= $k(2 CH_2(OO^{\bullet})CO(O^-)) - 8$
Pathway 1: $2 CH_3CH(OO^{\bullet})CO(O^-) \rightarrow 2 CH_3COCO(O^-) + H_2O_2$		$2.2 \cdot 10^7$			BR: 30%
Pathway 2: $2 CH_3CH(OO^{\bullet})CO(O^-) (+ 2 H_2O) \rightarrow 2 CH_3CHO + 2 CO_2 + H_2O_2 + 2 OH^-$		$2.2 \cdot 10^7$			BR: 30%
Pathway 3: $2 CH_3CH(OO^{\bullet})CO(O^-) \rightarrow CH_3COCO(O^-) + CH_3CH(OH)CO(O^-) + O_2$		$2.2 \cdot 10^7$			BR: 30%
Pathway 4: $2 CH_3CH(OO^{\bullet})CO(O^-) \rightarrow 2 CH_3CH(O^{\bullet})CO(O^-) + O_2$		$9.0 \cdot 10^6$			BR: 10%
$CH_3CH(O^{\bullet})CO(O^-) \rightarrow CH_3C^{\bullet}(OH)CO(O^-)$					4
$CH_3C^{\bullet}(OH)CO(O^-) + O_2 \rightarrow CH_3C(OH)(OO^{\bullet})CO(O^-)$		$2.0 \cdot 10^9$			5
$2 CH_3CH(OO^{\bullet})CO(O^-) \rightarrow 0.90 CH_3COCO(O^-) + 0.30 CH_3CH(OH)CO(O^-) + 0.20 CH_3C(OH)(OO^{\bullet})CO(O^-) + 0.60 CH_3CHO + 0.60 CO_2 + 0.60 H_2O_2 + 0.20 O_2 - 0.60 H_2O + 0.60 OH^-$	R(488)	$7.5 \cdot 10^7$			= $k(2 CH_2(OO^{\bullet})CO(O^-)) - 8$
Pathway 1: $2 CH_2(OO^{\bullet})CH_2CO(OH) \rightarrow 2 CO(OH)CH_2CHO + H_2O_2$		$3.0 \cdot 10^7$			BR: 20%
Pathway 2: $2 CH_2(OO^{\bullet})CH_2CO(OH) \rightarrow 2 CH_2(O^{\bullet})CH_2CO(OH) + O_2$		$1.3 \cdot 10^8$			BR: 80%
$CH_2(O^{\bullet})CH_2CO(OH) \rightarrow C^{\bullet}H(OH)CH_2CO(OH)$					4
$C^{\bullet}H(OH)CH_2CO(OH) + O_2 \rightarrow CH(OH)(OO^{\bullet})CH_2CO(OH)$		$2.0 \cdot 10^9$			5
$2 CH_2(OO^{\bullet})CH_2CO(OH) \rightarrow 0.40 CO(OH)CH_2CHO + 1.60 CH(OH)(OO^{\bullet})CH_2CO(OH) + 0.20 H_2O_2 - 0.80 O_2$	R(489)	$1.6 \cdot 10^8$	-1600		= $k(2 CH_3CH_2(OO^{\bullet})) - 8$
Pathway 1: $2 CH_2(OO^{\bullet})CH_2CO(O^-) \rightarrow 2 CO(O^-)CH_2CHO + H_2O_2$		$3.0 \cdot 10^7$			BR: 20%
Pathway 2: $2 CH_2(OO^{\bullet})CH_2CO(O^-) \rightarrow 2 CH_2(O^{\bullet})CH_2CO(O^-) + O_2$		$1.3 \cdot 10^8$			BR: 80%
$CH_2(O^{\bullet})CH_2CO(O^-) \rightarrow C^{\bullet}H(OH)CH_2CO(O^-)$					4
$C^{\bullet}H(OH)CH_2CO(O^-) + O_2 \rightarrow CH(OH)(OO^{\bullet})CH_2CO(O^-)$		$2.0 \cdot 10^9$			5
$2 CH_2(OO^{\bullet})CH_2CO(O^-) \rightarrow 0.40 CO(O^-)CH_2CHO + 1.60 CH(OH)(OO^{\bullet})CH_2CO(O^-) + 0.20 H_2O_2 - 0.80 O_2$	R(490)	$1.6 \cdot 10^8$	-1600		= $k(2 CH_3CH_2(OO^{\bullet})) - 8$
Oxidation of 3-oxopropionic acid					42
Pathway 1: $CO(OH)CH_2CH(OH)(OH) + HO^{\bullet} \rightarrow CO(OH)CH_2C^{\bullet}(OH)(OH) + H_2O$		$5.5 \cdot 10^8$			BR: 58% - 43
$CO(OH)CH_2C^{\bullet}(OH)(OH) + O_2 \rightarrow CO(OH)CH_2C(OH)(OH)(OO^{\bullet})$		$2.0 \cdot 10^9$			5
Pathway 2: $CO(OH)CH_2CH(OH)(OH) + HO^{\bullet} \rightarrow CO(OH)CH_2CH(OH)(O^{\bullet}) + H_2O$		$4.0 \cdot 10^8$			BR: 42% - 43
$CO(OH)CH_2CH(OH)(O^{\bullet}) \rightarrow CO(OH)CH_2C^{\bullet}(OH)(OH)$					4
$CO(OH)CH_2C^{\bullet}(OH)(OH) + O_2 \rightarrow CO(OH)CH_2C(OH)(OH)(OO^{\bullet})$		$2.0 \cdot 10^9$			5
$CO(OH)CH_2CH(OH)(OH) + HO^{\bullet} \rightarrow CO(OH)CH_2C(OH)(OH)(OO^{\bullet}) + H_2O - O_2$	R(491)	$9.5 \cdot 10^8$			13
$CO(OH)CH_2CH(OH)(OH) + NO_3^{\bullet} \rightarrow CO(OH)CH_2C^{\bullet}(OH)(OH) + NO_3^- + H^+$		$9.1 \cdot 10^5$			BR: 100%
$CO(OH)CH_2C^{\bullet}(OH)(OH) + O_2 \rightarrow CO(OH)CH_2C(OH)(OH)(OO^{\bullet})$		$2.0 \cdot 10^9$			5
$CO(OH)CH_2CH(OH)(OH) + NO_3^{\bullet} \rightarrow CO(OH)CH_2C(OH)(OH)(OO^{\bullet}) + NO_3^- + H^+ - O_2$	R(492)	$9.1 \cdot 10^5$	3971		= $k(CH_2(OH)CO(OH) + NO_3^{\bullet}) - 2$
Pathway 1: $CO(O^-)CH_2CH(OH)(OH) + HO^{\bullet} \rightarrow CO(O^-)CH_2C^{\bullet}(OH)(OH) + H_2O$		$1.0 \cdot 10^9$			BR: 59% - 44
$CO(O^-)CH_2C^{\bullet}(OH)(OH) + O_2 \rightarrow CO(O^-)CH_2C(OH)(OH)(OO^{\bullet})$		$2.0 \cdot 10^9$			5
Pathway 2: $CO(O^-)CH_2CH(OH)(OH) + HO^{\bullet} \rightarrow CO(O^-)CH_2CH(OH)(O^{\bullet}) + H_2O$		$4.3 \cdot 10^8$			BR: 25% - 44
$CO(O^-)CH_2CH(OH)(O^{\bullet}) \rightarrow CO(O^-)CH_2C^{\bullet}(OH)(OH)$					4
$CO(O^-)CH_2C^{\bullet}(OH)(OH) + O_2 \rightarrow CO(O^-)CH_2C(OH)(OH)(OO^{\bullet})$		$2.0 \cdot 10^9$			5
Pathway 3: $CO(O^-)CH_2CH(OH)(OH) + HO^{\bullet} \rightarrow CO(O^{\bullet})CH_2CH(OH)(OH) + OH^-$		$2.7 \cdot 10^8$			BR: 16% - 44
$CO(O^{\bullet})CH_2CH(OH)(OH) \rightarrow C^{\bullet}H_2CH(OH)(OH) + CO_2$					6 - 7
$C^{\bullet}H_2CH(OH)(OH) + O_2 \rightarrow CH(OH)(OH)CH_2(OO^{\bullet})$		$2.0 \cdot 10^9$			5
$CO(O^-)CH_2CH(OH)(OH) + HO^{\bullet} \rightarrow 0.84 CO(O^-)CH_2C(OH)(OH)(OO^{\bullet}) + 0.16 CH(OH)(OH)CH_2(OO^{\bullet}) + 0.16 CO_2 + 0.84 H_2O + 0.16 OH^- - O_2$	R(493)	$1.7 \cdot 10^9$			13
Pathway 1: $CO(O^-)CH_2CH(OH)(OH) + NO_3^{\bullet} \rightarrow CO(O^-)CH_2C^{\bullet}(OH)(OH) + NO_3^- + H^+$		$1.0 \cdot 10^7$			BR: 100%
$CO(O^-)CH_2C^{\bullet}(OH)(OH) + O_2 \rightarrow CO(O^-)CH_2C(OH)(OH)(OO^{\bullet})$		$2.0 \cdot 10^9$			5
$CO(O^-)CH_2CH(OH)(OH) + NO_3^{\bullet} \rightarrow CO(O^-)CH_2C(OH)(OH)(OO^{\bullet}) + NO_3^- + H^+ - O_2$	R(494)	$1.0 \cdot 10^7$	3008		= $k(CH_2(OH)CO(O^-) + NO_3^{\bullet}) - 2$
$CO(O^-)CH_2CHO + HO^{\bullet} \rightarrow CO(O^-)CH_2C^{\bullet}O + H_2O$		$3.0 \cdot 10^9$			BR: 100% - 45

Reactions		k_{298} ($M^{-n+1} s^{-1}$)	Ea/R (K)	References	Notes
$CO(O^-)CH_2C^*O + O_2 \rightarrow CO(O^-)CH_2CO(OO^*)$		$2.0 \cdot 10^9$			5
$CO(O^-)CH_2CHO + HO^* \rightarrow CO(O^-)CH_2CO(OO^*) + H_2O - O_2$	R(495)	$3.0 \cdot 10^9$			13
$CO(O^-)CH_2CHO + NO_3^* \rightarrow CO(O^-)CH_2C^*O + NO_3^- + H^+$		$1.0 \cdot 10^7$			BR: 100%
$CO(O^-)CH_2C^*O + O_2 \rightarrow CO(O^-)CH_2CO(OO^*)$		$2.0 \cdot 10^9$			5
$CO(O^-)CH_2CHO + NO_3^* \rightarrow CO(O^-)CH_2CO(OO^*) + NO_3^- + H^+ - O_2$	R(496)	$1.0 \cdot 10^7$	3008		= $k(CH_2(OH)CO(O^-) + NO_3^*) - 2$
$CO(OH)CH_2C(OH)(OH)(OO^*) + OH^- \rightarrow CO(OH)CH_2C(OH)(O^-)(OO^*) + H_2O$		$4.0 \cdot 10^9$			
$CO(OH)CH_2C(OH)(O^-)(OO^*) \rightarrow CO(OH)CH_2CO(OH) + O_2^{\bullet-} + H_2O$	R(497)	$4.0 \cdot 10^9$			9
$CO(OH)CH_2C(OH)(OH)(OO^*) + OH^- \rightarrow CO(OH)CH_2CO(OH) + HO_2^*$	R(498)	$1.0 \cdot 10^6$			= $k(CH_3CH(OH)(OO^*) + OH^-)$
$CO(O^-)CH_2C(OH)(OH)(OO^*) + OH^- \rightarrow CO(O^-)CH_2C(OH)(O^-)(OO^*) + H_2O$		$4.0 \cdot 10^9$			15
$CO(O^-)CH_2C(OH)(O^-)(OO^*) \rightarrow CO(OH)CH_2CO(O^-) + O_2^{\bullet-}$					9
$CO(O^-)CH_2C(OH)(OH)(OO^*) + OH^- \rightarrow CO(OH)CH_2CO(O^-) + O_2^{\bullet-} + H_2O$	R(499)	$4.0 \cdot 10^9$			= $k(CH_3CH(OH)(OO^*) + OH^-)$
$CO(O^-)CH_2C(OH)(OH)(OO^*) \rightarrow CO(OH)CH_2CO(O^-) + HO_2^*$	R(500)	$1.0 \cdot 10^6$			15
$2 CO(O^-)CH_2CO(OO^*) \rightarrow 2 CO(O^-)CH_2CO(O^*) + O_2$		$1.6 \cdot 10^8$			
$CO(O^-)CH_2CO(O^*) \rightarrow CO(O^-)C^*H_2 + CO_2$					6 - 7
$CO(O^-)C^*H_2 + O_2 \rightarrow CH_2(OO^*)CO(O^-)$		$2.0 \cdot 10^9$			5
$2 CO(O^-)CH_2CO(OO^*) \rightarrow 2 CH_2(OO^*)CO(O^-) + 2 CO_2 - O_2$	R(501)	$1.6 \cdot 10^8$	-1600		= $k(2 CH_3CH_2(OO^*)) - 8$
Oxidation of malonic acid					
$CO(OH)CH_2CO(OH) + HO^* \rightarrow CO(OH)C^*HCO(OH) + H_2O$		$1.9 \cdot 10^7$			BR: 100%
$CO(OH)C^*HCO(OH) + O_2 \rightarrow CO(OH)CH(OO^*)CO(OH)$		$2.0 \cdot 10^9$			5
$CO(OH)CH_2CO(OH) + HO^* \rightarrow CO(OH)CH(OO^*)CO(OH) + H_2O - O_2$	R(502)	$1.9 \cdot 10^7$		Wang et al., 2001	
$CO(OH)CH_2CO(OH) + NO_3^* \rightarrow CO(OH)C^*HCO(OH) + NO_3^- + H^+$		$5.1 \cdot 10^4$			BR: 100%
$CO(OH)C^*HCO(OH) + O_2 \rightarrow CO(OH)CH(OO^*)CO(OH)$		$2.0 \cdot 10^9$			5
$CO(OH)CH_2CO(OH) + NO_3^* \rightarrow CO(OH)CH(OO^*)CO(OH) + NO_3^- + H^+ - O_2$	R(503)	$5.1 \cdot 10^4$		De Semainville et al., 2010	2
Pathway 1: $CO(OH)CH_2CO(O^-) + HO^* \rightarrow CO(OH)C^*HCO(O^-) + H_2O$		$4.0 \cdot 10^6$			BR: 6% - 46
$CO(OH)C^*HCO(O^-) + O_2 \rightarrow CO(OH)CH(OO^*)CO(O^-)$		$2.0 \cdot 10^9$			5
Pathway 2: $CO(OH)CH_2CO(O^-) + HO^* \rightarrow CO(O^*)CH_2CO(OH) + OH^-$		$5.6 \cdot 10^7$			BR: 94% - 46
$CO(O^*)CH_2CO(OH) \rightarrow C^*H_2CO(OH) + CO_2$					6 - 7
$C^*H_2CO(OH) + O_2 \rightarrow CH_2(OO^*)CO(OH)$		$2.0 \cdot 10^9$			5
$CO(OH)CH_2CO(O^-) + HO^* \rightarrow 0.06 CO(OH)CH(OO^*)CO(O^-) + 0.94 CH_2(OO^*)CO(OH) + 0.94 CO_2 + 0.06 H_2O + 0.94 OH^- - O_2$	R(504)	$6.0 \cdot 10^7$	1300	Ervens et al., 2003	
Pathway 1: $CO(OH)CH_2CO(O^-) + NO_3^* \rightarrow CO(OH)C^*HCO(O^-) + NO_3^- + H^+$		$5.6 \cdot 10^6$	3369		
$CO(OH)C^*HCO(O^-) + O_2 \rightarrow CO(OH)CH(OO^*)CO(O^-)$		$2.0 \cdot 10^9$			5
$CO(OH)CH_2CO(O^-) + NO_3^* \rightarrow CO(OH)CH(OO^*)CO(O^-) + NO_3^- + H^+ - O_2$	R(505)	$5.6 \cdot 10^6$	3369	De Semainville et al., 2010	2
Pathway 1: $CO(O^-)CH_2CO(O^-) + HO^* \rightarrow CO(O^-)C^*HCO(O^-) + H_2O$		$1.0 \cdot 10^7$			BR: 6% - 47
$CO(O^-)C^*HCO(O^-) + O_2 \rightarrow CO(O^-)CH(OO^*)CO(O^-)$		$2.0 \cdot 10^9$			5
Pathway 2: $CO(O^-)CH_2CO(O^-) + HO^* \rightarrow CO(O^*)CH_2CO(O^-) + OH^-$		$1.0 \cdot 10^8$			BR: 94% - 47
$CO(O^*)CH_2CO(O^-) \rightarrow C^*H_2CO(O^-) + CO_2$					6 - 7
$C^*H_2CO(O^-) + O_2 \rightarrow CH_2(OO^*)CO(O^-)$		$2.0 \cdot 10^9$			5
$CO(O^-)CH_2CO(O^-) + HO^* \rightarrow 0.06 CO(O^-)CH(OO^*)CO(O^-) + 0.94 CH_2(OO^*)CO(O^-) + 0.94 CO_2 + 0.06 H_2O + 0.94 OH^- - O_2$	R(506)	$1.1 \cdot 10^8$		Wang et al., 2001	
Pathway 1: $CO(O^-)CH_2CO(O^-) + NO_3^* \rightarrow CO(O^-)C^*HCO(O^-) + NO_3^- + H^+$		$2.3 \cdot 10^7$	3008		
$CO(O^-)C^*HCO(O^-) + O_2 \rightarrow CO(O^-)CH(OO^*)CO(O^-)$		$2.0 \cdot 10^9$			5

Reactions		k ₂₉₈ (M ⁻ⁿ⁺¹ s ⁻¹)	Ea/R (K)	References	Notes
CO(O ⁻)CH ₂ CO(O ⁻) + NO ₃ [•] → CO(O ⁻)CH(OO [•])CO(O ⁻) + NO ₃ ⁻ + H ⁺ - O ₂	R(507)	2.3·10 ⁷	3008	De Semainville et al., 2010	2
Pathway 1: 2 CO(OH)CH(OO [•])CO(OH) → 2 CO(OH)COCO(OH) + H ₂ O ₂		2.2 10 ⁷			BR: 30%
Pathway 2: 2 CO(OH)CH(OO [•])CO(OH) (+ 2 H ₂ O) → 2 CHOCO(OH) + 2 CO ₂ + H ₂ O ₂ + 2 H ₂ O		2.2 10 ⁷			BR: 30%
Pathway 3: 2 CO(OH)CH(OO [•])CO(OH) → CO(OH)COCO(OH) + CO(OH)CH(OH)CO(OH) + O ₂		2.2 10 ⁷			BR: 30%
Pathway 4: 2 CO(OH)CH(OO [•])CO(OH) → 2 CO(OH)CH(O [•])CO(OH) + O ₂		9.0 10 ⁶			BR: 10%
CO(OH)CH(O [•])CO(OH) → CO(OH)C [•] (OH)CO(OH)					4
CO(OH)C [•] (OH)CO(OH) + O ₂ → CO(OH)C(OH)(OO [•])CO(OH)		2.0·10 ⁹			5
2 CO(OH)CH(OO [•])CO(OH) → 0.90 CO(OH)COCO(OH) + 0.30 CO(OH)CH(OH)CO(OH) + 0.20 CO(OH)C(OH)(OO [•])CO(OH) + 0.60 CHOCO(OH) + 0.60 CO ₂ + 0.60 H ₂ O ₂ + 0.20 O ₂	R(508)	7.5·10 ⁷			= k(2 CH ₂ (OO [•])CO(O ⁻)) - 8
Pathway 1: 2 CO(OH)CH(OO [•])CO(O ⁻) → 2 CO(OH)COCO(O ⁻) + H ₂ O ₂		2.2 10 ⁷			BR: 30%
Pathway 2: 2 CO(OH)CH(OO [•])CO(O ⁻) (+ 2 H ₂ O) → 2 CHOCO(OH) + 2 CO ₂ + H ₂ O ₂ + 2 OH ⁻		2.2 10 ⁷			BR: 30%
Pathway 3: 2 CO(OH)CH(OO [•])CO(O ⁻) → CO(OH)COCO(O ⁻) + CO(OH)CH(OH)CO(O ⁻) + O ₂		2.2 10 ⁷			BR: 30%
Pathway 4: 2 CO(OH)CH(OO [•])CO(O ⁻) → 2 CO(OH)CH(O [•])CO(O ⁻) + O ₂		9.0 10 ⁶			BR: 10%
CO(OH)CH(O [•])CO(O ⁻) → CO(OH)C [•] (OH)CO(O ⁻)					4
CO(OH)C [•] (OH)CO(O ⁻) + O ₂ → CO(OH)C(OH)(OO [•])CO(O ⁻)		2.0·10 ⁹			5
2 CO(OH)CH(OO [•])CO(O ⁻) → 0.90 CO(OH)COCO(O ⁻) + 0.30 CO(OH)CH(OH)CO(O ⁻) + 0.20 CO(OH)C(OH)(OO [•])CO(O ⁻) + 0.60 CHOCO(OH) + 0.60 CO ₂ + 0.60 H ₂ O ₂ + 0.60 OH ⁻ + 0.20 O ₂ - 0.60 H ₂ O	R(509)	7.5·10 ⁷			= k(2 CH ₂ (OO [•])CO(O ⁻)) - 8
Pathway 1: 2 CO(O ⁻)CH(OO [•])CO(O ⁻) → 2 CO(O ⁻)COCO(O ⁻) + H ₂ O ₂		2.2 10 ⁷			BR: 30%
Pathway 2: 2 CO(O ⁻)CH(OO [•])CO(O ⁻) (+ 2 H ₂ O) → 2 CHOCO(O ⁻) + 2 CO ₂ + H ₂ O ₂ + 2 OH ⁻		2.2 10 ⁷			BR: 30%
Pathway 3: 2 CO(O ⁻)CH(OO [•])CO(O ⁻) → CO(O ⁻)COCO(O ⁻) + CO(O ⁻)CH(OH)CO(O ⁻) + O ₂		2.2 10 ⁷			BR: 30%
Pathway 4: 2 CO(O ⁻)CH(OO [•])CO(O ⁻) → 2 CO(O ⁻)CH(O [•])CO(O ⁻) + O ₂		9.0 10 ⁶			BR: 10%
CO(O ⁻)CH(O [•])CO(O ⁻) → CO(O ⁻)C [•] (OH)CO(O ⁻)					4
CO(O ⁻)C [•] (OH)CO(O ⁻) + O ₂ → CO(O ⁻)C(OH)(OO [•])CO(O ⁻)		2.0·10 ⁹			5
2 CO(O ⁻)CH(OO [•])CO(O ⁻) → 0.90 CO(O ⁻)COCO(O ⁻) + 0.30 CO(O ⁻)CH(OH)CO(O ⁻) + 0.20 CO(O ⁻)C(OH)(OO [•])CO(O ⁻) + 0.60 CHOCO(O ⁻) + 0.60 CO ₂ + 0.60 H ₂ O ₂ + 0.60 OH ⁻ + 0.20 O ₂ - 0.60 H ₂ O	R(510)	7.5·10 ⁷			= k(2 CH ₂ (OO [•])CO(O ⁻)) - 8
Oxidation of tartronic acid					
Pathway 1: CO(OH)CH(OH)CO(OH) + HO [•] → CO(OH)C [•] (OH)CO(OH) + H ₂ O		2.0·10 ⁷			BR: 11% - 48
CO(OH)C [•] (OH)CO(OH) + O ₂ → CO(OH)C(OH)(OO [•])CO(OH)		2.0·10 ⁹			5
Pathway 2: CO(OH)CH(OH)CO(OH) + HO [•] → CO(OH)CH(O [•])CO(OH) + H ₂ O		1.5 10 ⁸			BR: 89% - 48
CO(OH)CH(O [•])CO(OH) → C [•] O(OH) + CHOCO(OH)					6 - 7
C [•] O(OH) + O ₂ → CO(OH)(OO [•])		2.0·10 ⁹			5
CO(OH)CH(OH)CO(OH) + HO [•] → 0.11 CO(OH)C(OH)(OO [•])CO(OH) + 0.89 CHOCO(OH) + 0.89 CO(OH)(OO [•]) + H ₂ O - O ₂	R(511)	1.7·10 ⁸		Schuchmann et al., 1995	
CO(OH)CH(OH)CO(OH) + NO ₃ [•] → CO(OH)C [•] (OH)CO(OH) + NO ₃ ⁻ + H ⁺		5.1·10 ⁴			BR: 100%
CO(OH)C [•] (OH)CO(OH) + O ₂ → CO(OH)C(OH)(OO [•])CO(OH)		2.0·10 ⁹			5
CO(OH)CH(OH)CO(OH) + NO ₃ [•] → CO(OH)C(OH)(OO [•])CO(OH) + NO ₃ ⁻ + H ⁺ - O ₂	R(512)	5.1·10 ⁴			= k(CO(OH)CH ₂ CO(OH) + NO ₃ [•]) - 2
Pathway 1: CO(OH)CH(OH)CO(O ⁻) + HO [•] → CO(OH)C [•] (OH)CO(O ⁻) + H ₂ O		4.0·10 ⁷			BR: 11% - 49
CO(OH)C [•] (OH)CO(O ⁻) + O ₂ → CO(OH)C(OH)(OO [•])CO(O ⁻)		2.0·10 ⁹			5
Pathway 2: CO(OH)CH(OH)CO(O ⁻) + HO [•] → CO(OH)CH(O [•])CO(O ⁻) + H ₂ O		1.5 10 ⁸			BR: 43% - 49
CO(OH)CH(O [•])CO(O ⁻) → C [•] O(OH) + CHOCO(O ⁻)					6 - 7
C [•] O(OH) + O ₂ → CO(OH)(OO [•])		2.0·10 ⁹			5
Pathway 3: CO(OH)CH(OH)CO(O ⁻) + HO [•] → CO(OH)CH(OH)CO(O [•]) + OH ⁻		1.7 10 ⁸			BR: 46% - 49
CO(OH)CH(OH)CO(O [•]) → CO(OH)C [•] H(OH) + CO ₂					6 - 7
CO(OH)C [•] H(OH) + O ₂ → CH(OH)(OO [•])CO(OH)		2.0·10 ⁹			5

Reactions		k_{298} ($M^{-n+1} s^{-1}$)	Ea/R (K)	References	Notes
$CO(OH)CH(OH)CO(O^-) + HO^\bullet \rightarrow 0.11 CO(OH)C(OH)(OO^\bullet)CO(O^-) + 0.46 CH(OH)(OO^\bullet)CO(OH) + 0.43 CO(OH)(OO^\bullet) + 0.46 CO_2 + 0.43 CHOCO(O^-) + 0.46 OH^- + 0.54 H_2O - O_2$ Pathway 1: $CO(OH)CH(OH)CO(O^-) + NO_3^\bullet \rightarrow CO(OH)C^\bullet(OH)CO(O^-) + NO_3^- + H^+$ $CO(OH)C^\bullet(OH)CO(O^-) + O_2 \rightarrow CO(OH)C(OH)(OO^\bullet)CO(O^-)$ $CO(OH)CH(OH)CO(O^-) + NO_3^\bullet \rightarrow CO(OH)C(OH)(OO^\bullet)CO(O^-) + NO_3^- + H^+ - O_2$	R(513)	$3.6 \cdot 10^8$		Schuchmann et al., 1995	
		$5.6 \cdot 10^6$	3369		BR: 100% 5
	R(514)	$5.6 \cdot 10^6$	3369		= $k(CO(OH)CH_2CO(O^-) + NO_3^\bullet) - 2$ BR: 13% - 50 5
Pathway 1: $CO(O^-)CH(OH)CO(O^-) + HO^\bullet \rightarrow CO(O^-)C^\bullet(OH)CO(O^-) + H_2O$ $CO(O^-)C^\bullet(OH)CO(O^-) + O_2 \rightarrow CO(O^-)C(OH)(OO^\bullet)CO(O^-)$ Pathway 2: $CO(O^-)CH(OH)CO(O^-) + HO^\bullet \rightarrow CO(O^-)CH(O^\bullet)CO(O^-) + H_2O$ $CO(O^-)CH(O^\bullet)CO(O^-) \rightarrow C^\bullet O(O^-) + CHOCO(O^-)$ $C^\bullet O(O^-) + O_2 \rightarrow CO(O^-)(OO^\bullet)$ $CO(O^-)(OO^\bullet) \rightarrow CO_2 + O_2^\bullet$ Pathway 3: $CO(O^-)CH(OH)CO(O^-) + HO^\bullet \rightarrow CO(O^-)CH(OH)CO(O^\bullet) + OH^-$ $CO(O^-)CH(OH)CO(O^\bullet) \rightarrow CO(O^-)C^\bullet H(OH) + CO_2$ $CO(O^-)C^\bullet H(OH) + O_2 \rightarrow CH(OH)(OO^\bullet)CO(O^-)$		$6.0 \cdot 10^7$			BR: 27% - 50 6 - 7
		$2.0 \cdot 10^9$			5
		$1.2 \cdot 10^8$			9
		$2.0 \cdot 10^9$			BR: 60% - 50 6 - 7 5
		$2.6 \cdot 10^8$			
$CO(O^-)CH(OH)CO(O^-) + HO^\bullet \rightarrow 0.13 CO(O^-)C(OH)(OO^\bullet)CO(O^-) + 0.27 CHOCO(O^-) + 0.60 CH(OH)(OO^\bullet)CO(O^-) + 0.87 CO_2 + 0.27 O_2^\bullet + 0.60 OH^- + 0.40 H_2O - O_2$ $CO(O^-)CH(OH)CO(O^-) + NO_3^\bullet \rightarrow CO(O^-)C^\bullet(OH)CO(O^-) + NO_3^- + H^+$ $CO(O^-)C^\bullet(OH)CO(O^-) + O_2 \rightarrow CO(O^-)C(OH)(OO^\bullet)CO(O^-)$ $CO(O^-)CH(OH)CO(O^-) + NO_3^\bullet \rightarrow CO(O^-)C(OH)(OO^\bullet)CO(O^-) + NO_3^- + H^+ - O_2$	R(515)	$4.4 \cdot 10^8$		Schuchmann et al., 1995	
		$2.3 \cdot 10^7$	3008		BR: 100% 5
		$2.0 \cdot 10^9$			
	R(516)	$2.3 \cdot 10^7$	3008		= $k(CO(O^-)CH_2CO(O^-) + NO_3^\bullet) - 2$ 9
		$4.0 \cdot 10^9$			
$CO(OH)C(OH)(OO^\bullet)CO(OH) + OH^- \rightarrow CO(OH)C(O^-)(OO^\bullet)CO(OH) + H_2O$ $CO(OH)C(O^-)(OO^\bullet)CO(OH) \rightarrow CO(OH)COCO(OH) + O_2^\bullet$ $CO(OH)C(OH)(OO^\bullet)CO(OH) + OH^- \rightarrow CO(OH)COCO(OH) + O_2^\bullet + H_2O$ $CO(OH)C(OH)(OO^\bullet)CO(OH) \rightarrow CO(OH)COCO(OH) + HO_2^\bullet$ $CO(OH)C(OH)(OO^\bullet)CO(O^-) + OH^- \rightarrow CO(OH)C(O^-)(OO^\bullet)CO(O^-) + H_2O$ $CO(OH)C(O^-)(OO^\bullet)CO(O^-) \rightarrow CO(OH)COCO(O^-) + O_2^\bullet$	R(517)	$4.0 \cdot 10^9$			= $k(CH_3CH(OH)(OO^\bullet) + OH^-)$
	R(518)	$1.1 \cdot 10^4$			= $k(CO(O^-)C(OH)(OO^\bullet)CO(O^-))$ 9
		$4.0 \cdot 10^9$			
	R(519)	$4.0 \cdot 10^9$			= $k(CH_3CH(OH)(OO^\bullet) + OH^-)$
	R(520)	$1.1 \cdot 10^4$			= $k(CO(O^-)C(OH)(OO^\bullet)CO(O^-))$ 9
		$4.0 \cdot 10^9$			
$CO(O^-)C(OH)(OO^\bullet)CO(O^-) + OH^- \rightarrow CO(O^-)COCO(O^-) + O_2^\bullet + H_2O$ $CO(O^-)C(OH)(OO^\bullet)CO(O^-) \rightarrow CO(O^-)COCO(O^-) + HO_2^\bullet$ $CO(O^-)C(OH)(OO^\bullet)CO(O^-) + OH^- \rightarrow CO(O^-)C(O^-)(OO^\bullet)CO(O^-) + H_2O$ $CO(O^-)C(O^-)(OO^\bullet)CO(O^-) \rightarrow CO(O^-)COCO(O^-) + O_2^\bullet$	R(521)	$4.0 \cdot 10^9$			= $k(CH_3CH(OH)(OO^\bullet) + OH^-)$
	R(522)	$1.1 \cdot 10^4$		Schuchmann et al., 1995	
Oxidation of mesoxalic acid (or ketomalonic acid)					51
$CO(OH)C(OH)(OH)CO(OH) + HO^\bullet \rightarrow CO(OH)C(O^\bullet)(OH)CO(OH) + H_2O$ $CO(OH)C(O^\bullet)(OH)CO(OH) \rightarrow CO(OH)CO(OH) + C^\bullet O(OH)$ $C^\bullet O(OH) + O_2 \rightarrow CO(OH)(OO^\bullet)$		$1.4 \cdot 10^8$			BR: 100% 6 - 7 5
		$2.0 \cdot 10^9$			
$CO(OH)C(OH)(OH)CO(OH) + HO^\bullet \rightarrow CO(OH)CO(OH) + CO(OH)(OO^\bullet) + H_2O - O_2$ $CO(OH)COCO(O^-) + HO^\bullet \rightarrow CO(OH)COCO(O^\bullet) + OH^-$ $CO(OH)COCO(O^\bullet) \rightarrow CO(OH)C^\bullet O + CO_2$ $CO(OH)C^\bullet O + O_2 \rightarrow CO(OH)CO(OO^\bullet)$	R(523)	$1.4 \cdot 10^8$		Schaefer, 2012	
		$3.2 \cdot 10^8$			BR: 100% 6 - 7 5
		$2.0 \cdot 10^9$			
$CO(OH)COCO(O^-) + HO^\bullet \rightarrow CO(OH)CO(OO^\bullet) + CO_2 + OH^- - O_2$ $CO(OH)C(OH)(OH)CO(O^-) + HO^\bullet \rightarrow CO(OH)C(O^\bullet)(OH)CO(O^-) + H_2O$ $CO(OH)C(O^\bullet)(OH)CO(O^-) \rightarrow CO(OH)CO(O^-) + C^\bullet O(OH)$ $C^\bullet O(OH) + O_2 \rightarrow CO(OH)(OO^\bullet)$	R(524)	$3.2 \cdot 10^8$			13 BR: 100% 6 - 7 5
		$1.4 \cdot 10^8$			
		$2.0 \cdot 10^9$			

Reactions		k ₂₉₈ (M ⁻ⁿ⁺¹ s ⁻¹)	Ea/R (K)	References	Notes
CO(OH)C(OH)(OH)CO(O ⁻) + HO [•] → CO(OH)CO(O ⁻) + CO(OH)(OO [•]) + H ₂ O - O ₂	R(525)	1.4 10 ⁸		Schaefer, 2012	
CO(O ⁻)COCO(O ⁻) + HO [•] → CO(O ⁻)COCO(O [•]) + OH ⁻		1.6 10 ⁸			BR: 100%
CO(O ⁻)COCO(O [•]) → CO(O ⁻)C [•] O + CO ₂					6 - 7
CO(O ⁻)C [•] O + O ₂ → CO(O ⁻)CO(OO [•])		2.0·10 ⁹			5
CO(O ⁻)COCO(O ⁻) + HO [•] → CO(O ⁻)CO(OO [•]) + CO ₂ + OH ⁻ - O ₂	R(526)	1.6 10 ⁸		Schaefer, 2012	
Oxidation of lactic acid					
Pathway 1: CH ₃ CH(OH)CO(OH) + HO [•] → CH ₃ C [•] (OH)CO(OH) + H ₂ O		1.8 10 ⁸			BR: 34% - 52
CH ₃ C [•] (OH)CO(OH) + O ₂ → CH ₃ C(OH)(OO [•])CO(OH)		2.0 10 ⁹			5
Pathway 2: CH ₃ CH(OH)CO(OH) + HO [•] → C [•] H ₂ CH(OH)CO(OH) + H ₂ O		2.3 10 ⁸			BR: 44% - 52
C [•] H ₂ CH(OH)CO(OH) + O ₂ → CH ₂ (OO [•])CH(OH)CO(OH)		2.0 10 ⁹			5
Pathway 3: CH ₃ CH(OH)CO(OH) + HO [•] → CH ₃ CH(O [•])CO(OH) + H ₂ O		1.1 10 ⁸			BR: 22% - 52
CH ₃ CH(O [•])CO(OH) → CH ₃ CHO + C [•] O(OH)					6 - 7
C [•] O(OH) + O ₂ → CO(OH)(OO [•])		2.0 10 ⁹			5
CH ₃ CH(OH)CO(OH) + HO [•] → 0.34 CH ₃ C(OH)(OO [•])CO(OH) + 0.44 CH ₂ (OO [•])CH(OH)CO(OH) + 0.22 CH ₃ CHO + 0.22 CO(OH)(OO [•]) + H ₂ O - O ₂	R(527)	5.3·10 ⁸	1120	Martin et al., 2008	
Pathway 1: CH ₃ CH(OH)CO(OH) + NO ₃ [•] → CH ₃ C [•] (OH)CO(OH) + NO ₃ ⁻ + H ⁺		9.0 10 ⁵			BR: 43%
CH ₃ C [•] (OH)CO(OH) + O ₂ → CH ₃ C(OH)(OO [•])CO(OH)		2.0 10 ⁹			5
Pathway 2: CH ₃ CH(OH)CO(OH) + NO ₃ [•] → C [•] H ₂ CH(OH)CO(OH) + NO ₃ ⁻ + H ⁺		1.2 10 ⁶			BR: 57%
C [•] H ₂ CH(OH)CO(OH) + O ₂ → CH ₂ (OO [•])CH(OH)CO(OH)		2.0 10 ⁹			5
CH ₃ CH(OH)CO(OH) + NO ₃ [•] → 0.43 CH ₃ C(OH)(OO [•])CO(OH) + 0.57 CH ₂ (OO [•])CH(OH)CO(OH) + NO ₃ ⁻ + H ⁺ - O ₂	R(528)	2.1·10 ⁶	3248	De Semainville et al., 2007	2
Pathway 1: CH ₃ CH(OH)CO(O ⁻) + HO [•] → CH ₃ C [•] (OH)CO(O ⁻) + H ₂ O		3.3 10 ⁸			BR: 42% - 53
CH ₃ C [•] (OH)CO(O ⁻) + O ₂ → CH ₃ C(OH)(OO [•])CO(O ⁻)		2.0 10 ⁹			5
Pathway 2: CH ₃ CH(OH)CO(O ⁻) + HO [•] → C [•] H ₂ CH(OH)CO(O ⁻) + H ₂ O		2.2 10 ⁸			BR: 28% - 53
C [•] H ₂ CH(OH)CO(O ⁻) + O ₂ → CH ₂ (OO [•])CH(OH)CO(O ⁻)		2.0 10 ⁹			5
Pathway 3: CH ₃ CH(OH)CO(O ⁻) + HO [•] → CH ₃ CH(O [•])CO(O ⁻) + H ₂ O		1.1 10 ⁸			BR: 14% - 53
CH ₃ CH(O [•])CO(O ⁻) → CH ₃ CHO + C [•] O(O ⁻)					6 - 7
C [•] O(O ⁻) + O ₂ → CO(O ⁻)(OO [•])		2.0 10 ⁹			5
CO(O ⁻)(OO [•]) → CO ₂ + O ₂ ^{•-}					9
Pathway 4: CH ₃ CH(OH)CO(O ⁻) + HO [•] → CH ₃ CH(OH)CO(O [•]) + OH ⁻		1.2 10 ⁸			BR: 16% - 53
CH ₃ CH(OH)CO(O [•]) → CH ₃ C [•] H(OH) + CO ₂					6 - 7
CH ₃ C [•] H(OH) + O ₂ → CH ₃ CH(OH)(OO [•])		2.0·10 ⁹			5
CH ₃ CH(OH)CO(O ⁻) + HO [•] → 0.42 CH ₃ C(OH)(OO [•])CO(O ⁻) + 0.28 CH ₂ (OO [•])CH(OH)CO(O ⁻) + 0.14 CH ₃ CHO + 0.16 CH ₃ CH(OH)(OO [•]) + 0.30 CO ₂ + 0.14 O ₂ ^{•-} + 0.16 OH ⁻ + 0.84 H ₂ O - O ₂	R(529)	7.9·10 ⁸	1294	Martin et al., 2008	
Pathway 1: CH ₃ CH(OH)CO(O ⁻) + NO ₃ [•] → CH ₃ C [•] (OH)CO(O ⁻) + NO ₃ ⁻ + H ⁺		6.0 10 ⁶			BR: 60%
CH ₃ C [•] (OH)CO(O ⁻) + O ₂ → CH ₃ C(OH)(OO [•])CO(O ⁻)		2.0 10 ⁹			5
Pathway 2: CH ₃ CH(OH)CO(O ⁻) + NO ₃ [•] → C [•] H ₂ CH(OH)CO(O ⁻) + NO ₃ ⁻ + H ⁺		4.0 10 ⁶			BR: 40%
C [•] H ₂ CH(OH)CO(O ⁻) + O ₂ → CH ₂ (OO [•])CH(OH)CO(O ⁻)		2.0 10 ⁹			5
CH ₃ CH(OH)CO(O ⁻) + NO ₃ [•] → 0.60 CH ₃ C(OH)(OO [•])CO(O ⁻) + 0.40 CH ₂ (OO [•])CH(OH)CO(O ⁻) + NO ₃ ⁻ + H ⁺ - O ₂	R(530)	1.0·10 ⁷	2646	De Semainville et al., 2007	2
CH ₃ C(OH)(OO [•])CO(OH) + OH ⁻ → CH ₃ C(O ⁻)(OO [•])CO(OH) + H ₂ O		4.0 10 ⁹			
CH ₃ C(O ⁻)(OO [•])CO(OH) → CH ₃ COCO(OH) + O ₂ ^{•-}					9
CH ₃ C(OH)(OO [•])CO(OH) + OH ⁻ → CH ₃ COCO(OH) + O ₂ ^{•-} + H ₂ O	R(531)	4.0 10 ⁹			= k(CH ₃ CH(OH)(OO [•]) + OH ⁻)
CH ₃ C(OH)(OO [•])CO(OH) → CH ₃ COCO(OH) + HO ₂ [•]	R(532)	665			10
CH ₃ C(OH)(OO [•])CO(O ⁻) + OH ⁻ → CH ₃ C(O ⁻)(OO [•])CO(O ⁻) + H ₂ O		4.0 10 ⁹			
CH ₃ C(O ⁻)(OO [•])CO(O ⁻) → CH ₃ COCO(O ⁻) + O ₂ ^{•-}					9
CH ₃ C(OH)(OO [•])CO(O ⁻) + OH ⁻ → CH ₃ COCO(O ⁻) + O ₂ ^{•-} + H ₂ O	R(533)	4.0 10 ⁹			= k(CH ₃ CH(OH)(OO [•]) + OH ⁻)

Reactions		k ₂₉₈ (M ⁻ⁿ⁺¹ s ⁻¹)	Ea/R (K)	References	Notes
CH ₃ C(OH)(OO•)CO(O ⁻) → CH ₃ COCO(O ⁻) + HO ₂ •	R(534)	665			10
Pathway 1: 2 CH ₂ (OO•)CH(OH)CO(OH) → 2 CHOCH(OH)CO(OH) + H ₂ O ₂		5.0 10 ⁷			BR: 50%
Pathway 2: 2 CH ₂ (OO•)CH(OH)CO(OH) → CHOCH(OH)CO(OH) + CH ₂ (OH)CH(OH)CO(OH) + O ₂		3.3 10 ⁷			BR: 33%
Pathway 3: 2 CH ₂ (OO•)CH(OH)CO(OH) → 2 CH ₂ (O•)CH(OH)CO(OH) + O ₂		1.7 10 ⁷			BR: 17%
CH ₂ (O•)CH(OH)CO(OH) → CH ₂ O + C•H(OH)CO(OH)					6 - 7
C•H(OH)CO(OH) + O ₂ → CH(OH)(OO•)CO(OH)		2.0·10 ⁹			5
2 CH ₂ (OO•)CH(OH)CO(OH) → 1.33 CHOCH(OH)CO(OH) + 0.33 CH ₂ (OH)CH(OH)CO(OH) + 0.34 CH(OH)(OO•)CO(OH) + 0.34 CH ₂ O + 0.50 H ₂ O ₂ + 0.16 O ₂	R(535)	1.0 10 ⁸			= k(2 CH ₂ (OH)CH ₂ (OO•)) - 8
Pathway 1: 2 CH ₂ (OO•)CH(OH)CO(O ⁻) → 2 CHOCH(OH)CO(O ⁻) + H ₂ O ₂		5.0 10 ⁷			BR: 50%
Pathway 2: 2 CH ₂ (OO•)CH(OH)CO(O ⁻) → CHOCH(OH)CO(O ⁻) + CH ₂ (OH)CH(OH)CO(O ⁻) + O ₂		3.3 10 ⁷			BR: 33%
Pathway 3: 2 CH ₂ (OO•)CH(OH)CO(O ⁻) → 2 CH ₂ (O•)CH(OH)CO(O ⁻) + O ₂		1.7 10 ⁷			BR: 17%
CH ₂ (O•)CH(OH)CO(O ⁻) → CH ₂ O + C•H(OH)CO(O ⁻)					6 - 7
C•H(OH)CO(O ⁻) + O ₂ → CH(OH)(OO•)CO(O ⁻)		2.0·10 ⁹			5
2 CH ₂ (OO•)CH(OH)CO(O ⁻) → 1.33 CHOCH(OH)CO(O ⁻) + 0.33 CH ₂ (OH)CH(OH)CO(O ⁻) + 0.34 CH(OH)(OO•)CO(O ⁻) + 0.34 CH ₂ O + 0.50 H ₂ O ₂ + 0.16 O ₂	R(536)	1.0 10 ⁸			= k(2 CH ₂ (OH)CH ₂ (OO•)) - 8
Oxidation of 2,3-dihydroxypropanoic acid					
Pathway 1: CH ₂ (OH)CH(OH)CO(OH) + HO• → C•H(OH)CH(OH)CO(OH) + H ₂ O		5.5·10 ⁸			BR: 85% - 54
C•H(OH)CH(OH)CO(OH) + O ₂ → CH(OH)(OO•)CH(OH)CO(OH)		2.0·10 ⁹			5
Pathway 2: CH ₂ (OH)CH(OH)CO(OH) + HO• → CH ₂ (O•)CH(OH)CO(OH) + H ₂ O		1.0·10 ⁸			BR: 15% - 54
CH ₂ (O•)CH(OH)CO(OH) → CH ₂ O + C•H(OH)CO(OH)					6 - 7
C•H(OH)CO(OH) + O ₂ → CH(OH)(OO•)CO(OH)		2.0·10 ⁹			5
CH ₂ (OH)CH(OH)CO(OH) + HO• → 0.85 CH(OH)(OO•)CH(OH)CO(OH) + 0.15 CH ₂ O + 0.15 CH(OH)(OO•)CO(OH) + H ₂ O - O ₂	R(537)	6.5 10 ⁸			13
CH ₂ (OH)CH(OH)CO(OH) + NO ₃ • → C•H(OH)CH(OH)CO(OH) + NO ₃ ⁻ + H ⁺		9.1 10 ⁵			BR: 100%
C•H(OH)CH(OH)CO(OH) + O ₂ → CH(OH)(OO•)CH(OH)CO(OH)		2.0·10 ⁹			5
CH ₂ (OH)CH(OH)CO(OH) + NO ₃ • → CH(OH)(OO•)CH(OH)CO(OH) + NO ₃ ⁻ + H ⁺ - O ₂	R(538)	9.1 10 ⁵	3971		= k(CH ₂ (OH)CO(OH) + NO ₃ •) - 2
Pathway 1: CH ₂ (OH)CH(OH)CO(O ⁻) + HO• → C•H(OH)CH(OH)CO(O ⁻) + H ₂ O		1.1·10 ⁹			BR: 87% - 55
C•H(OH)CH(OH)CO(O ⁻) + O ₂ → CH(OH)(OO•)CH(OH)CO(O ⁻)		2.0·10 ⁹			5
Pathway 2: CH ₂ (OH)CH(OH)CO(O ⁻) + HO• → CH ₂ (OH)C•(OH)CO(O ⁻) + H ₂ O		2.0·10 ⁸			BR: 13% - 55
CH ₂ (OH)C•(OH)CO(O ⁻) + O ₂ → CH ₂ (OH)C(OH)(OO•)CO(O ⁻)		2.0·10 ⁹			5
CH ₂ (OH)CH(OH)CO(O ⁻) + HO• → 0.87 CH(OH)(OO•)CH(OH)CO(O ⁻) + 0.13 CH ₂ (OH)C(OH)(OO•)CO(O ⁻) + H ₂ O - O ₂	R(539)	1.3 10 ⁹			13
Pathway 1: CH ₂ (OH)CH(OH)CO(O ⁻) + NO ₃ • → C•H(OH)CH(OH)CO(O ⁻) + NO ₃ ⁻ + H ⁺		8.7·10 ⁶			BR: 87%
C•H(OH)CH(OH)CO(O ⁻) + O ₂ → CH(OH)(OO•)CH(OH)CO(O ⁻)		2.0·10 ⁹			5
Pathway 2: CH ₂ (OH)CH(OH)CO(O ⁻) + NO ₃ • → CH ₂ (OH)C•(OH)CO(O ⁻) + NO ₃ ⁻ + H ⁺		1.3·10 ⁶			BR: 13%
CH ₂ (OH)C•(OH)CO(O ⁻) + O ₂ → CH ₂ (OH)C(OH)(OO•)CO(O ⁻)		2.0·10 ⁹			5
CH ₂ (OH)CH(OH)CO(O ⁻) + NO ₃ • → 0.87 CH(OH)(OO•)CH(OH)CO(O ⁻) + 0.13 CH ₂ (OH)C(OH)(OO•)CO(O ⁻) + NO ₃ ⁻ + H ⁺ - O ₂	R(540)	1.0 10 ⁷	3008		= k(CH ₂ (OH)CO(O ⁻) + NO ₃ •) - 2
CH(OH)(OO•)CH(OH)CO(OH) + OH ⁻ → CH(O ⁻)(OO•)CH(OH)CO(OH) + H ₂ O		4.0 10 ⁹			
CH(O ⁻)(OO•)CH(OH)CO(OH) → CHOCH(OH)CO(OH) + O ₂ • ⁻					9
CH(OH)(OO•)CH(OH)CO(OH) + OH ⁻ → CHOCH(OH)CO(OH) + O ₂ • ⁻ + H ₂ O	R(541)	4.0 10 ⁹			= k(CH ₃ CH(OH)(OO•) + OH ⁻)
CH(OH)(OO•)CH(OH)CO(OH) → CHOCH(OH)CO(OH) + HO ₂ •	R(542)	1.9 10 ²			10
CH(OH)(OO•)CH(OH)CO(O ⁻) + OH ⁻ → CH(O ⁻)(OO•)CH(OH)CO(O ⁻) + H ₂ O		4.0 10 ⁹			
CH(O ⁻)(OO•)CH(OH)CO(O ⁻) → CHOCH(OH)CO(O ⁻) + O ₂ • ⁻					9
CH(OH)(OO•)CH(OH)CO(O ⁻) + OH ⁻ → CHOCH(OH)CO(O ⁻) + O ₂ • ⁻ + H ₂ O	R(543)	4.0 10 ⁹			= k(CH ₃ CH(OH)(OO•) + OH ⁻)
CH(OH)(OO•)CH(OH)CO(O ⁻) → CHOCH(OH)CO(O ⁻) + HO ₂ •	R(544)	1.9 10 ²			10

Reactions		k_{298} ($M^{-n+1} s^{-1}$)	Ea/R (K)	References	Notes
$CH_2(OH)C(OH)(OO^{\bullet})CO(OH) + OH^{-} \rightarrow CH_2(OH)C(O^{-})(OO^{\bullet})CO(OH) + H_2O$		$4.0 \cdot 10^9$			
$CH_2(OH)C(O^{-})(OO^{\bullet})CO(OH) \rightarrow CH_2(OH)COCO(OH) + O_2^{\bullet-}$					9
$CH_2(OH)C(OH)(OO^{\bullet})CO(OH) + OH^{-} \rightarrow CH_2(OH)COCO(OH) + O_2^{\bullet-} + H_2O$	R(545)	$4.0 \cdot 10^9$			= $k(CH_3CH(OH)(OO^{\bullet}) + OH^{-})$
$CH_2(OH)C(OH)(OO^{\bullet})CO(OH) \rightarrow CH_2(OH)COCO(OH) + HO_2^{\bullet}$	R(546)	665			10
$CH_2(OH)C(OH)(OO^{\bullet})CO(O^{-}) + OH^{-} \rightarrow CH_2(OH)C(O^{-})(OO^{\bullet})CO(O^{-}) + H_2O$		$4.0 \cdot 10^9$			
$CH_2(OH)C(O^{-})(OO^{\bullet})CO(O^{-}) \rightarrow CH_2(OH)COCO(O^{-}) + O_2^{\bullet-}$					9
$CH_2(OH)C(OH)(OO^{\bullet})CO(O^{-}) + OH^{-} \rightarrow CH_2(OH)COCO(O^{-}) + O_2^{\bullet-} + H_2O$	R(547)	$4.0 \cdot 10^9$			= $k(CH_3CH(OH)(OO^{\bullet}) + OH^{-})$
$CH_2(OH)C(OH)(OO^{\bullet})CO(O^{-}) \rightarrow CH_2(OH)COCO(O^{-}) + HO_2^{\bullet}$	R(548)	665			10
Oxidation of 2-hydroxy, 3-oxopropanoic acid					56
Pathway 1: $CH(OH)(OH)CH(OH)CO(OH) + HO^{\bullet} \rightarrow CH(OH)(O^{\bullet})CH(OH)CO(OH) + H_2O$		$4.2 \cdot 10^8$			BR: 63% - 57
$CH(OH)(O^{\bullet})CH(OH)CO(OH) \rightarrow CHO(OH) + C^{\bullet}H(OH)CO(OH)$					6 - 7
$C^{\bullet}H(OH)CO(OH) + O_2 \rightarrow CH(OH)(OO^{\bullet})CO(OH)$		$2.0 \cdot 10^9$			5
Pathway 2: $CH(OH)(OH)CH(OH)CO(OH) + HO^{\bullet} \rightarrow C^{\bullet}(OH)(OH)CH(OH)CO(OH) + H_2O$		$2.5 \cdot 10^8$			BR: 37% - 57
$C^{\bullet}(OH)(OH)CH(OH)CO(OH) + O_2 \rightarrow CO(OH)CH(OH)C(OH)(OH)(OO^{\bullet})$		$2.0 \cdot 10^9$			5
$CH(OH)(OH)CH(OH)CO(OH) + HO^{\bullet} \rightarrow 0.37 CO(OH)CH(OH)C(OH)(OH)(OO^{\bullet}) + 0.63 CHO(OH) + 0.63$	R(549)	$6.7 \cdot 10^8$			13
$CH(OH)(OO^{\bullet})CO(OH) + H_2O - O_2$					
$CH(OH)(OH)CH(OH)CO(OH) + NO_3^{\bullet} \rightarrow C^{\bullet}(OH)(OH)CH(OH)CO(OH) + NO_3^{-} + H^{\bullet}$		$1.1 \cdot 10^6$			BR: 100%
$C^{\bullet}(OH)(OH)CH(OH)CO(OH) + O_2 \rightarrow CO(OH)CH(OH)C(OH)(OH)(OO^{\bullet})$		$2.0 \cdot 10^9$			5
$CH(OH)(OH)CH(OH)CO(OH) + NO_3^{\bullet} \rightarrow CO(OH)CH(OH)C(OH)(OH)(OO^{\bullet}) + NO_3^{-} + H^{\bullet} - O_2$	R(550)	$1.1 \cdot 10^6$			= $k(CH_2(OH)CH(OH)(OH) + NO_3^{\bullet}) - 2$
Pathway 1: $CH(OH)(OH)CH(OH)CO(O^{-}) + HO^{\bullet} \rightarrow CH(OH)(O^{\bullet})CH(OH)CO(O^{-}) + H_2O$		$4.9 \cdot 10^8$			BR: 49% - 58
$CH(OH)(O^{\bullet})CH(OH)CO(O^{-}) \rightarrow CHO(OH) + C^{\bullet}H(OH)CO(O^{-})$					6 - 7
$C^{\bullet}H(OH)CO(O^{-}) + O_2 \rightarrow CH(OH)(OO^{\bullet})CO(O^{-})$		$2.0 \cdot 10^9$			5
Pathway 2: $CH(OH)(OH)CH(OH)CO(O^{-}) + HO^{\bullet} \rightarrow C^{\bullet}(OH)(OH)CH(OH)CO(O^{-}) + H_2O$		$5.1 \cdot 10^8$			BR: 51% - 58
$C^{\bullet}(OH)(OH)CH(OH)CO(O^{-}) + O_2 \rightarrow CO(O^{-})CH(OH)C(OH)(OH)(OO^{\bullet})$		$2.0 \cdot 10^9$			5
$CH(OH)(OH)CH(OH)CO(O^{-}) + HO^{\bullet} \rightarrow 0.51 CO(O^{-})CH(OH)C(OH)(OH)(OO^{\bullet}) + 0.49 CH(OH)(OO^{\bullet})CO(O^{-}) + 0.49$	R(551)	$1.0 \cdot 10^9$			13
$CHO(OH) + H_2O - O_2$					
$CH(OH)(OH)CH(OH)CO(O^{-}) + NO_3^{\bullet} \rightarrow C^{\bullet}(OH)(OH)CH(OH)CO(O^{-}) + NO_3^{-} + H^{\bullet}$		$1.1 \cdot 10^6$			BR: 100%
$C^{\bullet}(OH)(OH)CH(OH)CO(O^{-}) + O_2 \rightarrow CO(O^{-})CH(OH)C(OH)(OH)(OO^{\bullet})$		$2.0 \cdot 10^9$			5
$CH(OH)(OH)CH(OH)CO(O^{-}) + NO_3^{\bullet} \rightarrow CO(O^{-})CH(OH)C(OH)(OH)(OO^{\bullet}) + NO_3^{-} + H^{\bullet} - O_2$	R(552)	$1.1 \cdot 10^6$			= $k(CH_2(OH)CH(OH)(OH) + NO_3^{\bullet}) - 2$
$CO(OH)CH(OH)C(OH)(OH)(OO^{\bullet}) + OH^{-} \rightarrow CO(OH)CH(OH)C(OH)(O^{-})(OO^{\bullet}) + H_2O$		$4.0 \cdot 10^9$			
$CO(OH)CH(OH)C(OH)(O^{-})(OO^{\bullet}) \rightarrow CO(OH)CH(OH)CO(OH) + O_2^{\bullet-}$					9
$CO(OH)CH(OH)C(OH)(OH)(OO^{\bullet}) + OH^{-} \rightarrow CO(OH)CH(OH)CO(OH) + O_2^{\bullet-} + H_2O$	R(553)	$4.0 \cdot 10^9$			= $k(CH_3CH(OH)(OO^{\bullet}) + OH^{-})$
$CO(OH)CH(OH)C(OH)(OH)(OO^{\bullet}) \rightarrow CO(OH)CH(OH)CO(OH) + HO_2^{\bullet}$	R(554)	$1.0 \cdot 10^6$			15
$CO(O^{-})CH(OH)C(OH)(OH)(OO^{\bullet}) + OH^{-} \rightarrow CO(O^{-})CH(OH)C(OH)(O^{-})(OO^{\bullet}) + H_2O$		$4.0 \cdot 10^9$			
$CO(O^{-})CH(OH)C(OH)(O^{-})(OO^{\bullet}) \rightarrow CO(OH)CH(OH)CO(O^{-}) + O_2^{\bullet-}$					9
$CO(O^{-})CH(OH)C(OH)(OH)(OO^{\bullet}) + OH^{-} \rightarrow CO(OH)CH(OH)CO(O^{-}) + O_2^{\bullet-} + H_2O$	R(555)	$4.0 \cdot 10^9$			= $k(CH_3CH(OH)(OO^{\bullet}) + OH^{-})$
$CO(O^{-})CH(OH)C(OH)(OH)(OO^{\bullet}) \rightarrow CO(OH)CH(OH)CO(O^{-}) + HO_2^{\bullet}$	R(556)	$1.0 \cdot 10^6$			15
$CH(OH)(OH)C(OH)(OO^{\bullet})CO(OH) + OH^{-} \rightarrow CH(OH)(OH)C(O^{-})(OO^{\bullet})CO(OH) + H_2O$		$4.0 \cdot 10^9$			
$CH(OH)(OH)C(O^{-})(OO^{\bullet})CO(OH) \rightarrow CH(OH)(OH)COCO(OH) + O_2^{\bullet-}$					9
$CH(OH)(OH)C(OH)(OO^{\bullet})CO(OH) + OH^{-} \rightarrow CH(OH)(OH)COCO(OH) + O_2^{\bullet-} + H_2O$	R(557)	$4.0 \cdot 10^9$			= $k(CH_3CH(OH)(OO^{\bullet}) + OH^{-})$
$CH(OH)(OH)C(OH)(OO^{\bullet})CO(OH) \rightarrow CH(OH)(OH)COCO(OH) + HO_2^{\bullet}$	R(558)	$1.9 \cdot 10^2$			15
$CH(OH)(OH)C(OH)(OO^{\bullet})CO(O^{-}) + OH^{-} \rightarrow CH(OH)(OH)C(O^{-})(OO^{\bullet})CO(O^{-}) + H_2O$		$4.0 \cdot 10^9$			
$CH(OH)(OH)C(O^{-})(OO^{\bullet})CO(O^{-}) \rightarrow CH(OH)(OH)COCO(O^{-}) + O_2^{\bullet-}$					9
$CH(OH)(OH)C(OH)(OO^{\bullet})CO(O^{-}) + OH^{-} \rightarrow CH(OH)(OH)COCO(O^{-}) + O_2^{\bullet-} + H_2O$	R(559)	$4.0 \cdot 10^9$			= $k(CH_3CH(OH)(OO^{\bullet}) + OH^{-})$

Reactions		k_{298} ($M^{-n+1} s^{-1}$)	Ea/R (K)	References	Notes
$CH(OH)(OH)C(OH)(OO^{\bullet})CO(O^-) \rightarrow CH(OH)(OH)COCO(O^-) + HO_2^{\bullet}$	R(560)	$1.9 \cdot 10^2$			15
Oxidation of 3-oxopyruvic acid					59
Pathway 1: $CO(OH)C(OH)(OH)CH(OH)(OH) + HO^{\bullet} \rightarrow CO(OH)C(OH)(OH)CH(OH)(O^{\bullet}) + H_2O$ $CO(OH)C(OH)(OH)CH(OH)(O^{\bullet}) \rightarrow CO(OH)C^{\bullet}(OH)(OH) + CHO(OH)$ $CO(OH)C^{\bullet}(OH)(OH) + O_2 \rightarrow CO(OH)C(OH)(OH)(OO^{\bullet})$		$3.6 \cdot 10^8$			BR: 52% - 60 6 - 7
Pathway 2: $CO(OH)C(OH)(OH)CH(OH)(OH) + HO^{\bullet} \rightarrow CO(OH)C(OH)(O^{\bullet})CH(OH)(OH) + H_2O$ $CO(OH)C(OH)(O^{\bullet})CH(OH)(OH) \rightarrow CH(OH)(OH)CO(OH) + C^{\bullet}O(OH)$ $C^{\bullet}O(OH) + O_2 \rightarrow CO(OH)(OO^{\bullet})$		$2.0 \cdot 10^9$ $2.4 \cdot 10^8$			5 BR: 35% - 60 6 - 7
Pathway 3: $CO(OH)C(OH)(OH)CH(OH)(OH) + HO^{\bullet} \rightarrow CO(OH)C(OH)(OH)C^{\bullet}(OH)(OH) + H_2O$ $CO(OH)C(OH)(OH)C^{\bullet}(OH)(OH) + O_2 \rightarrow CO(OH)C(OH)(OH)C(OH)(OH)(OO^{\bullet})$ $CO(OH)C(OH)(OH)CH(OH)(OH) + HO^{\bullet} \rightarrow 0.52 CHO(OH) + 0.52 CO(OH)C(OH)(OH)(OO^{\bullet}) + 0.13$ $CO(OH)C(OH)(OH)C(OH)(OH)(OO^{\bullet}) + 0.35 CH(OH)(OH)CO(OH) + 0.35 CO(OH)(OO^{\bullet}) + H_2O - O_2$ $CO(OH)C(OH)(OH)CH(OH)(OH) + NO_3^{\bullet} \rightarrow CO(OH)C(OH)(OH)C^{\bullet}(OH)(OH) + NO_3^- + H^+$ $CO(OH)C(OH)(OH)C^{\bullet}(OH)(OH) + O_2 \rightarrow CO(OH)C(OH)(OH)C(OH)(OH)(OO^{\bullet})$	R(561)	$2.0 \cdot 10^9$ $9.0 \cdot 10^7$ $2.0 \cdot 10^9$ $6.9 \cdot 10^8$			5 BR: 13% - 60 5 13
$CO(OH)C(OH)(OH)CH(OH)(OH) + NO_3^{\bullet} \rightarrow CO(OH)C(OH)(OH)C(OH)(OH)(OO^{\bullet}) + NO_3^- + H^+ - O_2$	R(562)	$1.1 \cdot 10^6$ $2.0 \cdot 10^9$ $1.0 \cdot 10^6$			BR:100% 5 = k(CH(OH)(OH)CO(OH) + NO ₃ [•]) - 2
Pathway 1: $CO(O^-)COCH(OH)(OH) + HO^{\bullet} \rightarrow CO(O^-)COCH(OH)(O^{\bullet}) + H_2O$ $CO(O^-)COCH(OH)(O^{\bullet}) \rightarrow CO(O^-)C^{\bullet}O + CHO(OH)$ $CO(O^-)C^{\bullet}O + O_2 \rightarrow CO(O^-)CO(OO^{\bullet})$		$3.6 \cdot 10^8$			BR: 65% - 61 6 - 7
Pathway 2: $CO(O^-)COCH(OH)(OH) + HO^{\bullet} \rightarrow CO(O^-)COC^{\bullet}(OH)(OH) + H_2O$ $CO(O^-)COC^{\bullet}(OH)(OH) + O_2 \rightarrow CO(O^-)COC(OH)(OH)(OO^{\bullet})$ $CO(O^-)COCH(OH)(OH) + HO^{\bullet} \rightarrow 0.65 CHO(OH) + 0.65 CO(O^-)CO(OO^{\bullet}) + 0.35 CO(O^-)COC(OH)(OH)(OO^{\bullet}) + H_2O - O_2$	R(563)	$2.0 \cdot 10^9$ $1.9 \cdot 10^8$ $2.0 \cdot 10^9$ $5.5 \cdot 10^8$			5 BR: 35% - 61 5 13
Pathway 1: $CO(O^-)COCH(OH)(OH) + NO_3^{\bullet} \rightarrow CO(O^-)COC^{\bullet}(OH)(OH) + NO_3^- + H^+$ $CO(O^-)COC^{\bullet}(OH)(OH) + O_2 \rightarrow CO(O^-)COC(OH)(OH)(OO^{\bullet})$ $CO(O^-)COCH(OH)(OH) + NO_3^{\bullet} \rightarrow CO(O^-)COC(OH)(OH)(OO^{\bullet}) + NO_3^- + H^+ - O_2$	R(564)	$1.8 \cdot 10^5$ $2.0 \cdot 10^9$ $1.8 \cdot 10^5$			BR: 100% 5 = k(CH(OH)(OH)CO(O^-) + NO ₃ [•]) - 2
Pathway 1: $CO(O^-)C(OH)(OH)CH(OH)(OH) + HO^{\bullet} \rightarrow CO(O^-)C(OH)(OH)CH(OH)(O^{\bullet}) + H_2O$ $CO(O^-)C(OH)(OH)CH(OH)(O^{\bullet}) \rightarrow CO(O^-)C^{\bullet}(OH)(OH) + CHO(OH)$ $CO(O^-)C^{\bullet}(OH)(OH) + O_2 \rightarrow CO(O^-)C(OH)(OH)(OO^{\bullet})$		$3.8 \cdot 10^8$			BR: 39% - 62 6 - 7
Pathway 2: $CO(O^-)C(OH)(OH)CH(OH)(OH) + HO^{\bullet} \rightarrow CO(O^-)C(OH)(O^{\bullet})CH(OH)(OH) + H_2O$ $CO(O^-)C(OH)(O^{\bullet})CH(OH)(OH) \rightarrow CH(OH)(OH)CO(OH) + CO_2^{\bullet-}$ $CO_2^{\bullet-} + O_2 \rightarrow CO_2 + O_2^{\bullet-}$		$2.0 \cdot 10^9$ $4.4 \cdot 10^8$			5 BR: 45% - 62 6 - 7
Pathway 3: $CO(O^-)C(OH)(OH)CH(OH)(OH) + HO^{\bullet} \rightarrow CO(O^-)C(OH)(OH)C^{\bullet}(OH)(OH) + H_2O$ $CO(O^-)C(OH)(OH)C^{\bullet}(OH)(OH) + O_2 \rightarrow CO(O^-)C(OH)(OH)C(OH)(OH)(OO^{\bullet})$ $CO(O^-)C(OH)(OH)CH(OH)(OH) + HO^{\bullet} \rightarrow 0.39 CHO(OH) + 0.39 CO(O^-)C(OH)(OH)(OO^{\bullet}) + 0.16 CO(O^-)$ $)C(OH)(OH)C(OH)(OH)(OO^{\bullet}) + 0.45 CH(OH)(OH)CO(OH) + 0.45 CO_2 + 0.45 O_2^{\bullet-} + H_2O - O_2$ $CO(OH)C(OH)(OH)CH(OH)(OH) + NO_3^{\bullet} \rightarrow CO(OH)C(OH)(OH)C^{\bullet}(OH)(OH) + NO_3^- + H^+$ $CO(OH)C(OH)(OH)C^{\bullet}(OH)(OH) + O_2 \rightarrow CO(OH)C(OH)(OH)C(OH)(OH)(OO^{\bullet})$	R(565)	$2.4 \cdot 10^9$ $1.6 \cdot 10^8$ $2.0 \cdot 10^9$ $9.8 \cdot 10^8$		Hislop and Bolton, 1999	BR: 16% - 62 5 13
$CO(O^-)C(OH)(OH)CH(OH)(OH) + NO_3^{\bullet} \rightarrow CO(O^-)C(OH)(OH)C(OH)(OH)(OO^{\bullet}) + NO_3^- + H^+ - O_2$	R(566)	$1.8 \cdot 10^5$ $2.0 \cdot 10^9$ $1.8 \cdot 10^5$			BR: 100% 5 = k(CH(OH)(OH)CO(O^-) + NO ₃ [•]) - 2
$CO(OH)C(OH)(OH)C(OH)(OH)(OO^{\bullet}) + OH^- \rightarrow CO(OH)C(OH)(OH)C(OH)(O^-)(OO^{\bullet}) + H_2O$ $CO(OH)C(OH)(OH)C(OH)(O^-)(OO^{\bullet}) \rightarrow CO(OH)C(OH)(OH)CO(OH) + O_2^{\bullet-}$ $CO(OH)C(OH)(OH)C(OH)(OH)(OO^{\bullet}) + OH^- \rightarrow CO(OH)C(OH)(OH)CO(OH) + O_2^{\bullet-} + H_2O$ $CO(OH)C(OH)(OH)C(OH)(OH)(OO^{\bullet}) \rightarrow CO(OH)C(OH)(OH)CO(OH) + HO_2^{\bullet}$ $CO(O^-)COC(OH)(OH)(OO^{\bullet}) + OH^- \rightarrow CO(O^-)COC(OH)(O^-)(OO^{\bullet}) + H_2O$ $CO(O^-)COC(OH)(O^-)(OO^{\bullet}) \rightarrow CO(OH)COCO(O^-) + O_2^{\bullet-}$	R(567) R(568)	$4.0 \cdot 10^9$ $4.0 \cdot 10^9$ $1.0 \cdot 10^6$ $4.0 \cdot 10^9$			9 = k(CH ₃ CH(OH)(OO [•]) + OH ⁻) 15 9

Reactions		k_{298} ($M^{-n+1} s^{-1}$)	Ea/R (K)	References	Notes
$CO(O^-)COC(OH)(OH)(OO^\bullet) + OH^- \rightarrow CO(OH)COCO(O^-) + O_2^{\bullet-} + H_2O$	R(569)	$4.0 \cdot 10^9$			= k(CH ₃ CH(OH)(OO [•]) + OH ⁻)
$CO(O^-)COC(OH)(OH)(OO^\bullet) \rightarrow CO(OH)COCO(O^-) + HO_2^\bullet$	R(570)	$1.0 \cdot 10^6$			15
$CO(O^-)C(OH)(OH)C(OH)(OH)(OO^\bullet) + OH^- \rightarrow CO(O^-)C(OH)(OH)C(OH)(O^-)(OO^\bullet) + H_2O$		$4.0 \cdot 10^9$			9
$CO(O^-)C(OH)(OH)C(OH)(O^-)(OO^\bullet) \rightarrow CO(O^-)C(OH)(OH)CO(OH) + O_2^{\bullet-}$					
$CO(O^-)C(OH)(OH)C(OH)(OH)(OO^\bullet) + OH^- \rightarrow CO(O^-)C(OH)(OH)CO(OH) + O_2^{\bullet-} + H_2O$	R(571)	$4.0 \cdot 10^9$			= k(CH ₃ CH(OH)(OO [•]) + OH ⁻)
$CO(O^-)C(OH)(OH)C(OH)(OH)(OO^\bullet) \rightarrow CO(O^-)C(OH)(OH)CO(OH) + HO_2^\bullet$	R(572)	$1.0 \cdot 10^6$			15
Oxidation of 3-hydroxypyruvic acid					63
Pathway 1: $CH_2(OH)C(OH)(OH)CO(OH) + HO^\bullet \rightarrow CH_2(OH)C(OH)(O^\bullet)CO(OH) + H_2O$		$2.5 \cdot 10^8$			BR: 48% - 64
$CH_2(OH)C(OH)(O^\bullet)CO(OH) \rightarrow CH_2(OH)CO(OH) + C^\bullet O(OH)$					6 - 7
$C^\bullet O(OH) + O_2 \rightarrow CO(OH)(OO^\bullet)$		$2.0 \cdot 10^9$			5
Pathway 2: $CH_2(OH)C(OH)(OH)CO(OH) + HO^\bullet \rightarrow C^\bullet H(OH)C(OH)(OH)CO(OH) + H_2O$		$1.9 \cdot 10^8$			BR: 36% - 64
$C^\bullet H(OH)C(OH)(OH)CO(OH) + O_2 \rightarrow CH(OH)(OO^\bullet)C(OH)(OH)CO(OH)$		$2.0 \cdot 10^9$			5
Pathway 3: $CH_2(OH)C(OH)(OH)CO(OH) + HO^\bullet \rightarrow CH_2(O^\bullet)C(OH)(OH)CO(OH) + H_2O$		$8.0 \cdot 10^7$			BR: 16% - 64
$CH_2(O^\bullet)C(OH)(OH)CO(OH) \rightarrow CH_2O + C^\bullet(OH)(OH)CO(OH)$					6 - 7
$C^\bullet(OH)(OH)CO(OH) + O_2 \rightarrow CO(OH)C(OH)(OH)(OO^\bullet)$		$2.0 \cdot 10^9$			5
$CH_2(OH)C(OH)(OH)CO(OH) + HO^\bullet \rightarrow 0.36 CH(OH)(OO^\bullet)C(OH)(OH)CO(OH) + 0.48 CH_2(OH)CO(OH) + 0.48$	R(573)	$5.2 \cdot 10^8$			13
$CO(OH)(OO^\bullet) + 0.16 CH_2O + 0.16 CO(OH)C(OH)(OH)(OO^\bullet) + H_2O - O_2$					
$CH_2(OH)C(OH)(OH)CO(OH) + NO_3^\bullet \rightarrow C^\bullet H(OH)C(OH)(OH)CO(OH) + NO_3^- + H^\bullet$		$1.1 \cdot 10^6$			BR: 100%
$C^\bullet H(OH)C(OH)(OH)CO(OH) + O_2 \rightarrow CH(OH)(OO^\bullet)C(OH)(OH)CO(OH)$		$2.0 \cdot 10^9$			5
$CH_2(OH)C(OH)(OH)CO(OH) + NO_3^\bullet \rightarrow CH(OH)(OO^\bullet)C(OH)(OH)CO(OH) + NO_3^- + H^\bullet - O_2$	R(574)	$1.0 \cdot 10^6$			= k(CH(OH)(OH)CO(OH) + NO ₃ [•]) - 2
$CH_2(OH)COCO(O^-) + HO^\bullet \rightarrow C^\bullet H(OH)COCO(O^-) + H_2O$		$5.0 \cdot 10^8$			BR: 100% - 65
$C^\bullet H(OH)COCO(O^-) + O_2 \rightarrow CH(OH)(OO^\bullet)COCO(O^-)$		$2.0 \cdot 10^9$			5
$CH_2(OH)COCO(O^-) + HO^\bullet \rightarrow CH(OH)(OO^\bullet)COCO(O^-) + H_2O - O_2$	R(575)	$5.0 \cdot 10^8$			13
$CH_2(OH)COCO(O^-) + NO_3^\bullet \rightarrow C^\bullet H(OH)COCO(O^-) + NO_3^- + H^\bullet$		$3.1 \cdot 10^6$			BR: 100%
$C^\bullet H(OH)COCO(O^-) + O_2 \rightarrow CH(OH)(OO^\bullet)COCO(O^-)$		$2.0 \cdot 10^9$			5
$CH_2(OH)COCO(O^-) + NO_3^\bullet \rightarrow CH(OH)(OO^\bullet)COCO(O^-) + NO_3^- + H^\bullet - O_2$	R(576)	$3.1 \cdot 10^6$			= k(CH ₂ (OH)CHO + NO ₃ [•])- 2
$CH(OH)(OO^\bullet)C(OH)(OH)CO(OH) + OH^- \rightarrow CH(O^-)(OO^\bullet)C(OH)(OH)CO(OH) + H_2O$		$4.0 \cdot 10^9$			9
$CH(O^-)(OO^\bullet)C(OH)(OH)CO(OH) \rightarrow CO(OH)C(OH)(OH)CHO + O_2^{\bullet-}$					
$CH(OH)(OO^\bullet)C(OH)(OH)CO(OH) + OH^- \rightarrow CO(OH)C(OH)(OH)CHO + O_2^{\bullet-} + H_2O$	R(577)	$4.0 \cdot 10^9$			= k(CH ₃ CH(OH)(OO [•]) + OH ⁻)
$CH(OH)(OO^\bullet)C(OH)(OH)CO(OH) \rightarrow CO(OH)C(OH)(OH)CHO + HO_2^\bullet$	R(578)	$1.9 \cdot 10^2$			10
$CH(OH)(OO^\bullet)COCO(O^-) + OH^- \rightarrow CH(O^-)(OO^\bullet)COCO(O^-) + H_2O$		$4.0 \cdot 10^9$			
$CH(O^-)(OO^\bullet)COCO(O^-) \rightarrow CO(O^-)COCHO + O_2^{\bullet-}$					9
$CH(OH)(OO^\bullet)COCO(O^-) + OH^- \rightarrow CO(O^-)COCHO + O_2^{\bullet-} + H_2O$	R(579)	$4.0 \cdot 10^9$			= k(CH ₃ CH(OH)(OO [•]) + OH ⁻)
$CH(OH)(OO^\bullet)COCO(O^-) \rightarrow CO(O^-)COCHO + HO_2^\bullet$	R(580)	$1.9 \cdot 10^2$			10
Oxidation of pyruvic acid					66
$CH_3COCO(OH) + h\nu \rightarrow CH_3COCO(OH)^*$	R(581)	Calculated		Reed Harris et al., 2014 ; Griffith et al., 2013	
$CH_3COCO(OH)^* + CH_3C(OH)(OH)CO(OH) \rightarrow CH_3C^\bullet(OH)CO(OH) + CH_3C^\bullet(OH)(OH) + CO_2$					67
$CH_3C^\bullet(OH)CO(OH) + O_2 \rightarrow CH_3C(OH)(OO^\bullet)CO(OH)$		$2.0 \cdot 10^9$			5
$CH_3C^\bullet(OH)(OH) + O_2 \rightarrow CH_3C(OH)(OH)(OO^\bullet)$		$2.0 \cdot 10^9$			5
$CH_3COCO(OH)^* + CH_3C(OH)(OH)CO(OH) \rightarrow CH_3C(OH)(OO^\bullet)CO(OH) + CH_3C(OH)(OH)(OO^\bullet) + CO_2 - 2 O_2$	R(582)	$1.0 \cdot 10^{10}$		Reed Harris et al., 2014 ; Griffith et al., 2013	
$CH_3COCO(O^-)^* + CH_3C(OH)(OH)CO(OH) \rightarrow CH_3C^\bullet(OH)CO(O^-) + CH_3C^\bullet(OH)(OH) + CO_2$					67

Reactions		k_{298} ($M^{-n+1} s^{-1}$)	Ea/R (K)	References	Notes
$CH_3C^*(OH)CO(O^-) + O_2 \rightarrow CH_3C(OH)(OO^*)CO(O^-)$		$2.0 \cdot 10^9$			5
$CH_3C^*(OH)(OH) + O_2 \rightarrow CH_3C(OH)(OH)(OO^*)$		$2.0 \cdot 10^9$			5
$CH_3COCO(OH)^* + CH_3C(OH)(OH)CO(O^-) \rightarrow CH_3C(OH)(OO^*)CO(O^-) + CH_3C(OH)(OH)(OO^*) + CO_2 - 2 O_2$	R(583)	$1.0 \cdot 10^{10}$		Reed Harris et al., 2014 ; Griffith et al., 2013	
$CH_3COCO(OH) + HO^* \rightarrow CO(OH)COCH_2(OO^*) + H_2O - O_2$	R(584)	$3.2 \cdot 10^8$	1804	Schaefer et al., 2012	
$CH_3COCO(OH) + NO_3^* \rightarrow CO(OH)COCH_2(OO^*) + NO_3^- + H^+ - O_2$	R(585)	$2.4 \cdot 10^6$	1804	De Semainville et al., 2007	
$CH_3COCO(OH) + H_2O_2 \rightarrow CH_3CO(OH) + CO_2 + H_2O$	R(586)	$1.2 \cdot 10^{-1}$		Schöne and Herrmann, 2014	
$CH_3COCO(O^-) + h\nu \rightarrow CH_3COCO(O^-)^*$	R(587)	Calculated		Reed Harris et al., 2014 ; Griffith et al., 2013	
$CH_3COCO(O^-) + HO^* \rightarrow CO(O^-)COCH_2(OO^*) + H_2O - O_2$	R(588)	$7.1 \cdot 10^8$	3007	Schaefer et al., 2012	
$CH_3COCO(O^-) + NO_3^* \rightarrow CO(O^-)COCH_2(OO^*) + NO_3^- + H^+ - O_2$	R(589)	$1.9 \cdot 10^7$	2887	De Semainville et al., 2007	
$CH_3COCO(O^-) + H_2O_2 \rightarrow CH_3CO(O^-) + CO_2 + H_2O$	R(590)	$7.5 \cdot 10^{-1}$		Schöne and Herrmann, 2014	
Pathway 1: $2 CO(OH)COCH_2(OO^*) \rightarrow 2 CO(OH)COCHO + H_2O_2$		$1.8 \cdot 10^8$			BR: 45%
Pathway 2: $2 CO(OH)COCH_2(OO^*) \rightarrow CO(OH)COCHO + CH_2(OH)COCO(OH) + O_2$		$8.0 \cdot 10^7$			BR: 20%
Pathway 3: $2 CO(OH)COCH_2(OO^*) \rightarrow 2 CO(OH)COCH_2(O^*) + O_2$		$1.4 \cdot 10^8$			BR: 35%
$CO(OH)COCH_2(O^*) \rightarrow CO(OH)COC^*H(OH)$					4
$CO(OH)COC^*H(OH) + O_2 \rightarrow CH(OH)(OO^*)COCO(OH)$		$2.0 \cdot 10^9$			5
$2 CO(OH)COCH_2(OO^*) \rightarrow 1.10 CO(OH)COCHO + 0.20 CH_2(OH)COCO(OH) + 0.70 CH(OH)(OO^*)COCO(OH) + 0.45 H_2O_2 - 0.15 O_2$	R(591)	$4.0 \cdot 10^8$			= $k(2 CH_3COCH_2(OO^*)) - 8$
Pathway 1: $2 CO(O^-)COCH_2(OO^*) \rightarrow 2 CO(O^-)COCHO + H_2O_2$		$1.8 \cdot 10^8$			BR: 45%
Pathway 2: $2 CO(O^-)COCH_2(OO^*) \rightarrow CO(O^-)COCHO + CH_2(OH)COCO(O^-) + O_2$		$8.0 \cdot 10^7$			BR: 20%
Pathway 3: $2 CO(O^-)COCH_2(OO^*) \rightarrow 2 CO(O^-)COCH_2(O^*) + O_2$		$1.4 \cdot 10^8$			BR: 35%
$CO(O^-)COCH_2(O^*) \rightarrow CO(O^-)COC^*H(OH)$					4
$CO(O^-)COC^*H(OH) + O_2 \rightarrow CH(OH)(OO^*)COCO(O^-)$		$2.0 \cdot 10^9$			5
$2 CO(O^-)COCH_2(OO^*) \rightarrow 1.10 CO(O^-)COCHO + 0.20 CH_2(OH)COCO(O^-) + 0.70 CH(OH)(OO^*)COCO(O^-) + 0.45 H_2O_2 - 0.15 O_2$	R(592)	$4.0 \cdot 10^8$			= $k(2 CH_3COCH_2(OO^*)) - 8$
Pathway 1: $CH_3C(OH)(OH)CO(OH) + HO^* \rightarrow CH_3C(OH)(O^*)CO(OH) + H_2O$		$2.7 \cdot 10^8$			BR: 85% - 68
$CH_3C(OH)(O^*)CO(OH) \rightarrow CH_3CO(OH) + C^*O(OH)$					6 - 7
$C^*O(OH) + O_2 \rightarrow CO(OH)(OO^*)$		$2.0 \cdot 10^9$			5
Pathway 2: $CH_3C(OH)(OH)CO(OH) + HO^* \rightarrow C^*H_2C(OH)(OH)CO(OH) + H_2O$		$5.0 \cdot 10^7$			BR: 15% - 68
$C^*H_2C(OH)(OH)CO(OH) + O_2 \rightarrow CO(OH)C(OH)(OH)CH_2(OO^*)$		$2.0 \cdot 10^9$			5
$CH_3C(OH)(OH)CO(OH) + HO^* \rightarrow 0.15 CO(OH)C(OH)(OH)CH_2(OO^*) + 0.85 CH_3CO(OH) + 0.85 CO(OH)(OO^*) + H_2O - O_2$	R(593)	$3.2 \cdot 10^8$			13
$CH_3C(OH)(OH)CO(OH) + NO_3^* \rightarrow C^*H_2C(OH)(OH)CO(OH) + NO_3^- + H^+$		$1.1 \cdot 10^6$			BR: 100%
$C^*H_2C(OH)(OH)CO(OH) + O_2 \rightarrow CO(OH)C(OH)(OH)CH_2(OO^*)$		$2.0 \cdot 10^9$			5
$CH_3C(OH)(OH)CO(OH) + NO_3^* \rightarrow CO(OH)C(OH)(OH)CH_2(OO^*) + NO_3^- + H^+ - O_2$	R(594)	$1.0 \cdot 10^6$			= $k(CH(OH)(OH)CO(OH) + NO_3^*) - 2$
Pathway 1: $2 CO(OH)C(OH)(OH)CH_2(OO^*) \rightarrow 2 CO(OH)C(OH)(OH)CHO + H_2O_2$		$5.0 \cdot 10^7$			BR: 50%
Pathway 2: $2 CO(OH)C(OH)(OH)CH_2(OO^*) \rightarrow CO(OH)C(OH)(OH)CHO + CH_2(OH)C(OH)(OH)CO(OH) + O_2$		$3.3 \cdot 10^7$			BR: 33%
Pathway 3: $2 CO(OH)C(OH)(OH)CH_2(OO^*) \rightarrow 2 CH_2(O^*)C(OH)(OH)CO(OH) + O_2$		$1.7 \cdot 10^7$			BR: 17%
$CH_2(O^*)C(OH)(OH)CO(OH) \rightarrow CH_2O + C^*(OH)(OH)CO(OH)$					6 - 7

Reactions		k_{298} ($M^{-n+1} s^{-1}$)	Ea/R (K)	References	Notes
$C^*(OH)(OH)CO(OH) + O_2 \rightarrow CO(OH)C(OH)(OH)(OO^*)$ $2 CO(OH)C(OH)(OH)CH_2(OO^*) \rightarrow 1.33 CO(OH)C(OH)(OH)CHO + 0.33 CH_2(OH)C(OH)(OH)CO(OH) + 0.34 CH_2O + 0.34 CO(OH)C(OH)(OH)(OO^*) + 0.50 H_2O_2 + 0.16 O_2$	R(595)	$2.0 \cdot 10^9$ $1.0 \cdot 10^8$			5 = k(2 CH ₂ (OH)CH ₂ (OO*)) - 8
Oxidation of acrolein (from C4)					69
CH ₂ =CHCHO + H ₂ O → CH ₂ (OH)CH ₂ CHO	R(596)	9.4 10 ⁻⁶		Pressman and Lucas, 1942	
CH ₂ (OH)CH ₂ CHO → CH ₂ =CHCHO + H ₂ O	R(597)	7.8 10 ⁻⁷		Pressman and Lucas, 1942	
CH ₂ =CHCHO + HO• → CH ₂ (OH)C•HCHO		7.0 10 ⁹			BR: 100% - 70
CH ₂ (OH)C•HCHO + O ₂ → CH ₂ (OH)CH(OO•)CHO		2.0 10 ⁹			5
CH ₂ =CHCHO + HO• → CH ₂ (OH)CH(OO•)CHO - O ₂	R(598)	7.0 10 ⁹		Lilie and Henglein, 1970	
Pathway 1: 2 CH ₂ (OH)CH(OO•)CHO → 2 CH ₂ (OH)COCHO + H ₂ O ₂		5.0 10 ⁷			BR: 50%
Pathway 2: 2 CH ₂ (OH)CH(OO•)CHO → CH ₂ (OH)COCHO + CH ₂ (OH)CH(OH)CHO + O ₂		3.3 10 ⁷			BR: 33%
Pathway 3: 2 CH ₂ (OH)CH(OO•)CHO → 2 CH ₂ (OH)CH(O•)CHO + O ₂		1.7 10 ⁷			BR: 17%
CH ₂ (OH)CH(O•)CHO → 0.50 C•H ₂ (OH) + 0.50 CHOCHO + 0.50 C•HO + 0.50 CH ₂ (OH)CHO					6 - 7
C•H ₂ (OH) + O ₂ → CH ₂ (OH)(OO•)		2.0·10 ⁹			5
C•HO + O ₂ → CHO(OO•)		2.0 10 ⁹			5
2 CH ₂ (OH)CH(OO•)CHO → 1.33 CH ₂ (OH)COCHO + 0.33 CH ₂ (OH)CH(OH)CHO + 0.17 CHOCHO + 0.17 CH ₂ (OH)CHO + 0.17 CH ₂ (OH)(OO•) + 0.17 CHO(OO•) + 0.5 H ₂ O ₂ + 0.16 O ₂	R(599)	1.0 10 ⁸			= k(2 CH ₂ (OH)CH ₂ (OO*)) - 8
Pathway 1: 2 CH ₂ (OH)CH(OO•)CH(OH)(OH) → 2 CH ₂ (OH)COCH(OH)(OH) + H ₂ O ₂		5.0 10 ⁷			BR: 50%
Pathway 2: 2 CH ₂ (OH)CH(OO•)CH(OH)(OH) → CH ₂ (OH)COCH(OH)(OH) + CH ₂ (OH)CH(OH)CH(OH)(OH) + O ₂		3.3 10 ⁷			BR: 33%
Pathway 3: 2 CH ₂ (OH)CH(OO•)CH(OH)(OH) → 2 CH ₂ (OH)CH(O•)CH(OH)(OH) + O ₂		1.7 10 ⁷			BR: 17%
CH ₂ (OH)CH(O•)CH(OH)(OH) → 0.50 C•H ₂ (OH) + 0.50 CHOCH(OH)(OH) + 0.50 C•H(OH)(OH) + 0.50 CH ₂ (OH)CHO					6 - 7
C•H ₂ (OH) + O ₂ → CH ₂ (OH)(OO•)		2.0·10 ⁹			5
C•H(OH)(OH) + O ₂ → CH(OH)(OH)(OO•)		2.0 10 ⁹			5
2 CH ₂ (OH)CH(OO•)CH(OH)(OH) → 1.33 CH ₂ (OH)COCH(OH)(OH) + 0.33 CH ₂ (OH)CH(OH)CH(OH)(OH) + 0.17 CHOCH(OH)(OH) + 0.17 CH ₂ (OH)CHO + 0.17 CH ₂ (OH)(OO•) + 0.17 CH(OH)(OH)(OO•) + 0.50 H ₂ O ₂ + 0.16 O ₂	R(600)	1.0 10 ⁸			= k(2 CH ₂ (OH)CH ₂ (OO*)) - 8
Oxidation of 2,3-dihydroxypropanal (or glyceraldehyde)					71
Pathway 1: CH ₂ (OH)CH(OH)CH(OH)(OH) + HO• → C•H(OH)CH(OH)CH(OH)(OH) + H ₂ O		8.3 10 ⁸			BR: 49% - 72
C•H(OH)CH(OH)CH(OH)(OH) + O ₂ → CH(OH)(OH)CH(OH)CH(OH)(OO•)		2.0·10 ⁹			5
Pathway 2: CH ₂ (OH)CH(OH)CH(OH)(OH) + HO• → CH ₂ (OH)CH(OH)CH(OH)(O•) + H ₂ O		4.6 10 ⁸			BR: 27% - 72
CH ₂ (OH)CH(OH)CH(OH)(O•) → CH ₂ (OH)C•H(OH) + CHO(OH)					6 - 7
CH ₂ (OH)C•H(OH) + O ₂ → CH ₂ (OH)CH(OH)(OO•)		2.0·10 ⁹			5
Pathway 3: CH ₂ (OH)CH(OH)CH(OH)(OH) + HO• → CH ₂ (OH)CH(OH)C•(OH)(OH) + H ₂ O		4.1 10 ⁸			BR: 24% - 72
CH ₂ (OH)CH(OH)C•(OH)(OH) + O ₂ → CH ₂ (OH)CH(OH)C(OH)(OH)(OO•)		2.0·10 ⁹			5
CH ₂ (OH)CH(OH)CH(OH)CH(OH)(OH) + HO• → 0.49 CH(OH)(OH)CH(OH)CH(OH)(OH)(OO•) + 0.27 CHO(OH) + 0.27 CH ₂ (OH)CH(OH)(OO•) + 0.24 CH ₂ (OH)CH(OH)C(OH)(OH)(OO•) + H ₂ O - O ₂	R(601)	1.7 10 ⁹			13
Pathway 1: CH ₂ (OH)CH(OH)CH(OH)CH(OH)(OH) + NO ₃ • → C•H(OH)CH(OH)CH(OH)CH(OH)(OH) + NO ₃ ⁻ + H ⁺		7.4 10 ⁵			BR: 67%
C•H(OH)CH(OH)CH(OH)CH(OH)(OH) + O ₂ → CH(OH)(OH)CH(OH)CH(OH)(OO•)		2.0·10 ⁹			5
Pathway 2: CH ₂ (OH)CH(OH)CH(OH)CH(OH)(OH) + NO ₃ • → CH ₂ (OH)CH(OH)C•(OH)(OH) + NO ₃ ⁻ + H ⁺		3.6 10 ⁵			BR: 33%
CH ₂ (OH)CH(OH)C•(OH)(OH) + O ₂ → CH ₂ (OH)CH(OH)C(OH)(OH)(OO•)		2.0·10 ⁹			5
CH ₂ (OH)CH(OH)CH(OH)CH(OH)(OH) + NO ₃ • → 0.67 CH(OH)(OH)CH(OH)CH(OH)(OO•) + 0.33 CH ₂ (OH)CH(OH)C(OH)(OH)(OO•) + NO ₃ ⁻ + H ⁺ - O ₂	R(602)	1.1 10 ⁶			= k(CH ₂ (OH)CH(OH)CH(OH)CH(OH) + NO ₃ •) - 2
CH(OH)(OH)CH(OH)CH(OH)(OO•) + OH• → CH(OH)(OH)CH(OH)CH(O•)(OO•) + H ₂ O		4.0 10 ⁹			
CH(OH)(OH)CH(OH)CH(O•)(OO•) → CHOCH(OH)CH(OH)(OH) + O ₂ •					9

Reactions		k_{298} ($M^{-n+1} s^{-1}$)	Ea/R (K)	References	Notes
$CH(OH)(OH)CH(OH)CH(OH)(OO^{\bullet}) + OH^{\bullet} \rightarrow CHOCH(OH)CH(OH)(OH) + O_2^{\bullet-} + H_2O$	R(603)	$4.0 \cdot 10^9$			$= k(CH_3CH(OH)(OO^{\bullet}) + OH^{\bullet})$
$CH(OH)(OH)CH(OH)CH(OH)(OO^{\bullet}) \rightarrow CHOCH(OH)CH(OH)(OH) + HO_2^{\bullet}$	R(604)	$1.9 \cdot 10^2$			10
$CH_2(OH)CH(OH)C(OH)(OH)(OO^{\bullet}) + OH^{\bullet} \rightarrow CH_2(OH)CH(OH)C(OH)(O^{\bullet})(OO^{\bullet}) + H_2O$		$4.0 \cdot 10^9$			9
$CH_2(OH)CH(OH)C(OH)(O^{\bullet})(OO^{\bullet}) \rightarrow CH_2(OH)CH(OH)CO(OH) + O_2^{\bullet-}$					
$CH_2(OH)CH(OH)C(OH)(OH)(OO^{\bullet}) + OH^{\bullet} \rightarrow CH_2(OH)CH(OH)CO(OH) + O_2^{\bullet-} + H_2O$	R(605)	$4.0 \cdot 10^9$			$= k(CH_3CH(OH)(OO^{\bullet}) + OH^{\bullet})$
$CH_2(OH)CH(OH)C(OH)(OH)(OO^{\bullet}) \rightarrow CH_2(OH)CH(OH)CO(OH) + HO_2^{\bullet}$	R(606)	$1.0 \cdot 10^6$			15
Oxidation of hydroxypropanedial					73
Pathway 1: $CH(OH)(OH)CH(OH)CH(OH)(OH) + HO^{\bullet} \rightarrow CH(OH)(OH)CH(OH)CH(OH)(O^{\bullet}) + H_2O$		$8.6 \cdot 10^8$			BR: 54% - 74
$CH(OH)(OH)CH(OH)CH(OH)(O^{\bullet}) \rightarrow CH(OH)(OH)C^{\bullet}H(OH) + CHO(OH)$					6 - 7
$CH(OH)(OH)C^{\bullet}H(OH) + O_2 \rightarrow CH(OH)(OO^{\bullet})CH(OH)(OH)$		$2.0 \cdot 10^9$			5
Pathway 2: $CH(OH)(OH)CH(OH)CH(OH)(OH) + HO^{\bullet} \rightarrow CH(OH)(OH)CH(OH)C^{\bullet}(OH)(OH) + H_2O$		$7.4 \cdot 10^8$			BR: 46% - 74
$CH(OH)(OH)CH(OH)C^{\bullet}(OH)(OH) + O_2 \rightarrow CH(OH)(OH)CH(OH)C(OH)(OH)(OO^{\bullet})$		$2.0 \cdot 10^9$			5
$CH(OH)(OH)CH(OH)CH(OH)(OH) + HO^{\bullet} \rightarrow 0.54 CHO(OH) + 0.54 CH(OH)(OO^{\bullet})CH(OH)(OH) + 0.46$	R(607)	$1.6 \cdot 10^9$			13
$CH(OH)(OH)CH(OH)C(OH)(OH)(OO^{\bullet}) + H_2O - O_2$					
$CH(OH)(OH)CH(OH)CH(OH)(OH) + NO_3^{\bullet} \rightarrow CH(OH)(OH)CH(OH)C^{\bullet}(OH)(OH) + NO_3^{\bullet-} + H^+$		$1.1 \cdot 10^6$			BR: 100%
$CH(OH)(OH)CH(OH)C^{\bullet}(OH)(OH) + O_2 \rightarrow CH(OH)(OH)CH(OH)C(OH)(OH)(OO^{\bullet})$		$2.0 \cdot 10^9$			5
$CH(OH)(OH)CH(OH)CH(OH)(OH) + NO_3^{\bullet} \rightarrow CH(OH)(OH)CH(OH)C(OH)(OH)(OO^{\bullet}) + NO_3^{\bullet-} + H^+ - O_2$	R(608)	$1.1 \cdot 10^6$			$= k(CH_2(OH)CH(OH)(OH) + NO_3^{\bullet}) - 2$
$CH(OH)(OH)CH(OH)C(OH)(OH)(OO^{\bullet}) + OH^{\bullet} \rightarrow CH(OH)(OH)CH(OH)C(OH)(O^{\bullet})(OO^{\bullet}) + H_2O$		$4.0 \cdot 10^9$			9
$CH(OH)(OH)CH(OH)C(OH)(O^{\bullet})(OO^{\bullet}) \rightarrow CH(OH)(OH)CH(OH)CO(OH) + O_2^{\bullet-}$					
$CH(OH)(OH)CH(OH)C(OH)(OH)(OO^{\bullet}) + OH^{\bullet} \rightarrow CH(OH)(OH)CH(OH)CO(OH) + O_2^{\bullet-} + H_2O$	R(609)	$4.0 \cdot 10^9$			$= k(CH_3CH(OH)(OO^{\bullet}) + OH^{\bullet})$
$CH(OH)(OH)CH(OH)C(OH)(OH)(OO^{\bullet}) \rightarrow CH(OH)(OH)CH(OH)CO(OH) + HO_2^{\bullet}$	R(610)	$1.0 \cdot 10^6$			15

1 - Branching ratios are calculated by the SAR from Monod and Doussin (2013): 16% for CH₃, 16% for CH₂, 65% for CH₂(OH) and 3% for OH. The 3 first pathways are considered corresponding to 97% of the total reactivity. They have been scaled to 15/15/70%.

2 - The oxidation by the radicals (NO₃[•], SO₄^{•-}, Cl[•], Cl₂^{•-}, CO₃^{•-}) is supposed to produce the same R(OO[•]) as the oxidation by HO[•] with the same branching ratios. The electron transfer pathways are not considered for these radicals. The H abstraction on an (OH) group by the NO₃[•] radical is also neglected because this reaction is thermodynamically disfavored.

3 - We assume the same branching ratio than for the self reaction of CH₃CH₂(OO[•]), *i.e.*, 80/20%. The "alkoxy" pathway (pathway 2) is more likely to occur (80%).

4 - DeCosta and Pincock (1989) showed that electron transfer proceeds with a rate constant around $1.0 \cdot 10^{10} s^{-1}$. We assumed that the electron transfer is non limiting.

5 - We assumed a fast rate constant equal to $2.0 \cdot 10^9 M^{-1} s^{-1}$ based on values compiled in Neta et al. (1990). This reaction is not a rate-determining step.

6 - Hilborn and Pincock (1991) showed that acyl alkoxyl radical RCO(O[•]) are fragmented with a rate constant around $1.0 \cdot 10^9 s^{-1}$. We assumed that the alkoxy fragmentation is non limiting.


7 - For alkoxyl radical, we assume an electron transfer reaction. When an oxygenated functional group is in β-position, we assume a fragmentation of the corresponding c-c bond. When there are two oxygenated function in β-position, we assume that the fragmentation occurs in priority on the C-CO(OH) bond.

8 - For self-reaction of peroxy radicals, we follow these similarity criteria:

Peroxy categories	Model compounds	References
>C(OO [•])CO(OH)/>C(OO [•])CO(O [•])	CH ₂ (OO [•])CO(O [•])	Schuchmann et al. (1985)
>C(OH)C(OO [•])<	CH ₂ (OH)CH ₂ (OO [•])	Piesiak et al. (1984)
>COC(OO [•])<	CH ₃ COCH ₂ (OO [•])	Zegota et al. (1986b)
Others	CH ₃ CH ₂ (OO [•])	Monod et al. (2007)

9 - Non-limiting reaction following Bothe et al. (1978).

10 - The HO₂[•] elimination rate constant depends on the substituent attached to the carbon atom bearing the peroxy function.



Von Sonntag (1987) compiled the following rate constants for :

R	R'	k (s ⁻¹)
H	H	<10
H	CH ₃	52
H	CH ₂ (OH)	190
CH ₃	CH ₃	665

For secondary carbon atom bearing the peroxy function, we assumed a rate of 665 s⁻¹.

11 - The hydration constant is calculated with the GROMHE method; $K_h = 2.49$. We consider the reactivity of the hydrated and non-hydrated forms since the hydrated form represents 71% of the total species.

12 - Branching ratios are calculated by the SAR from Doussin and Monod (2013) : 60% for CHO, 36% for CH₂(OH), 2% for (OH) and 2% for CH₂. The 2 first pathways are considered corresponding to 96% of the total reactivity. They have been scaled to 62/38%.

13 - Rate constant calculated from Doussin and Monod (2013).

14 - Branching ratios are calculated by the SAR from Doussin and Monod (2013) : 27% for H on CH(OH)(OH), 55% for CH₂ on CH₂(OH), 3% for (OH) on CH₂(OH), 2% for CH₂ and 13% for OH on CH(OH)(OH). The 2 first pathways are considered corresponding to 82% of the total reactivity. They have been scaled to 33/67%.

15 - Von Sonntag (1987) et Schuchmann & Von Sonntag (1988) have shown that the HO₂• elimination for RC(OH)(OH)(OO•) species is fast. This is confirmed by McElroy and Waygood (1991) for hydrated formaldehyde. We supposed a kinetic constant equal to 1.0 10⁶ s⁻¹.

16 - Branching ratios are calculated by the SAR from Doussin and Monod (2013) : 42% for CH₂, 38% for CH on CH(OH), 10% for CH₃, 5% for OH on CH₂(OH) and 5% for OH on CH(OH). The 2 first pathways are considered corresponding to 80% of the total reactivity. They have been scaled to 60/40%.

17 - The hydration constant is calculated with the GROMHE method; $K_h = 6.1$. We only consider the reactivity of the hydrated form since the hydrated form represents 86% of the total species.

18 - Branching ratios are calculated by the SAR from Doussin and Monod (2013) : 29% for OH on CH(OH)(OH), 28% for CH on CH(OH)(OH), 22% for CH on CH(OH), 14% for CH₃ and 7% for OH on CH(OH). The 3 first pathways are considered corresponding to 79% of the total reactivity. They have been scaled to 37/35/28%.

19 - The hydration constant is calculated with the GROMHE method. The K_h for the mono-hydrated and the di-hydrated form is equal to 22 and 100, respectively. The di-hydrated form represent 81% of the total species. We only consider the reactivity of di-hydrated form.

20 - Branching ratios are calculated by the SAR from Doussin and Monod (2013) : 32% for OH on CH(OH)(OH), 66% for CH on CH(OH)(OH), 2% for CH₂. The 2 first pathways are considered corresponding to 98% of the total reactivity. They have been scaled to 33/67%.

21 - Branching ratios are calculated by the SAR from Doussin and Monod (2013) : 91% for CH₂ on CH₂(OH), 7% for on OH CH₂(OH) and 2% for CH₂. The first pathway is considered corresponding to 91% of the total reactivity. It has been scaled to 100%.

22 - Branching ratios are calculated by the SAR from Doussin and Monod (2013) : 82% for CH₂ on CH₂(OH), 4% for OH on CH₂(OH), 3% for CH₂ and 11% for the electron-transfer on CO(O•). The first pathway is considered corresponding to 82% of the total reactivity. It has been scaled to 100%.

23 - Branching ratios are determined by Padmaja and Huie (1993): 87% for CH on CH(OH) and 13% for CH₃. Branching ratios calculated by the SAR from Doussin and Monod (2013) are close: 78% for CH on CH(OH) and 18% for CH₃, and 4% for OH on CH(OH). 4%.

24 - The hydration constant is equal to 0.87. The hydrated form represents 47% of the total species. We consider the reactivity of the hydrated and non-hydrated forms.

25 - Branching ratios are calculated by the SAR from Doussin and Monod (2013) : 82% for CHO, 12% for CH₃, 6% for CH₂. The 2 first pathways are considered corresponding to 94% of the total reactivity. They have been scaled to 87/13%.

26 - Branching ratios are calculated by the SAR from Doussin and Monod (2013) : 47% for CH on CH(OH)(OH), 23% for CH₃, 21% for OH on CH(OH)(OH), 9% for CH₂. The 3 first pathways are considered corresponding to 91% of the total reactivity. They have been scaled to 52/25/23%.

27 - The mono-hydrated form CH₃COCH(OH)(OH) and the di-hydrated form represent respectively 66% and 32% of the total species (the non-hydrated + the two mono-hydrated + the di-hydrated forms). We only consider the reactivity of this mono-hydrated form and of the di-hydrated form.

28 - Branching ratios are calculated by the SAR from Doussin and Monod (2013) : 57% for OH on CH(OH)(OH), 29% for CH on CH(OH)(OH), 14% for CH₃. The 3 pathways are considered corresponding to 100% of the total reactivity.

29 - Branching ratios are calculated by the SAR from Doussin and Monod (2013) : 41% for OH on C(OH)(OH), 37% for OH on CH(OH)(OH), 15% for CH on CH(OH)(OH) and 7% for CH₃. The 2 first pathways are considered corresponding to 78% of the total reactivity. They have been scaled to 53/47%.

30 - The non-hydrated form and the mono-hydrated form on the ketone function are negligible. The mono-hydrated form CH₂(OH)COCH(OH)(OH) and the di-hydrated form CH₂(OH)C(OH)(OH)CH(OH)(OH) are the main forms (respectively 18% and 81% of the total species). The reactivity of these two forms are considered.

31 - Branching ratios are calculated by the SAR from Doussin and Monod (2013) : 37% for OH on CH(OH)(OH), 36% for CH₂ on CH₂(OH), 18% for CH on CH(OH)(OH), 9% for (OH) on CH₂(OH). The 3 first pathways are considered corresponding to 91% of the total reactivity. They have been scaled to 41/40/19%.

32 - Branching ratios are calculated by the SAR from Doussin and Monod (2013) : 31% for OH on C(OH)(OH), 29% for OH on CH(OH)(OH), 22% for CH₂ on CH₂(OH), 11% for CH on CH(OH)(OH) and 7% for OH on CH₂(OH). The 3 first pathways are considered corresponding to 82% of the total reactivity. They have been scaled to 38/35/27%.

33 - The tri-hydrated form is dominant (97%). We only consider its reactivity. This compound is also produced under its mono-hydrated form in the oxidation of methyl glyoxal and 2-oxo,3-hydroxy propanal.

34 - Branching ratios are calculated by the SAR from Doussin and Monod (2013) : 54% for OH on CH(OH)(OH), 27% for OH on C(OH)(OH), 19% for CH on CH(OH)(OH). The 2 first pathways are considered corresponding to 81% of the total reactivity. They have been scaled to 67/33%.

35 - $K_h = 2.0 \cdot 10^{-3}$ (Bell, 1966). The hydrated form is not significant. Only the reactivity of the non-hydrated is considered. The equilibrium is not considered because the hydrated form cannot be produced through reactivity.

36 - $K_h = 7.0 \cdot 10^{-2}$ estimated by GROMHE. The hydrated form is not significant. Only the reactivity of the non-hydrated is considered. The equilibrium is not considered because the hydrated form cannot be produced through reactivity.

37 - Branching ratios are calculated by the SAR from Doussin and Monod (2013) : 16% for CH₃, 69% for CH₂ on CH₂(OH), 15% for OH on CH₂(OH). The 3 first pathways are considered corresponding to 100% of the total reactivity.

38 - $K_h = 0.32$ estimated by GROMHE. The hydrated form represent 24% of the total species. Only the reactivity of the non-hydrated is considered.

39 - Branching ratios are calculated by the SAR from Doussin and Monod (2013) : 81% for CH₂, 19% for OH on CH₂(OH). The first pathway is considered corresponding to 81% of the total reactivity. The branching ratio is scaled to 100%.

40 - Branching ratios are calculated by the SAR from Doussin and Monod (2013) : 17% for CH₂, 83% for CH₃. The two pathways are considered corresponding to 100% of the total reactivity.

41 - Branching ratios are calculated by the SAR from Doussin and Monod (2013) : 28% for CH₂, 72% for CH₃. The two pathways are considered corresponding to 100% of the total reactivity.

42 - The hydration constant of the acid form CO(OH)CH₂CHO is equal to 10 by GROMHE. The hydrated form represents 91% of the total species. We only consider the reactivity of the hydrated form. For CO(O⁻)CH₂CHO, the hydration constant is 1.4 estimated by GROMHE. The hydrated form represents 58% of the total species. We consider the reactivity of the hydrated and non-hydrated forms.

43 - Branching ratios are calculated by the SAR from Doussin and Monod (2013) : 57% for CH on CH(OH)(OH), 42% for OH on CH(OH)(OH) and 1% for CH₂. The two pathways are considered corresponding to 99% of the total reactivity. They have been scaled to 58/42%.

44 - Branching ratios are calculated by the SAR from Doussin and Monod (2013) : 59% for CH on CH(OH)(OH), 25% for OH on CH(OH)(OH), 16% for the electron-transfer on CO(O⁻). The 3 pathways are considered corresponding to 100% of the total reactivity.

45 - Branching ratios are calculated by the SAR from Doussin and Monod (2013) : 91% for CHO, 8% for the electron-transfer on CO(O⁻), 1% for CH₂. The first pathway is considered corresponding to 91% of the total reactivity.

46 - Branching ratios are calculated by the SAR from Doussin and Monod (2013) : 6% for CH₂ and 94% for the electron-transfer on CO(O⁻). The 2 pathways are considered corresponding to 100% of the total reactivity.

47 - Branching ratios are calculated by the SAR from Doussin and Monod (2013) : 6% for CH₂ and 94% for the electron-transfer on CO(O⁻). The 2 pathways are considered corresponding to 100% of the total reactivity.

48 - Branching ratios are calculated by the SAR from Doussin and Monod (2013) : 11% for CH on CH(OH) and 89% for OH on CH(OH). The 2 pathways are considered corresponding to 100% of the total reactivity.

49 - Branching ratios are calculated by the SAR from Doussin and Monod (2013) : 11% for CH on CH(OH), 43% for OH on CH(OH) and 46% for the electron-transfer on CO(O⁻). The 3 pathways are considered corresponding to 100% of the total reactivity.

50 - Branching ratios are calculated by the SAR from Doussin and Monod (2013) : 13% for CH on CH(OH), 27% for OH on CH(OH) and 60% for the electron-transfer on CO(O⁻). The 3 pathways are considered corresponding to 100% of the total reactivity.

51 - The hydration constant of the acid form CO(OH)COCO(OH) is measured at 100 by Le Henaff (1968). We only consider the reactivity of the hydrated form. For the mono-anion CO(OH)COCO(O⁻), the hydration constant is equal to 2.4; the hydrated form represents 71% of the total species. Therefore, we consider the reactivity of the hydrated and non-hydrated forms. For the di-anion CO(O⁻)COCO(O⁻), the hydration constant is low and estimated at $1.6 \cdot 10^{-2}$ by GROMHE. We only consider the reactivity of the non-hydrated form.

52 - Branching ratios are calculated by the SAR from Doussin and Monod (2013) : 34% for CH on CH(OH), 44% for CH₃ and 22% for OH on CH(OH). The 3 pathways are considered corresponding to 100% of the total reactivity.

53 - Branching ratios are calculated by the SAR from Doussin and Monod (2013) : 42% for CH on CH(OH), 28% for CH₃, 14% for OH on CH(OH) and 16% for the electron-transfer on CO(O⁻). The 4 pathways are considered corresponding to 100% of the total reactivity.

54 - Branching ratios are calculated by the SAR from Doussin and Monod (2013) : 72% for CH₂, 13% for OH on CH₂(OH), 9% for OH on CH(OH) and 6% for CH on CH(OH). The two pathways are considered corresponding to 85% of the total reactivity. They have been scaled to 85/15%.

55 - Branching ratios are calculated by the SAR from Doussin and Monod (2013) : 66% for CH₂, 10% for CH on CH(OH), 9% for the electron-transfer on CO(O⁻), 8% for OH on CH(OH) and 7% OH on CH₂(OH). The two pathways are considered corresponding to 76% of the total reactivity. They have been scaled to 87/13%.

56 - $K_h = 46$ for the acid form CHOCH(OH)CO(OH) estimated by GROMHE. The hydrated form represent 98% of the total species. Only the reactivity of the hydrated form is considered. $K_h = 6.1$ for the anion $\text{CHOCH(OH)CO(O}^-)$ estimated by GROMHE. The hydrated form represent 86% of the total species. Only the reactivity of the hydrated is considered.

57 - Branching ratios are calculated by the SAR from Doussin and Monod (2013) : 57% for OH on CH(OH)(OH) , 33% for CH on CH(OH)(OH) , 8% for OH on CH(OH) , 2% for CH on CH(OH) . The two pathways are considered corresponding to 90% of the total reactivity. They have been scaled to 63/37%.

58 - Branching ratios are calculated by the SAR from Doussin and Monod (2013) : 37% for OH on CH(OH)(OH) , 38% for CH on CH(OH)(OH) , 10% for the electron-transfer on $\text{CO(O}^-)$, 10% for OH on CH(OH) , 5% for CH on CH(OH) . The two pathways are considered corresponding to 75% of the total reactivity. They have been scaled to 49/51%.

59 - For the acid form, the di-hydrate is the main form (99%) due to its high K_h ($1.3 \cdot 10^5$ calculated with GROMHE). Only the reactivity of the di-hydrated form is considered. For the anionic form, the mono-hydrate $\text{CO(O}^-)\text{COCH(OH)(OH)}$ is dominant (49%) together with the di-hydrated form (49%). Only the reactivity of the di-hydrated form and of one of the mono-hydrated form ($\text{CO(O}^-)\text{COCH(OH)(OH)}$) is considered.

60 - Branching ratios are calculated by the SAR from Doussin and Monod (2013) : 52% for OH on CH(OH)(OH) , 35% for OH on C(OH)(OH) , 13% for CH on CH(OH)(OH) . The three pathways are considered corresponding to 100% of the total reactivity.

61 - Branching ratios are calculated by the SAR from Doussin and Monod (2013) : 60% for OH on CH(OH)(OH) , 32% for CH on CH(OH)(OH) , 8% for the electron-transfer on $\text{CO(O}^-)$. The two pathways are considered corresponding to 92% of the total reactivity. They have been scaled to 65/35%.

62 - Branching ratios are calculated by the SAR from Doussin and Monod (2013) : 37% for OH on CH(OH)(OH) , 43% for OH on C(OH)(OH) , 16% for CH on CH(OH)(OH) and 4% for the electron-transfer on $\text{CO(O}^-)$. The three first pathways are considered corresponding to 96% of the total reactivity. They have been scaled to 39/45/16%.

63 - For the acid form, the hydrated form is the main form (92%) due to its high K_h (10.8 calculated with GROMHE). Only the reactivity of the hydrated form is considered. For the anionic form, the hydrated form is negligible (7%). Only the reactivity of the non-hydrated form is considered.

64 - Branching ratios are calculated by the SAR from Doussin and Monod (2013) : 48% for OH on C(OH)(OH) , 36% for CH_2 on $\text{CH}_2(\text{OH})$, 16% for OH on $\text{CH}_2(\text{OH})$. The three pathways are considered corresponding to 100% of the total reactivity.

65 - Branching ratios are calculated by the SAR from Doussin and Monod (2013) : 75% for CH_2 on $\text{CH}_2(\text{OH})$, 15% for OH on $\text{CH}_2(\text{OH})$ and 10% for the electron-transfer on $\text{CO(O}^-)$. The first pathway is considered corresponding to 75% of the total reactivity. This has been scaled to 100%.

66 - For the acid form, $K_h = 1.5$ following Poker et al. (1969). The hydrated form represents 60% of the total species. We consider the reactivity of the hydrated and non-hydrated forms. For the anion, $K_h = 5.7 \cdot 10^{-2}$ from Poker et al. (1969). Therefore, we only consider the reactivity of the non-hydrated form.

67 - We suppose this reaction to be non limiting.

68 - Branching ratios are calculated by the SAR from Doussin and Monod (2013) : 85% for OH on C(OH)(OH) , 15% for CH_3 . The two pathways are considered corresponding to 100% of the total reactivity.

69 - Acrolein establishes a very slow equilibrium with 3-hydroxypropionaldehyde following Pressmann and Lucas (1942).

70 - The addition of OH on the internal carbon of the $\text{C}=\text{C}$ bond is highly favored (>80%). Von Sonntag et Schuchmann (1997) have shown that for ethene, this pathway is the most efficient.

71 - $K_h = 11.2$ estimated by GROMHE. The hydrated form represent the main form of the total species. Only the reactivity of the hydrated form is considered.

72 - Branching ratios are calculated by the SAR from Doussin and Monod (2013) : 41% for CH_2 , 22% for OH on CH(OH)(OH) , 20% for CH on CH(OH)(OH) , 7% for CH on CH(OH) , 5% for OH on $\text{CH}_2(\text{OH})$ and 5% for OH on CH(OH) . The three first pathways are considered corresponding to 83% of the total reactivity. They have been scaled to 49/27/24%.

73 - The hydration constant is calculated with the GROMHE method. The K_h for the mono-hydrated and the di-hydrated form is equal to 98 and $2.0 \cdot 10^3$, respectively. The di-hydrated form represent 95% of the total species. We only consider the reactivity of di-hydrated form.

74 - Branching ratios are calculated by the SAR from Doussin and Monod (2013) : 49% for OH on CH(OH)(OH) , 42% for CH on CH(OH)(OH) , 5% for OH on CH(OH) , 3% for CH on CH(OH) . The two first pathways are considered corresponding to 91% of the total reactivity. They have been scaled to 54/46%.

References:

- Bell, R. P.: The reversible hydration of carbonyl compounds, in: *Advances in Physical Organic Chemistry*, Elsevier Inc ed., V. Gold, Londres (Royaume-Uni), 1-29, 1966.
- Bothe, E., Schuchmann, M. N., Schulte-Frohlinde, D., and Von Sonntag, C.: HO2 Elimination from α -hydroxyalkylperoxy radicals in aqueous solution, *Photochem. Photobiol.*, 28, 639-643, 1978.
- DeCosta, D. P., and Pincock, J. A.: Control of product distribution by Marcus type electron-transfer rates for the radical pair generated in benzylic ester photochemistry, *J. Am. Chem. Soc.*, 111, 8948-8950, 1989.
- Doussin, J. F., and Monod, A.: Structure–activity relationship for the estimation of OH-oxidation rate constants of carbonyl compounds in the aqueous phase, *Atmos. Chem. Phys.*, 13, 11625-11641, 10.5194/acp-13-11625-2013, 2013.
- Hilborn, J. W., and Pincock, J. A.: Rates of decarboxylation of acyloxy radicals formed in the photocleavage of substituted 1-naphthylmethyl alkanoates, *J. Am. Chem. Soc.*, 113, 2683-2686, 1991.
- Le Henaff, P.: Methodes d'étude et propriétés des hydrates hémiacétals et hémithioacétals dérivés des aldéhydes et cétones, *P. Bull. Soc. Chim. Fr.*, 11, 4687-4698, 1968.
- McElroy, W. J., and Waygood, S. J.: Oxidation of formaldehyde by the hydroxyl radical in aqueous solution, *J. Chem. Soc. Far. Trans.*, 87, 1513-1521, 1991.
- Monod, A., Chevallier, E., Durand Jolibois, R., Doussin, J. F., Picquet-Varrault, B., and Carlier, P.: Photooxidation of methylhydroperoxide and ethylhydroperoxide in the aqueous phase under simulated cloud droplet conditions, *Atmos. Environ.*, 41, 2412-2426, 2007.

-
- Neta, P., Huie, R., and Ross, A.: Rate constants for reactions of peroxy radicals in fluid solutions, J. Phys. Chem. Ref. Data, 19, 413, 1990.
- Padmaja, S., Neta, P., and Huie, R. E.: Rate constant for some reactions of inorganic radicals with inorganic ions. Temperature dependence and solvent dependence, J. Phys. Chem., 25, 445-455, 1993.
- Piesiak, A., Schuchmann, M. N., Zegota, H., and von Sonntag, C.: β -Hydroxyethylperoxy radicals: A study of the γ -radiolysis and pulse radiolysis of ethylene in oxygenated aqueous solutions, Z. Naturforsch., 39b, 1262-1267, 1984.
- Pocker, Y., Meany, J. E., Nist, B. J., and Zadorojny, C.: Reversible hydration of pyruvic acid. I. Equilibrium studies, J. Phys. Chem., 73, 2879-2882, 1969.
- Pressman, D., and Lucas, H. J.: Hydration of unsaturated compounds. XI. Acrolein and acrylic acid, J. Am. Chem. Soc., 64, 1953-1957, 1942.
- Schuchmann, M. N., Zegota, H., and Von Sonntag, C.: Acetate peroxy radicals $\cdot\text{O}_2\text{CH}_2\text{CO}_2^-$: A study on the γ -radiolysis and pulse radiolysis of acetate in oxygenated aqueous solutions, Z. Naturforsch., 40b, 215-221, 1985.
- Von Sonntag, C.: The chemical basis of radiation biology, Taylor & Francis, 1987.
- Von Sonntag, C., Dowideit, P., Fang, X., Mertens, R., Pan, X., Schuchmann, M. N., and Schuchmann, H.-P.: The fate of peroxy radicals in aqueous solution, W. Sci. and Tech., 35, 9-15, 1997.
- Schuchmann, M. N., and Von Sonntag, C.: The rapid hydration of the acetyl radical. A pulse radiolysis study of acetaldehyde in aqueous solution, J. Am. Chem. Soc., 110, 5698-5701, 1988.
- Zegota, H., Schuchmann, M. N., and Von Sonntag, C.: Elucidation of the mechanism of peroxy radical reactions in aqueous solutions using the pulse radiolysis technique, J. Radioanal. Nucl. Chem., 101, 199-207, 1986b.

Equilibria

Species		K _a or K _h	-ΔH/R (K)	References	Notes
C3 compounds					
3-hydroxypropionaldehyde					
CH ₂ (OH)CH ₂ CHO + H ₂ O ↔ CH ₂ (OH)CH ₂ CH(OH)(OH)	T(48)	2.5		Estimated with GROMHE	
CH ₂ (OH)CH ₂ CO(OO [•]) + H ₂ O ↔ CH ₂ (OH)CH ₂ C(OH)(OH)(OO [•])	T(49)	2.5			1 = K _h (CH ₂ (OH)CH ₂ CHO/CH ₂ (OH)CH ₂ CH(OH)(OH))
CHOCH ₂ CH(OH)(OO [•]) + H ₂ O ↔ CH(OH)(OH)CH ₂ CH(OH)(OO [•])	T(50)	2.5			1 = K _h (CH ₂ (OH)CH ₂ CHO/CH ₂ (OH)CH ₂ CH(OH)(OH))
2-hydroxypropionaldehyde					
CH ₃ CH(OH)CHO + H ₂ O ↔ CH ₃ CH(OH)CH(OH)(OH)	T(51)	6.1		Estimated with GROMHE	
CH ₃ CH(OH)CO(OO [•]) + H ₂ O ↔ CH ₃ CH(OH)C(OH)(OH)(OO [•])	T(52)	1.0 10 ⁻³			2
CH ₃ C(OH)(OO [•])CHO + H ₂ O ↔ CH ₃ C(OH)(OO [•])CH(OH)(OH)	T(53)	6.1			1 = K _h (CH ₃ CH(OH)CHO/CH ₃ CH(OH)CH(OH)(OH))
Propanedial					
CHOCH ₂ CHO + H ₂ O ↔ CHOCH ₂ CH(OH)(OH)	T(54)	2.2 10 ¹		Estimated with GROMHE	
CHOCH ₂ CHO + 2 H ₂ O ↔ CH(OH)(OH)CH ₂ CH(OH)(OH)	T(55)	1.0 10 ²		Estimated with GROMHE	
CHOCH ₂ CO(OO [•]) + H ₂ O ↔ CHOCH ₂ C(OH)(OH)(OO [•])	T(56)	1.0 10 ⁻³			2
CHOCH ₂ CO(OO [•]) + H ₂ O ↔ CH(OH)(OH)CH ₂ CO(OO [•])	T(57)	2.2 10 ¹			1 = K _h (CHOCH ₂ CHO/CHOCH ₂ CH(OH)(OH)) - 3
CHOCH ₂ CO(OO [•]) + 2 H ₂ O ↔ CH(OH)(OH)CH ₂ C(OH)(OH)(OO [•])	T(58)	1.0 10 ²			1 = K _h (CHOCH ₂ CHO/CH(OH)(OH)CH ₂ CH(OH)(OH))
3-hydroxypropionic acid					
CH ₂ (OH)CH ₂ CO(OH) ↔ CH ₂ (OH)CH ₂ CO(O ⁻) + H ⁺	T(59)	3.1 10 ⁻⁵		Lide, 2005	
CH(OH)(OO [•])CH ₂ CO(OH) ↔ CH(OH)(OO [•])CH ₂ CO(O ⁻) + H ⁺	T(60)	3.1 10 ⁻⁵			4 = K _a (CH ₂ (OH)CH ₂ CO(OH)/CH ₂ (OH)CH ₂ CO(O ⁻))
Propionaldehyde					
CH ₃ CH ₂ CHO + H ₂ O ↔ CH ₃ CH ₂ CH(OH)(OH)	T(61)	0.87	2715	Socrates, 1969	
CH ₃ CH ₂ CO(OO [•]) + H ₂ O ↔ CH ₃ CH ₂ C(OH)(OH)(OO [•])	T(62)	0.87			1 = K _h (CH ₃ CH ₂ CHO/CH ₃ CH ₂ CH(OH)(OH))
CHOCH ₂ CH ₂ (OO [•]) + H ₂ O ↔ CH(OH)(OH)CH ₂ CH ₂ (OO [•])	T(63)	0.87			1 = K _h (CH ₃ CH ₂ CHO/CH ₃ CH ₂ CH(OH)(OH))
Methylglyoxal					
CH ₃ COCHO + H ₂ O ↔ CH ₃ COCH(OH)(OH)	T(64)	1.8 10 ³		Wasa and Musha, 1970	5
CH ₃ COCHO + H ₂ O ↔ CH ₃ C(OH)(OH)CHO	T(65)	5.4 10 ¹		Wasa and Musha, 1970	5
CH ₃ COCHO + 2 H ₂ O ↔ CH ₃ C(OH)(OH)CH(OH)(OH)	T(66)	8.7 10 ²		Wasa and Musha, 1970	5

Species		K _a or K _h	-ΔH/R (K)	References	Notes
CH ₃ COCO(OO•) + H ₂ O ↔ CH ₃ COC(OH)(OH)(OO•)	T(67)	1.0 10 ⁻³			2
CH ₃ COCO(OO•) + H ₂ O ↔ CH ₃ C(OH)(OH)CO(OO•)	T(68)	5.4 10 ¹			1 = K _h (CH ₃ COCHO/CH ₃ C(OH)(OH)CHO)
CH ₃ COCO(OO•) + 2 H ₂ O ↔ CH ₃ C(OH)(OH)C(OH)(OH)(OO•)	T(69)	8.7 10 ²			1 = K _h (CH ₃ COCHO/CH ₃ C(OH)(OH)CH(OH)(OH))
CHOCOCH ₂ (OO•) + H ₂ O ↔ CH(OH)(OH)COCH ₂ (OO•)	T(70)	1.8 10 ³			1 = K _h (CH ₃ COCHO/CH ₃ COCH(OH)(OH))
CHOCOCH ₂ (OO•) + H ₂ O ↔ CHOC(OH)(OH)CH ₂ (OO•)	T(71)	5.4 10 ¹			1 = K _h ([CH ₃ COCHO/CH ₃ C(OH)(OH)CHO])
CHOCOCH ₂ (OO•) + 2 H ₂ O ↔ CH(OH)(OH)C(OH)(OH)CH ₂ (OO•)	T(72)	8.7 10 ²			1 = K _h ([CH ₃ COCHO/CH ₃ C(OH)(OH)CH(OH)(OH)])
2-oxo, 3-hydroxypropanal					
CH ₂ (OH)COCHO + H ₂ O ↔ CH ₂ (OH)COCH(OH)(OH)	T(73)	2.0 10 ²		Estimated with GROMHE	
CH ₂ (OH)COCHO + H ₂ O ↔ CH ₂ (OH)C(OH)(OH)CHO	T(74)	1.3 10 ¹		Estimated with GROMHE	
CH ₂ (OH)COCHO + 2 H ₂ O ↔ CH ₂ (OH)C(OH)(OH)CH(OH)(OH)	T(75)	9.2 10 ²		Estimated with GROMHE	
CH ₂ (OH)COCO(OO•) + H ₂ O ↔ CH ₂ (OH)COC(OH)(OH)(OO•)	T(76)	1.0 10 ⁻³			2
CH ₂ (OH)COCO(OO•) + H ₂ O ↔ CH ₂ (OH)C(OH)(OH)CO(OO•)	T(77)	1.3 10 ¹			1 = K _h (CH ₂ (OH)COCHO/CH ₂ (OH)C(OH)(OH)CHO)
CH ₂ (OH)COCO(OO•) + 2 H ₂ O ↔ CH ₂ (OH)C(OH)(OH)C(OH)(OH)(OO•)	T(78)	9.2 10 ²			1 = K _h (CH ₂ (OH)COCHO/CH ₂ (OH)C(OH)(OH)CH(OH)(OH))
CHOCOCH(OH)(OO•) + H ₂ O ↔ CH(OH)(OH)COCH(OH)(OO•)	T(79)	2.0 10 ²			1 = K _h (CH ₂ (OH)COCHO/CH ₂ (OH)COCH(OH)(OH))
CHOCOCH(OH)(OO•) + H ₂ O ↔ CHOC(OH)(OH)CH(OH)(OO•)	T(80)	1.3 10 ¹			1 = K _h (CH ₂ (OH)COCHO/CH ₂ (OH)C(OH)(OH)CHO)
CHOCOCH(OH)(OO•) + 2 H ₂ O ↔ CH(OH)(OH)C(OH)(OH)CH(OH)(OO•)	T(81)	9.2 10 ²			1 = K _h (CH ₂ (OH)COCHO/CH ₂ (OH)C(OH)(OH)CH(OH)(OH))
Oxopropanedial					
CHOCOCHO + H ₂ O ↔ CHOCOCH(OH)(OH)	T(82)	1.7 10 ³		Estimated with GROMHE	
CHOCOCHO + H ₂ O ↔ CHOC(OH)(OH)CHO	T(83)	5.1 10 ²		Estimated with GROMHE	
CHOCOCHO + 2 H ₂ O ↔ CH(OH)(OH)COCH(OH)(OH)	T(84)	6.3 10 ⁵		Estimated with GROMHE	
CHOCOCHO + 2 H ₂ O ↔ CHOC(OH)(OH)CH(OH)(OH)	T(85)	3.3 10 ⁵		Estimated with GROMHE	
CHOCOCHO + 3 H ₂ O ↔ CH(OH)(OH)C(OH)(OH)CH(OH)(OH)	T(86)	3.4 10 ⁷		Estimated with GROMHE	
CHOCOCO(OO•) + H ₂ O ↔ CHOCOC(OH)(OH)(OO•)	T(87)	1.0 10 ⁻³			2

Species		K _a or K _h	-ΔH/R (K)	References	Notes
CHOCOCO(OO [•]) + H ₂ O ↔ CHOC(OH)(OH)CO(OO [•])	T(88)	5.1 10 ²			1 = K _h (CHOCOCHO/CHOC(OH)(OH)CHO)
CHOCOCO(OO [•]) + 2 H ₂ O ↔ CH(OH)(OH)COC(OH)(OH)(OO [•])	T(89)	6.3 10 ⁵			1 = K _h (CHOCOCHO/CH(OH)(OH)COCH(OH)(OH))
CHOCOCO(OO [•]) + 2 H ₂ O ↔ CHOC(OH)(OH)C(OH)(OH)(OO [•])	T(90)	3.3 10 ⁵			1 = K _h (CHOCOCHO/CHOC(OH)(OH)CH(OH)(OH))
CHOCOCO(OO [•]) + 2 H ₂ O ↔ CH(OH)(OH)C(OH)(OH)CO(OO [•])	T(91)	3.3 10 ⁵			1 = K _h (CHOCOCHO/CHOC(OH)(OH)CH(OH)(OH))
CHOCOCO(OO [•]) + 3 H ₂ O ↔ CH(OH)(OH)C(OH)(OH)C(OH)(OH)(OO [•])	T(92)	3.4 10 ⁷			1 = K _h (CHOCOCHO/CH(OH)(OH)C(OH)(OH)CH(OH)(OH))
Dihydroxyacetone					
CH ₂ (OH)COCH ₂ (OH) + H ₂ O ↔ CH ₂ (OH)C(OH)(OH)CH ₂ (OH)	T(93)	3.2 10 ⁻¹		Estimated with GROMHE	
CH ₂ (OH)COCH(OH)(OO [•]) + H ₂ O ↔ CH ₂ (OH)C(OH)(OH)CH(OH)(OO [•])	T(94)	3.2 10 ⁻¹			1 = K _h (CH ₂ (OH)COCH ₂ (OH)/CH ₂ (OH)C(OH)(OH)CH ₂ (OH))
Propionic acid					
CH ₃ CH ₂ CO(OH) ↔ CH ₃ CH ₂ CO(O ⁻) + H ⁺	T(95)	1.3·10 ⁻⁵		Lide, 2005	
CH ₃ CH(OO [•])CO(OH) ↔ CH ₃ CH(OO [•])CO(O ⁻) + H ⁺	T(96)	1.3·10 ⁻⁵			4 = K _a (CH ₃ CH ₂ CO(OH)/CH ₃ CH ₂ CO(O ⁻))
CH ₂ (OO [•])CH ₂ CO(OH) ↔ CH ₂ (OO [•])CH ₂ CO(O ⁻) + H ⁺	T(97)	1.3·10 ⁻⁵			4 = K _a (CH ₃ CH ₂ CO(OH)/CH ₃ CH ₂ CO(O ⁻))
3-oxopropionic acid					
CO(OH)CH ₂ CHO ↔ CO(O ⁻)CH ₂ CHO + H ⁺	T(98)	1.3·10 ⁻⁵			= K _a (CH ₃ CH ₂ CO(OH)/CH ₃ CH ₂ CO(O ⁻))
CO(OH)CH ₂ CHO + H ₂ O ↔ CO(OH)CH ₂ CH(OH)(OH)	T(99)	1.0 10 ¹		Estimated with GROMHE	
CO(O ⁻)CH ₂ CHO + H ₂ O ↔ CO(O ⁻)CH ₂ CH(OH)(OH)	T(100)	1.4		Estimated with GROMHE	
CO(OH)CH ₂ C(OH)(OH)(OO [•]) ↔ CO(O ⁻)CH ₂ C(OH)(OH)(OO [•]) + H ⁺	T(101)	1.3·10 ⁻⁵			4 = K _a (CO(OH)CH ₂ CHO/CO(O ⁻)CH ₂ CHO)
CO(OH)CH ₂ CO(OO [•]) + H ₂ O ↔ CO(OH)CH ₂ C(OH)(OH)(OO [•])	T(102)	1.0 10 ⁻³			2
CO(O ⁻)CH ₂ CO(OO [•]) + H ₂ O ↔ CO(O ⁻)CH ₂ C(OH)(OH)(OO [•])	T(103)	1.0 10 ⁻³			2
Malonic acid					
CO(OH)CH ₂ CO(OH) ↔ CO(OH)CH ₂ CO(O ⁻) + H ⁺	T(104)	1.4·10 ⁻³		Lide, 2005	
CO(OH)CH ₂ CO(O ⁻) ↔ CO(O ⁻)CH ₂ CO(O ⁻) + H ⁺	T(105)	2.0·10 ⁻⁶		Lide, 2005	
CO(OH)CH(OO [•])CO(OH) ↔ CO(OH)CH(OO [•])CO(O ⁻) + H ⁺	T(106)	1.4·10 ⁻³			4 = K _a (CO(OH)CH ₂ CO(OH)/CO(OH)CH ₂ CO(O ⁻))
CO(OH)CH(OO [•])CO(O ⁻) ↔ CO(O ⁻)CH(OO [•])CO(O ⁻) + H ⁺	T(107)	2.0·10 ⁻⁶			4 = K _a (CO(OH)CH ₂ CO(O ⁻)/[CO(O ⁻)CH ₂ CO(O ⁻)])
Tartronic acid					
CO(OH)CH(OH)CO(OH) ↔ CO(OH)CH(OH)CO(O ⁻) + H ⁺	T(108)	3.8·10 ⁻³		Lide, 2005	
CO(OH)CH(OH)CO(O ⁻) ↔ CO(O ⁻)CH(OH)CO(O ⁻) + H ⁺	T(109)	2.9·10 ⁻⁵		Lide, 2005	

Species		K _a or K _b	-ΔH/R (K)	References	Notes
CO(OH)C(OH)(OO [•])CO(OH) ↔ CO(OH)C(OH)(OO [•])CO(O ⁻) + H ⁺	T(110)	3.8·10 ⁻³			4 = K _a (CO(OH)CH(OH)CO(OH)/CO(OH)CH(OH)CO(O ⁻))
CO(OH)C(OH)(OO [•])CO(O ⁻) ↔ CO(O ⁻)C(OH)(OO [•])CO(O ⁻) + H ⁺	T(111)	2.9·10 ⁻⁵			4 = K _a (CO(OH)CH(OH)CO(O ⁻)/CO(O ⁻)CH(OH)CO(O ⁻))
Mesoxalic acid					
CO(OH)C(OH)(OH)CO(OH) ↔ CO(OH)C(OH)(OH)CO(O ⁻) + H ⁺	T(112)	3.2·10 ⁻³		Albalat et al., 1989	
CO(OH)C(OH)(OH)CO(O ⁻) ↔ CO(O ⁻)C(OH)(OH)CO(O ⁻) + H ⁺	T(113)	1.3·10 ⁻⁴		Albalat et al., 1989	
CO(OH)COCO(OH) + H ₂ O ↔ CO(OH)C(OH)(OH)CO(OH)	T(114)	1.0 10 ²		Le Henaff, 1968	
CO(OH)COCO(O ⁻) + H ₂ O ↔ CO(OH)C(OH)(OH)CO(O ⁻)	T(115)	2.4		Estimated with GROMHE	
CO(O ⁻)COCO(O ⁻) + H ₂ O ↔ CO(O ⁻)C(OH)(OH)CO(O ⁻)	T(116)	1.6 10 ⁻²		Estimated with GROMHE	
Lactic acid					
CH ₃ CH(OH)CO(OH) ↔ CH ₃ CH(OH)CO(O ⁻) + H ⁺	T(117)	1.4·10 ⁻⁴		Lide, 2005	
CH ₃ C(OH)(OO [•])CO(OH) ↔ CH ₃ C(OH)(OO [•])CO(O ⁻) + H ⁺	T(118)	1.4·10 ⁻⁴			4 = K _a (CH ₃ CH(OH)CO(OH)/CH ₃ CH(OH)CO(O ⁻))
CH ₂ (OO [•])CH(OH)CO(OH) ↔ CH ₂ (OO [•])CH(OH)CO(O ⁻) + H ⁺	T(119)	1.4·10 ⁻⁴			4 = K _a (CH ₃ CH(OH)CO(OH)/CH ₃ CH(OH)CO(O ⁻))
2,3-dihydroxypropanoic acid - glyceric acid					
CH ₂ (OH)CH(OH)CO(OH) ↔ CH ₂ (OH)CH(OH)CO(O ⁻) + H ⁺	T(120)	3.0·10 ⁻⁴		Lide, 2005	
CH(OH)(OO [•])CH(OH)CO(OH) ↔ CH(OH)(OO [•])CH(OH)CO(O ⁻) + H ⁺	T(121)	3.0·10 ⁻⁴			4 = K _a (CH ₂ (OH)CH(OH)CO(OH)/CH ₂ (OH)CH(OH)CO(O ⁻))
CH ₂ (OH)C(OH)(OO [•])CO(OH) ↔ CH ₂ (OH)C(OH)(OO [•])CO(O ⁻) + H ⁺	T(122)	3.0·10 ⁻⁴			4 = K _a (CH ₂ (OH)CH(OH)CO(OH)/CH ₂ (OH)CH(OH)CO(O ⁻))
2-hydroxy, 3-oxopropanoic acid					
CHOCH(OH)CO(OH) ↔ CHOCH(OH)CO(O ⁻) + H ⁺	T(123)	3.0·10 ⁻⁴			= K _a (CH ₂ (OH)CH(OH)CO(OH)/CH ₂ (OH)CH(OH)CO(O ⁻))
CHOCH(OH)CO(OH) + H ₂ O ↔ CH(OH)(OH)CH(OH)CO(OH)	T(124)	4.6 10 ¹		Estimated with GROMHE	
CHOCH(OH)CO(O ⁻) + H ₂ O ↔ CH(OH)(OH)CH(OH)CO(O ⁻)	T(125)	6.1		Estimated with GROMHE	
CO(OH)CH(OH)C(OH)(OH)(OO [•]) ↔ CO(O ⁻)CH(OH)C(OH)(OH)(OO [•]) + H ⁺	T(126)	3.0·10 ⁻⁴			4 = K _a (CHOCH(OH)CO(OH)/CHOCH(OH)CO(O ⁻))
CO(OH)CH(OH)CO(OO [•]) + H ₂ O ↔ CO(OH)CH(OH)C(OH)(OH)(OO [•])	T(127)	1.0 10 ⁻³			2
CO(O ⁻)CH(OH)CO(OO [•]) + H ₂ O ↔ CO(O ⁻)CH(OH)C(OH)(OH)(OO [•])	T(128)	1.0 10 ⁻³			2
3-oxopyruvic acid					
CO(OH)COCHO ↔ CO(O ⁻)COCHO + H ⁺	T(129)	4.1·10 ⁻³			= K _a (CH ₃ COCO(OH)/CH ₃ COCO(O ⁻))
CO(OH)COCHO + H ₂ O ↔ CO(OH)COCH(OH)(OH)	T(130)	8.1 10 ²		Estimated with GROMHE	
CO(OH)COCHO + H ₂ O ↔ CO(OH)C(OH)(OH)CHO	T(131)	4.4 10 ²		Estimated with GROMHE	

Species		K _a or K _h	-ΔH/R (K)	References	Notes
CO(OH)COCHO + 2 H ₂ O ↔ CO(OH)C(OH)(OH)CH(OH)(OH)	T(132)	1.3 10 ⁵		Estimated with GROMHE	
CO(O ⁻)COCHO + H ₂ O ↔ CO(O ⁻)COCH(OH)(OH)	T(133)	1.1 10 ²		Estimated with GROMHE	
CO(O ⁻)COCHO + H ₂ O ↔ CO(O ⁻)C(OH)(OH)CHO	T(134)	2.8		Estimated with GROMHE	
CO(O ⁻)COCHO + 2 H ₂ O ↔ CO(O ⁻)C(OH)(OH)CH(OH)(OH)	T(135)	1.1 10 ²		Estimated with GROMHE	
CO(OH)C(OH)(OH)C(OH)(OH)(OO [•]) ↔ CO(O ⁻)C(OH)(OH)C(OH)(OH)(OO [•]) + H ⁺	T(136)	4.1·10 ⁻³			4 = K _a (CO(OH)COCHO/CO(O ⁻)COCHO)
CO(OH)COCO(OO [•]) + H ₂ O ↔ CO(OH)COC(OH)(OH)(OO [•])	T(137)	1.0 10 ⁻³			2
CO(OH)COCO(OO [•]) + H ₂ O ↔ CO(OH)C(OH)(OH)CO(OO [•])	T(138)	4.4 10 ²			1 = K _h (CO(OH)COCHO/CO(OH)C(OH)(OH)CHO)
CO(OH)COCO(OO [•]) + 2 H ₂ O ↔ CO(OH)C(OH)(OH)C(OH)(OH)(OO [•])	T(139)	1.3 10 ⁵			1 = K _h (CO(OH)COCHO/CO(OH)C(OH)(OH)CH(OH)(OH))
CO(O ⁻)COCO(OO [•]) + H ₂ O ↔ CO(O ⁻)COC(OH)(OH)(OO [•])	T(140)	1.0 10 ⁻³			2
CO(O ⁻)COCO(OO [•]) + H ₂ O ↔ CO(O ⁻)C(OH)(OH)CO(OO [•])	T(141)	2.8			1 = K _h (CO(O ⁻)COCHO/CO(O ⁻)C(OH)(OH)CHO)
CO(O ⁻)COCO(OO [•]) + 2 H ₂ O ↔ CO(O ⁻)C(OH)(OH)C(OH)(OH)(OO [•])	T(142)	1.1 10 ²			1 = K _h (CO(O ⁻)COCHO/CO(O ⁻)C(OH)(OH)CH(OH)(OH))
3-hydroxypyruvic acid					
CH ₂ (OH)COCO(OH) ↔ CH ₂ (OH)COCO(O ⁻) + H ⁺	T(143)	4.1·10 ⁻³			= K _a (CH ₃ COCO(OH)/CH ₃ COCO(O ⁻))
CH ₂ (OH)COCO(OH) + H ₂ O ↔ CH ₂ (OH)C(OH)(OH)CO(OH)	T(144)	1.1 10 ¹		Estimated with GROMHE	
CH ₂ (OH)COCO(O ⁻) + H ₂ O ↔ CH ₂ (OH)C(OH)(OH)CO(O ⁻)	T(145)	7.0 10 ⁻²		Estimated with GROMHE	
CH(OH)(OO [•])COCO(OH) ↔ CH(OH)(OO [•])COCO(O ⁻) + H ⁺	T(146)	4.1·10 ⁻³			4 = K _a (CH ₃ COCO(OH)/CH ₃ COCO(O ⁻))
CH(OH)(OO [•])COCO(OH) + H ₂ O ↔ CH(OH)(OO [•])C(OH)(OH)CO(OH)	T(147)	1.1 10 ¹			1 = K _h (CH ₂ (OH)COCO(OH)/CH ₂ (OH)C(OH)(OH)CO(OH))
CH(OH)(OO [•])COCO(O ⁻) + H ₂ O ↔ CH(OH)(OO [•])C(OH)(OH)CO(O ⁻)	T(148)	7.0 10 ⁻²			1 = K _h (CH ₂ (OH)COCO(O ⁻)/CH ₂ (OH)C(OH)(OH)CO(O ⁻))
Pyruvic acid					
CH ₃ COCO(OH) ↔ CH ₃ COCO(O ⁻) + H ⁺	T(149)	4.1·10 ⁻³		Lide, 2005	
CH ₃ COCO(OH) + H ₂ O ↔ CH ₃ C(OH)(OH)CO(OH)	T(150)	1.5		Pocker et al., 1969	
CH ₃ COCO(O ⁻) + H ₂ O ↔ CH ₃ C(OH)(OH)CO(O ⁻)	T(151)	5.7 10 ⁻²		Pocker et al., 1969	
CO(OH)COCH ₂ (OO [•]) ↔ CO(O ⁻)COCH ₂ (OO [•]) + H ⁺	T(152)	4.1·10 ⁻³			4 = K _a (CH ₃ COCO(OH)/CH ₃ COCO(O ⁻))
CO(OH)COCH ₂ (OO [•]) + H ₂ O ↔ CO(OH)C(OH)(OH)CH ₂ (OO [•])	T(153)	1.5			1 = K _h (CH ₃ COCO(OH)/CH ₃ C(OH)(OH)CO(OH))
CO(O ⁻)COCH ₂ (OO [•]) + H ₂ O ↔ CO(O ⁻)C(OH)(OH)CH ₂ (OO [•])	T(154)	5.7 10 ⁻²			1 = K _h (CH ₃ COCO(O ⁻)/CH ₃ C(OH)(OH)CO(O ⁻))

Species		K _a or K _h	-ΔH/R (K)	References	Notes
Acroleine					
CH ₂ (OH)CH(OO•)CHO + H ₂ O ↔ CH ₂ (OH)CH(OO•)CH(OH)(OH)	T(155)	1.1 10 ¹			1 = K _h (CH ₂ (OH)CH(OH)CHO/CH ₂ (OH)CH(OH)CH(OH)(OH))
2,3 dihydroxypropanal - glyceraldehyde					
CH ₂ (OH)CH(OH)CHO + H ₂ O ↔ CH ₂ (OH)CH(OH)CH(OH)(OH)	T(156)	1.1 10 ¹		Estimated with GROMHE	
CHOCH(OH)CH(OH)(OO•) + H ₂ O ↔ CH(OH)(OH)CH(OH)CH(OH)(OO•)	T(157)	1.1 10 ¹			1 = K _h (CH ₂ (OH)CH(OH)CHO/CH ₂ (OH)CH(OH)CH(OH)(OH))
CH ₂ (OH)CH(OH)CO(OO•) + H ₂ O ↔ CH ₂ (OH)CH(OH)C(OH)(OH)(OO•)	T(158)	1.0 10 ⁻³			2
Hydroxypropanedial					
CHOCH(OH)CHO + H ₂ O ↔ CHOCH(OH)CH(OH)(OH)	T(159)	9.8 10 ¹		Estimated with GROMHE	
CHOCH(OH)CHO + 2 H ₂ O ↔ CH(OH)(OH)CH(OH)CH(OH)(OH)	T(160)	2.0 10 ³		Estimated with GROMHE	
CHOCH(OH)CO(OO•) + H ₂ O ↔ CHOCH(OH)C(OH)(OH)(OO•)	T(161)	1.0 10 ⁻³			2
CHOCH(OH)CO(OO•) + 2 H ₂ O ↔ CH(OH)(OH)CH(OH)C(OH)(OH)(OO•)	T(162)	2.0 10 ³			1 = K _h (CHOCH(OH)CHO/CH(OH)(OH)CH(OH)CH(OH)(OH))

1 - For peroxy radicals, we assumed that the hydration constant is similar to the parent species.

2 - For acyl peroxy radicals, we assumed that hydration is not favored on the –CO(OO•) moiety. This is based on the similarity between the CO moiety in this function and the CO moiety in carboxylic (-CO(OH)) or percarboxylic acid (-CO(OOH)) organic functions which is not readily hydrated. Therefore we apply an arbitrarily low value (K_h = 10⁻³) to the hydration constant of these species.

3 - The parent species is symmetrical whereas the corresponding R(OO•) is asymmetrical. We assume that hydration occurs mainly on the CO(OO•) function.

4 - For peroxy radicals, we assumed that the acidity constant is similar to the parent species.

5 - For methylglyoxal, Wasa and Musha (1970) measured the total hydration constant. Each fractional hydration constant is calculated from the measured total hydration constant weighted by the relative contribution of each hydrate estimated with GROMHE.

References:

Wasa, T., and Musha, S.: Polarographic behavior of glyoxal and its related compounds, Bull. Univ. Osaka Pref. A, Eng. Nat. Sci., 19, 169-180, 1970.

Henry's law constants

Species		H (298K) (M atm ⁻¹)	-ΔH/R (K)	References	Notes
C3 compounds					
1-propanol CH ₃ CH ₂ CH ₂ (OH)	T(42)	1.3 10 ²	7470	Sander, 2015	
3-hydroxypropionaldehyde CH ₂ (OH)CH ₂ CHO	T(43)	2.3 10 ³	6014	Estimated	1 - 2 - 3
Propane-1,2-diol CH ₃ CH(OH)CH ₂ (OH)	T(44)	3.0 10 ⁶	6014	Estimated	3 - 4
2-hydroxypropionaldehyde CH ₃ CH(OH)CHO	T(45)	2.2 10 ³	6014	Estimated	1 - 2 - 3
Propanedial CHOCH ₂ CHO	T(46)	3.3 10 ⁴	6014	Estimated	1 - 2 - 3
3-hydroxypropionic acid CH ₂ (OH)CH ₂ CO(OH)	T(47)	2.0 10 ⁵	6014	Estimated	1 - 3
2-propanol CH ₃ CH(OH)CH ₃	T(48)	1.3 10 ²	7470	Sander, 2015	
Propionaldehyde CH ₃ CH ₂ CHO	T(49)	1.0 10 ¹	4330	Sander, 2015	2
Methylglyoxal CH ₃ COCHO	T(50)	3.7 10 ³	7540	Betterton and Hoffmann, 1988	2
2-oxo, 3-hydroxypropanal CH ₂ (OH)COCHO	T(51)	1.3 10 ⁷	6014	Estimated	1 - 2 - 3
Oxopropanedial CHOCOCHO	T(52)	9.8 10 ¹⁰	6014	Estimated	1 - 2 - 3
Acetone CH ₃ COCH ₃	T(53)	2.8 10 ¹	5050	Sander, 2015	2
Hydroxyacetone CH ₃ COCH ₂ (OH)	T(54)	1.6 10 ³	6014	Estimated	1 - 2 - 3
Dihydroxyacetone CH ₂ (OH)COCH ₂ (OH)	T(55)	1.1 10 ⁷	6014	Estimated	1 - 2 - 3
Propionic acid CH ₃ CH ₂ CO(OH)	T(56)	5.7 10 ³		Khan et al., 1995	
3-oxopropionic acid CO(OH)CH ₂ CHO	T(57)	9.0 10 ⁵	6014	Estimated	1 - 2 - 3

Species		H (298K) (M atm ⁻¹)	-ΔH/R (K)	References	Notes
Malonic acid CO(OH)CH ₂ CO(OH)	T(58)	3.9 10 ¹⁰	11000	Compernelle and Müller, 2014	
Tartronic acid CO(OH)CH(OH)CO(OH)	T(59)	1.0 10 ⁷	6014	Estimated	1 - 3
Mesoxalic acid CO(OH)COCO(OH)	T(60)	2.7 10 ¹⁰	6014	Estimated	1 - 2 - 3
Lactic acid CH ₃ CH(OH)CO(OH)	T(61)	9.6 10 ⁴	6014	Estimated	1 - 3
2,3-dihydroxypropanoic acid CH ₂ (OH)CH(OH)CO(OH)	T(62)	8.1 10 ⁷	6014	Estimated	1 - 3
2-hydroxy, 3-oxopropanoic acid CHOCH(OH)CO(OH)	T(63)	1.6 10 ⁶	6014	Estimated	1 - 2 - 3
3-oxopyruvic acid CO(OH)COCHO	T(64)	1.2 10 ¹¹	6014	Estimated	1 - 2 - 3
3-hydroxypyruvic acid CH ₂ (OH)COCO(OH)	T(65)	4.3 10 ⁷	6014	Estimated	1 - 2 - 3
Pyruvic acid CH ₃ COCO(OH)	T(66)	3.1 10 ⁵	5090	Sander, 2015	
Acrolein CH ₂ =CHCHO	T(67)	7.4	5100	Snider and Dawson, 1985	
2,3-dihydroxypropanal - glyceraldehyde CH ₂ (OH)CH(OH)CHO	T(68)	3.2 10 ⁵	6014	Estimated	1 - 2 - 3
Hydroxypropanedial CHOCH(OH)CHO	T(69)	2.3 10 ⁵	6014	Estimated	1 - 2 - 3

1 - Estimated by the SAR GROMHE (Raventos-Duran et al., 2010).

2 - Effective Henry's law constant.

3 - When unavailable, the temperature dependence (enthalpy of dissolution) is set at 50 kJ mol⁻¹; -ΔH/R = 6014 K.

4 - Saxena and Hildemann (1996) estimated the Henry's law constant between 10⁵ and 6.0 10⁶ M atm⁻¹. We suppose the Henry's law constant equal to 3.0 10⁶ M atm⁻¹.

References:

Raventos-Duran, T., Camredon, M., Valorso, R., Mouchel-Vallon, C., and Aumont, B.: Structure-activity relationships to estimate the effective Henry's law constants of organics of atmospheric interest, Atmos. Chem. Phys., 10, 7643-7654, 2010.

Saxena, P., and Hildemann, L. M.: Water-soluble organics in atmospheric particles: A critical review of the literature and application of thermodynamics to identify candidate compounds, J. Atmos. Chem., 24, 57-109, 1996.

Accommodation coefficients

Species		α (298K)	$-\Delta H$ (J/mol)	$-\Delta S$ (J/mol/K)	References	Notes
C3 compounds		1				
1-propanol CH ₃ CH ₂ CH ₂ (OH)	T(42)	6.0 10 ⁻³	3.8 10 ⁴	1.7 10 ²	Davidovits et al., 2011	
3-hydroxypropionaldehyde CH ₂ (OH)CH ₂ CHO	T(43)	5.0 10 ⁻²			Estimated	2
Propane-1,2-diol CH ₃ CH(OH)CH ₂ (OH)	T(44)	5.0 10 ⁻²			Estimated	2
2-hydroxypropionaldehyde CH ₃ CH(OH)CHO	T(45)	5.0 10 ⁻²			Estimated	2
Propanedial CHOCH ₂ CHO	T(46)	5.0 10 ⁻²			Estimated	2
3-hydroxypropionic acid CH ₂ (OH)CH ₂ CO(OH)	T(47)	5.0 10 ⁻²			Estimated	2
2-propanol CH ₃ CH(OH)CH ₃	T(48)	7.0 10 ⁻³	4.1 10 ⁴	1.8 10 ²	Davidovits et al., 2011	
Propionaldehyde CH ₃ CH ₂ CHO	T(49)	5.0 10 ⁻²			Estimated	2
Methylglyoxal CH ₃ COCHO	T(50)	1.0 10 ⁻⁴			Sander, 2015	
2-oxo, 3-hydroxypropanal CH ₂ (OH)COCHO	T(51)	5.0 10 ⁻²			Estimated	2
Oxopropanedial CHOCOCHO	T(52)	5.0 10 ⁻²			Estimated	2
Acetone CH ₃ COCH ₃	T(53)	4.0 10 ⁻³	5.3 10 ⁴	2.2 10 ²	Davidovits et al., 2011	
Hydroxyacetone CH ₃ COCH ₂ (OH)	T(54)	5.0 10 ⁻²			Estimated	2
Dihydroxyacetone CH ₂ (OH)COCH ₂ (OH)	T(55)	5.0 10 ⁻²			Estimated	2
Propionic acid CH ₃ CH ₂ CO(OH)	T(56)	5.0 10 ⁻²			Estimated	2
3-oxopropionic acid CO(OH)CH ₂ CHO	T(57)	5.0 10 ⁻²			Estimated	2
Malonic acid CO(OH)CH ₂ CO(OH)	T(58)	5.0 10 ⁻²			Estimated	2

Species		α (298K)	$-\Delta H$ (J/mol)	$-\Delta S$ (J/mol/K)	References	Notes
Tartronic acid CO(OH)CH(OH)CO(OH)	T(59)	5.0 10 ⁻²			Estimated	2
Mesoxalic acid CO(OH)COCO(OH)	T(60)	5.0 10 ⁻²			Estimated	2
Lactic acid CH ₃ CH(OH)CO(OH)	T(61)	5.0 10 ⁻²			Estimated	2
2,3-dihydroxypropanoic acid CH ₂ (OH)CH(OH)CO(OH)	T(62)	5.0 10 ⁻²			Estimated	2
2-hydroxy, 3-oxopropanoic acid CHOCH(OH)CO(OH)	T(63)	5.0 10 ⁻²			Estimated	2
3-oxopyruvic acid CO(OH)COCHO	T(64)	5.0 10 ⁻²			Estimated	2
3-hydroxypyruvic acid CH ₂ (OH)COCO(OH)	T(65)	5.0 10 ⁻²			Estimated	2
Pyruvic acid CH ₃ COCO(OH)	T(66)	5.0 10 ⁻²			Estimated	2
Acrolein CH ₂ =CHCHO	T(67)	5.0 10 ⁻²			Estimated	2
2,3-dihydroxypropanal - glyceraldehyde CH ₂ (OH)CH(OH)CHO	T(68)	5.0 10 ⁻²			Estimated	2
Hydroxypropanedial CHOCH(OH)CHO	T(69)	5.0 10 ⁻²			Estimated	2

1 - α can be calculated with ΔH and ΔS ; this allows considering the temperature dependency of α following Jayne et al. (1997): $\frac{\alpha}{1-\alpha} = \exp\left(\frac{-\Delta G}{RT}\right)$; $\Delta G = \Delta H - T\Delta S$

2 - Estimated equal 5.0 10⁻² following Lelieveld and Crutzen (1991) and Davidovits et al. (2011).

References:

Davidovits, P., Kolb, C. E., Williams, L. R., Jayne, J. T., and Worsnop, D. R.: Update 1 of: Mass accommodation and chemical reactions at gas-liquid interfaces, Chem. Rev., 111, 2011.
Lelieveld, J., and Crutzen, P. J.: The role of clouds in tropospheric photochemistry, J. Atmos. Chem., 12, 229-267, 1991.

REFERENCES:

- Adams, G. E., and Willson, R. L.: Pulse radiolysis studies on the oxidation of organic radicals in aqueous solution, *Trans.Faraday Soc.*, 65, 1969.
- Albalat, R., Claret, J., Gómez, E., Müller, C., and Sarret, M.: Electrocatalytic oxidation of mesoxalic acid on a polycrystalline platinum electrode in acid medium, *Electrochimica Acta*, 34, 611-618, 1989.
- Betterton, E. A., and Hoffmann, M. R.: Henry's law constants of some environmentally important aldehydes, *Environ. Sci. Tech.*, 22, 1415-1418, 1988.
- Compernelle, S., and Müller, J. F.: Henry's law constants of diacids and hydroxy polyacids: recommended values, *Atmos. Chem. Phys.*, 14, 2699-2712, 2014.
- Davidovits, P., Kolb, C. E., Williams, L. R., Jayne, J. T., and Worsnop, D. R.: Update 1 of: Mass accommodation and chemical reactions at gas-liquid interfaces, *Chem. Rev.*, 111, 2011.
- De Semainville, P., Hoffmann, D., George, C., and Herrmann, H.: Study of nitrate radical (NO_3) reactions with carbonyls and acids in aqueous solution as a function of temperature, *Phys. Chem. Chem. Phys.*, 9, 958-968, 2007.
- De Semainville, P. G., D Anna, B., and George, C.: Aqueous phase reactivity of nitrate radicals (NO_3) toward dicarboxylic acids, *Z. Phys. Chem.*, 224, 1247-1260, 2010.
- Ervens, B., Gligorovski, S., and Herrmann, H.: Temperature-dependent rate constants for hydroxyl radical reactions with organic compounds in aqueous solutions, *Phys. Chem. Chem. Phys.*, 5, 2003.
- Gligorovski, S., Rousse, D., George, C. H., and Herrmann, H.: Rate constants for the OH reactions with oxygenated organic compounds in aqueous solution, *Inter. J. Chem. Kin.*, 41, 309-326, 2009.
- Griffith, E. C., Carpenter, B. K., Shoemaker, R. K., and Vaida, V.: Photochemistry of aqueous pyruvic acid, *PNAS*, 110, 11714-11719, 2013.
- Herrmann, H., Exner, M., and Zellner, R.: Reactivity trends in reactions of the nitrate radical (NO_3) with inorganic and organic cloudwater constituents, *Geochim.Cosmochim. Acta*, 58, 3239-3244, 1994.
- Herrmann, H., and Zellner, R.: Reaction of NO_3 radicals in aqueous solution, in: *N-Centered Radicals*, edited by: Alfassi, Z., Wiley, London, 291-343, 1998.
- Hislop, K. A., and Bolton, J. R.: The photochemical generation of hydroxyl radicals in the UV-vis/Ferrioxalate/ H_2O_2 system, *Environ. Sci.Tech.*, 33, 3119-3126, 1999.
- Hoffmann, D., Weigert, B., Barzaghi, P., and Herrmann, H.: Reactivity of poly-alcohols towards OH, NO_3 and SO_4^- in aqueous solution, *Phys. Chem. Chem. Phys.*, 11, 9351-9363, 2009.
- Ilan, Y., Rabani, J., and Henglein, A.: Pulse radiolytic investigations of peroxy radicals produced from 2-propanol and methanol, *J. Phys. Chem.*, 80, 1558-1562, 1976.
- Khan, I., Brimblecombe, P., and Clegg, S. L.: Solubilities of pyruvic acid and the lower (C1-C6) carboxylic acids. Experimental determination of equilibrium vapour pressures above pure aqueous and salt solutions, *J Atmos Chem*, 22, 285-302, 1995.
- Le Henaff, P.: Methodes d'étude et propriétés des hydrates hémiacétals et hémithioacétals dérivés des aldéhydes et cétones, *P. Bull. Soc. Chim. Fr.*, 11, 4687-4698, 1968.
- Lide, D. R.: Dissociation constants of organic acids and bases, in: *CRC Handbook of Chemistry and Physics*, Internet Version 2005, edited by: Lide, D. R., CRC Press, Boca Raton, FL, 2005.
- Lilie, J., and Henglein, A.: Pulsradiolytische Untersuchung der Oxidation von ungesättigten Carbonylverbindungen in wässriger Lösung: Hydratisierte Enole als Zwischenprodukte, *Berichte der Bunsengesellschaft für physikalische Chemie*, 74, 388-393, 1970.
- Martin, L. R., Mezyk, S. P., and Mincher, B. J.: Determination of Arrhenius and thermodynamic parameters for the aqueous reaction of the hydroxyl radical with lactic acid, *J. Phys. Chem. A*, 113, 141-145, 2008.
- Monod, A., Poulain, L., Grubert, S., Voisin, D., and Wortham, H.: Kinetics of OH-initiated oxidation of oxygenated organic compounds in the aqueous phase: new rate constants, structure-activity relationships and atmospheric implications, *Atmos. Environ.*, 39, 7667-7688, 2005.
- Padmaja, S., Neta, P., and Huie, R. E.: Rate constant for some reactions of inorganic radicals with inorganic ions. Temperature dependence and solvent dependence, *J. Phys. Chem.*, 25, 445-455, 1993.
- Pocker, Y., Meany, J. E., Nist, B. J., and Zadorojny, C.: Reversible hydration of pyruvic acid. I. Equilibrium studies, *J. Phys. Chem.*, 73, 2879-2882, 1969.
- Pressman, D., and Lucas, H. J.: Hydration of unsaturated compounds. XI. Acrolein and acrylic acid, *J. Am. Chem. Soc.*, 64, 1953-1957, 1942.
- Reed Harris, A. E., Ervens, B., Shoemaker, R. K., Kroll, J. A., Rapf, R. J., Griffith, E. C., Monod, A., and Vaida, V.: Photochemical kinetics of pyruvic acid in aqueous solution, *J. Phys. Chem. A*, 118, 8505-8516, 2014.
- Rousse, D., and George, C.: A novel long path photolysis cell-application to the reactivity of selected organic compounds toward the nitrate radical (NO_3), *Phys. Chem. Chem. Phys.*, 6, 3408-3414, 2004.
- Sander, R.: Compilation of Henry's law constants (version 4.0) for water as solvent, *Atmos. Chem. Phys.*, 15, 4399-4981, 10.5194/acp-15-4399-2015, 2015.
- Schaefer, T.: Kinetische und mechanistische Untersuchungen der radikalischen Oxidation organischer kurzketziger Carbonylverbindungen in wässriger Lösung (Kinetic and mechanistic investigations of the radical oxidation of organic short-chain carbonyl compounds in the aqueous phase), Faculty of Chemistry and Mineralogy of the University of Leipzig, Leipzig, 174 pp., 2012.
- Schaefer, T., Schindelka, J., Hoffmann, D., and Herrmann, H.: Laboratory kinetic and mechanistic studies on the OH-initiated oxidation of acetone in aqueous solution, *J. Phys. Chem. A*, 116, 6317-6326, 2012.
- Schaefer, T., van Pinxteren, D., and Herrmann, H.: Multiphase chemistry of glyoxal: revised kinetics of the alkyl radical reaction with molecular oxygen and the reaction of glyoxal with OH, NO_3 , and SO_4^- in aqueous solution, *Environ. Sci. Tech.*, 49, 343-350, 10.1021/es505860s, 2014.
- Schöne, L., and Herrmann, H.: Kinetic measurements of the reactivity of hydrogen peroxide and ozone towards small atmospherically relevant aldehydes, ketones and organic acids in aqueous solutions, *Atmos. Chem. Phys.*, 14, 4503-4514, 2014.
- Schuchmann, M. N., Schuchmann, H.-P., and von Sonntag, C.: Oxidation of hydroxymalonic acid by OH radicals in the presence and in the absence of molecular oxygen. A pulse-radiolysis and product study, *J. Phys. Chem.*, 99, 9122-9129, 1995.
- Snider, J. R., and Dawson, G. A.: Tropospheric light alcohols, carbonyls, and acetonitrile: Concentrations in the southwestern united states and Henry's law data, *J. Geophys. Res.*, 90, 3797-3805, 1985.
- Socrates, G.: Hydration study of acetaldehyde and propionaldehyde, *J. Org. Chem.*, 34, 2958-2961, 1969.

Wang, W.-F., Schuchmann, M. N., Schuchmann, H.-P., and von Sonntag, C.: The Importance of mesomerism in the termination of α -carboxymethyl radicals from aqueous malonic and acetic acids, Chemistry – A European Journal, 7, 791-795, 2001.

Wasa, T., and Musha, S.: Polarographic behavior of glyoxal and its related compounds, Bull. Univ. Osaka Pref. A, Eng. Nat. Sci., 19, 169-180, 1970.

Zegota, H., Schuchmann, M. N., Schulz, D., and Sonntag, C. V.: Acetonylperoxyl radicals, $\text{CH}_3\text{COCH}_2\text{O}_2\cdot$: a study on the γ -radiolysis and pulse radiolysis of acetone in oxygenated aqueous solutions, Z. Naturforsch., 41b, 1015-1022, 1986a.