

Table S1. Primer sequences used in qRT-PCR.

Gene	Forward primer	Reverse primer
<i>Actin 5C</i>	CAGAGCAAGCGTGGTATCCT	CTCATTGTAGAAGGTGTGGTGC
<i>Cyp4e3</i>	CGGTCCTTTGAGGTCCTTCC	CGCGGATAGAATGGGATCGT
<i>E(spl)m4</i> <i>-BFM</i>	GCCCTCAAGGCCAACTCTTT	CCTCAATTTTCGCAGGACTGG
<i>Ets21C</i>	CGGATCCGGAAACGAGACC	CCGGAATCCGAGTCTGTTGA
<i>Hsp70Ba</i>	CCAAGATGCATCAGCAGGGA	GTCCAGAGTAGCCGCCAAAT
<i>IM3</i>	TCACTCGCCTTCGTTTTGGG	TTAGGCCCTCACATTGCAGAC
<i>Lip3</i>	CCCCCAAACGGGTGAATCT	CATTGCCCATCCACACATCG
<i>LManV</i>	GCCATCACTGGAACCGAGAA	GCAGGCAACTCTCGAACTCT
<i>Lst</i>	TGCGATCCAAGCCGTACTTT	GGACGACTCAGCTTCTGGAC
<i>Pcp</i>	TCGGTATGCCTACGAGACCT	TAGCCAAACTCATCGGCCAC
<i>Prat2</i>	GTGCTGCCAGAAGACAAAGG	ATTCGCAGGTCATTCCGGTG
<i>Sit</i>	AGGTGGTCGGTGTGCTTTAC	GACAGAGCCTTCTCCTTGGC
<i>Socs36E</i>	GGCGTGGTGACAAATCGAAC	CCATGGTTGTCGTGCAGTTG
<i>TotC</i>	CTACTATGCCTTGCCCTGCT	AGCAGCTCTTGGGTTCGATT