

**Carvacrol preserves antioxidant status and attenuates kidney fibrosis via
modulation of the TGF- β 1/Smad signaling and inflammation**

Chetan Ram¹, Shobhit Gairola¹, Abu Mohammad Syed¹, Shobhit Verma², Madhav
Nilakanth Mugale², Bidya Dhar Sahu^{1*}

¹ Department of Pharmacology & Toxicology, National Institute of Pharmaceutical Education and Research (NIPER)-Guwahati, Changsari, PIN-781101, Assam, India.

² Toxicology & Experimental Medicine, CSIR- Central Drug Research Institute (CDRI), Lucknow-226 031, India.

*Corresponding Author:

Dr. Bidya Dhar Sahu, Ph.D., Assistant Professor, Department of Pharmacology & Toxicology, National Institute of Pharmaceutical Education and Research (NIPER)-Guwahati, Changsari-781101, Assam, India.

Email ID: bidyasha@gmail.com; bidya.sahu@niperguwahati.ac.in

Raw western blot images

Fig. 3A: Uncropped raw western blot images of TGF- β , p-Smad2/3, Smad2/3, and β -actin in kidneys.




Fig. 3A: Uncropped raw western blot images of Fibronectin, collagen III, collagen I, and β -actin in kidneys.




Fig. 5A: Uncropped raw western blot images of α -SMA, vimentin, E-cadherin, and β -actin in kidneys.




Fig. 5F: Uncropped raw western blot images of E-cadherin, vimentin, and β -actin in NRK 52E cells.




Fig. 6D: Uncropped raw western blot images of Nrf2, NQO1, and β -actin in kidneys.




Fig. 7F: Uncropped raw western blot images of NF-kB (p65), phospho-NF-kB (p65), and β -actin in kidneys.




Fig. 7H: Uncropped raw western blot images of COX-2 and β -actin in kidneys.

