

Table S1 Effects of AKP on liver Index in mice

Group	Body weight	Liver weight	Liver weight/body weight (%)
Control	32.48±1.77	1.29±0.16	3.97±0.33
Model	33.44±3.34	1.98±0.13 ^{##}	5.68±0.15 ^{##}
Silybin	30.02±0.49	1.51±0.08 [*]	5.00±0.21 ^{**}
AKP	30.05±2.70	1.52±0.13 [*]	5.14±0.19 ^{**}

Note: # $P < 0.05$, and ## $P < 0.01$ compared with the Control group, and * $P < 0.05$, and ** $P < 0.01$

compared with the Model group. Data are given as the mean ± SEM (n = 8).

Table S2 Primers used for the quantitative RT-PCR analysis

Gene	Forward primer	Reverse primer
CYP2E1	5'-CCGACCTGTTCTTTGCAGGA-3'	5'-GGCTTGGCCCAATAACCCTG-3'
Keap1	5'-CGCTGTGCTTAGTCACCGT-3'	5'-GCATGGGGTTCCTGATGACA-3'
Nrf2	5'-CACATCCAGACAGACACCAGT-3'	5'-CTTGTTTGGGAATGTGGGCA-3'
HO-1	5'-ACAGAAGAGGCTAAGACCGC-3'	5'-GACGAAGTGACGCCATCTGT-3'
NQO1	5'-ACGACAACGGTCCTTTCCAG-3'	5'-ATGACGTTTCATGTCCCCGTG-3'
β-actin	5'-GGCTGTATTCCCCTCCATCG-3'	5'-GTTGGGGTTTGCCTCTCCC-3'