

RSC Med Chem

Article

Inhibition of MARK4 by serotonin is an attractive therapeutic approach to combat Alzheimer's disease and neuroinflammation

Anas Shamsi^{1,2*}, Debarati DasGupta³, Fahad A. Alhumaydhi⁴, Mohd Shahnawaz Khan⁵, Suliman A. Alsagaby⁶, Waleed Al Abdulmonem⁷, Md. Imtaiyaz Hassan^{1*}, Dharmendra Kumar Yadav⁸

¹ Centre for Interdisciplinary Research in Basic Sciences, Jamia Millia Islamia, Jamia Nagar, New Delhi 110025. E-mail: anas.shamsi18@gmail.com (AS); mihassan@jmi.ac.in (MIH)

² Centre of Medical and Bio-Allied Health Sciences Research, Ajman University, United Arab Emirates.

³ College of Pharmacy, University of Michigan, Ann Arbor, United States of America. Email: debarati@med.umich.edu (DDG)

⁴ Department of Medical Laboratories, College of Applied Medical Sciences, Qassim University, Buraydah, Saudi Arabia. Email: f.alhumaydhi@qu.edu.sa (FAA)

⁵ Department of Biochemistry, College of Sciences, King Saud University, Riyadh 11451, Saudi Arabia. Email: moskhan@ksu.edu.sa (MSK)

⁶ Department of Medical Laboratory Sciences, College of Applied Medical Sciences, Majmaah University, Majmaah 11932, Saudi Arabia. Email: s.alsagaby@mu.edu.sa (SAA)

⁷ Department of Pathology, College of Medicine, Qassim University, Buraydah, Saudi Arabia. Email: waleedmonem@qumed.edu.sa (WAA)

⁸ College of Pharmacy, Gachon University of Medicine and Science, Hambakmoeiro, Yeonsu-gu, Incheon City, 21924, South Korea. Email: dharmendra30oct@gmail.com (DKY)

*Correspondence: anas.shamsi18@gmail.com (A.S.); mihassan@jmi.ac.in (M.I.H.)




Figure S1. Structure of Serotonin after washing in MOE at pH 7

Table S1. Critical descriptor of the ligand.

Mol weight	H bond donors	H bond acceptors	No of single rotatable bonds	No of aromatic rings	SlogP	Flexibility index	Lipinski Drug Like Test
177.2	2	1	2	2	0.66	1.54	1