

Immobilization of Zn₄ complex on functionalized Layered HUS-7: Synthesis, Structural Investigation and Catalytic Activity

Divya Jadav¹, Pooja Shukla¹, Rajib Bandyopadhyay², Plaban Jyoti Sarma³, Ramesh Chandra Deka³, Rajesh Kumar⁴, Sourav Das^{1*}, Nao Tsunoji^{4*}, Mahuya Bandyopadhyay^{1*}

¹Department of Basic Sciences, Institute of Infrastructure, Technology, Research and Management, IITRAM, Ahmedabad, Gujarat, India

²Department of Chemistry, School of Technology, Pandit Deendayal Energy University, Gandhinagar, Gujarat, India

³Department of Chemical Sciences, Tezpur University, Tezpur, Assam, India

⁴Graduate School of Advanced Science and Engineering, Hiroshima University, Higashi-Hiroshima 739-8527, Japan




Figure S1. Figure S1. N₂ adsorption/desorption isotherm with pore size distribution curve of HUS-7-SO₃H and HUS-7-SO₃-Zn₄ samples.




Figure S2. TGA profile of HUS-7-SO₃-Zn₄ and pure Zn₄ complex.




Figure S3. SEM image of spent HUS-7-SO₃-Zn₄ catalyst in (A) morpholine system and (B) piperidine system.