

Supporting Information

New UV-light initiated intramolecular Se-N bond formation

Agata J. Pacuła-Miszewska^a, Magdalena Obieziurska-Fabisiak^a, Anna Laskowska^a,
Halina Kaczmarek^b and Jacek Ścianowski^a

Table of Content

I. Synthesis of diselenides 21a and 22a	2
II. Synthesis of benzoselenazol-2(H)-ones 21 and 22	3
III. ¹ H, ¹³ C and ⁷⁷ Se NMR spectra for compounds 21a , 22a , 21 and 22	4
IV. ¹ H NMR spectra for compounds 10 , 11 , 12a , 12 , 13a , 13 , 14a , 14 , 15a , 15 , 16a , 16 , 17a , 17 , 18a , 18 , 19a , 19 , 20a , 20 ,	10
V. References.....	20

II. Synthesis of diselenides **21a** and **22a**

To a solution of benzoselenazol-3(2H)-one (1.0 mmol; obtained by our previously published procedure ([Method A](#)): A.J Pacuła, K. B. Kaczor, A. Wojtowicz, J. Antosiewicz, A. Janecka, A. Długosz, T. Janecki and J. Scianowski, *Bioorg. Med. Chem.*, 2017, **25**, 126–131) in methanol (10 ml) cooled to 0°C, sodium borohydride (1.0 mmol) was added and the mixture was stirred for 1h. Water (15ml) was added and the mixture was oxidized with air for 1h. Formed precipitate was filtered and dried in air.

2,2'-Diselenobis((4-(Trifluoromethyl)phenyl)benzamide) 21a

Yield: 70%, mp 228-230°C;

¹H NMR (700 MHz, DMSO) δ = 7.42 (dt, J =7.0, 0.7 Hz, 1H_{ar}), 7.47 (dt, J =7.0, 1.4 Hz, 1H_{ar}), 7.75 (d, J =8.4, 2H_{ar}), 7.78 (dd, J =7.7, 0.7 Hz, 1H_{ar}), 7.97-7.99 (m, 3H_{ar}), 10.86 (s, NH) ppm; ¹³C NMR (100.61 Hz, DMSO) δ = 120.24, 120.89, 123.47, 124.40, 124.72, 126.17, 126.48 (q), 127.03, 129.40, 130.82, 132.13, 132.63, 132.88, 133.85, 139.08, 142.80, 167.26 (C=O) ppm; ⁷⁷Se (133.55 MHz, DMSO), δ = 446.68 ppm; IR: 3314, 1649, 1615, 1600, 1563, 1523, 1517, 1461, 1429, 1409, 1322, 1269, 1256, 1186, 1157, 1114, 1067, 1047, 1017 cm⁻¹; Elemental Anal. Calcd for C₂₈H₁₈F₆N₂O₂Se₂ (687.96): C, 49.00; H, 2.64, Found: C, 48.89; H, 2.59. Elemental Anal. Calcd for C₂₈H₁₈F₆N₂O₂Se₂ (687.96): C, 49.00; H, 2.64, Found: C, 49.23; H, 2.71.

2,2'-Diselenobis((2-(Trifluoromethyl)phenyl)benzamide) 22a

Yield: 74%, mp 277-279°C;

¹H NMR (700 MHz, DMSO) δ = 7.45 (dt, J =7.7, 1.4 Hz, 1H_{ar}), 7.49 (dt, J =8.4, 1.4 Hz, 1H_{ar}), 7.59-7.61 (m, 2H_{ar}), 7.78-7.81 (m, 2H_{ar}), 7.85 (d, J =7.7, 1H_{ar}), 8.00 (d, J =7.0, 1H_{ar}), 10.51 (s, NH) ppm; ¹³C NMR (100.61 Hz, DMSO) δ = 122.71, 125.43, 126.95, 127.13 (q), 128.33, 129.02, 130.66, 131.81, 132.75, 133.01, 133.80, 135.72, 167.89 (C=O) ppm; ⁷⁷Se (133.67MHz, DMSO), δ = 446.29 ppm; IR: 3280, 1643, 1610, 1587, 1525, 1487, 1454, 1321, 1306, 1288, 1267, 1256, 1173, 1141, 1110, 1060, 1036 cm⁻¹; Elemental Anal. Calcd for C₂₈H₁₈F₆N₂O₂Se₂ (687.96): C, 49.00; H, 2.64, Found: C, 48.89; H, 2.59

II. Synthesis of benzoselenazol-2(H)-ones **21** and **22**

Method A: To a solution of amine (2.0 mmol) and triethylamine (4.0 mmol) in dichloromethane (10 ml) 2-(chloroseleno)benzoyl chloride (2.0 mmol) was added. The mixture was stirred for 24h at room temperature, poured on water and extracted with DCM. The combined organic layers were dried over anhydrous magnesium sulfate and evaporated. The crude product was purified by column chromatography (silica gel, dichlorometane).

Method B: A solution of the starting diselenide **21a/22a** in acetonitrile (0.01M) was placed in a 5ml quartz cuvette and irradiated for 1h by 250 nm wavelength UV lamp. The solution was poured into a 10ml flask and evaporated. The final product was isolated by column chromatography (DCM, neutral aluminium oxide).

N*-4-(Trifluoromethyl)phenyl-1,2-benzisoselenazol-3(2*H*)-one **21*

Yield: 46% (Method A); Yield: 90% (Method B); mp 228-230°C;

^1H NMR (700 MHz, DMSO) δ = 7.48 (dt, $J=7.7$, 0.7 Hz, 1 H_{ar}), 7.69 (dt, $J=8.4$, 1.4 Hz, 1 H_{ar}), 7.78 (d, $J=8.4$, 2 H_{ar}), 7.91-7.93 (m, 3 H_{ar}), 8.09 (d, $J=7.7$, 1 H_{ar}) ppm; ^{13}C NMR (176.10 MHz, DMSO) δ = 123.85, 124.74, 125.39, 125.78, 125.96, 126.35, 126.80 (q), 128.61, 128.87, 133.19, 139.16, 144.17, 165.95 (C=O) ppm; ^{77}Se (133.55 MHz, DMSO), δ = 975.57 ppm; IR: 2922, 1624, 1597, 1568, 1513, 1445, 1419, 1320, 1305, 1268, 1192, 1174, 1107, 1070, 1014 cm^{-1} ; Elemental Anal. Calcd for $\text{C}_{14}\text{H}_8\text{F}_3\text{NOSe}$ (342.97): C, 49.14; H, 2.36, Found: C, 48.95; H, 2.29.



N*-2-(Trifluoromethyl)phenyl-1,2-benzisoselenazol-3(2*H*)-one **22*


Yield: 41% (Method A); Yield: 89% (Method B); (mp 199-201°C;

^1H NMR (700 MHz, DMSO) δ = 7.47 (dt, $J=7.7$, 0.7 Hz, 1 H_{ar}), 7.49 (d, $J=7.7$, 1 H_{ar}), 7.65 (t, $J=7.7$, 1 H_{ar}), 7.68 (dt, $J=7.7$, 0.7 Hz, 1 H_{ar}), 7.77 (t, $J=7.0$, 1 H_{ar}), 7.85 (d, $J=8.4$, 2 H_{ar}), 8.05 (d, $J=8.4$, 1 H_{ar}) ppm; ^{13}C NMR (176.10 MHz, DMSO) δ = 123.01, 124.57, 126.31, 126.69, 126.83, 127.64 (q), 128.39, 128.83, 129.00, 129.52, 132.75, 132.96, 134.15, 137.49, 140.90, 167.36 (C=O) ppm; ^{77}Se (133.6 MHz, DMSO), δ = 975.63 ppm; IR: 3064, 1590, 1561, 1496, 1454, 1443, 1347, 1317, 1272, 1260, 1221, 1175, 1164, 1128, 1117, 1109, 1058, 1034, 1025 cm^{-1} ; Elemental Anal. Calcd for $\text{C}_{14}\text{H}_8\text{F}_3\text{NOSe}$ (342.97): C, 49.14; H, 2.36, Found: C, 49.33; H, 2.39.


III. ^1H , ^{13}C and ^{77}Se NMR spectra for compounds **21a**, **22a**, **21** and **22**



N*-4-(Trifluoromethyl)phenyl-1,2-benziselenazol-3(2*H*)-one **21*








***N*-(2-(Trifluoromethyl)phenyl)-1,2-benziselenazol-3(2H)-one 22**








2,2'-Diselenobis((4-(Trifluoromethyl)phenyl)benzamide) 21a






2,2'-Diselenobis((2-(Trifluoromethyl)phenyl)benzamide) 22a






IV. ¹H NMR spectra for compounds **10-20** and **12a-20a**.


2,2'-Diselenobis(*N*-butylbenzamide) 10¹




***N*-butyl-1,2-benzisoselenazol-3(2*H*)-one 11²**




2,2'-Diselenobis(*N*-ethylbenzamide) 12a³




N-ethyl-1,2-benzisoselenazol-3(2*H*)-one 12⁴




2,2'-Diselenobis(*N*-propylbenzamide) 13a³




N-propyl-1,2-benzisoselenazol-3(2*H*)-one 13⁴




2,2'-diselenobis[*N*-(*R*)-(-)-*sec*-butylbezamide 14a⁵




N-[(*R*)-(-)-*sec*-butyl]-1,2-benzisoselenazol-3(2*H*)-one 14⁵




2,2'-diselenobis[*N*-(*S*)-(+)-1-hydroxy-2-butanylbezamide] 15a⁵




***N*-[(*S*)-(+)-1-hydroxy-2-butanyl]-1,2-benzisoselenazol-3(2*H*)-one 15⁵**




2,2'-diselenobis[*N*-(*S*)-(-)- α -methylbenzylbezamide]16a⁵




***N*-[(*S*)-(-)- α -methylbenzyl]-1,2-benzisoselenazol-3(2*H*)-one 16⁵**




2,2'-diselenobis[*N*-(1*S*,2*S*)-(-)-*trans*-2-hydroxy-1-indanyl]bezamide] 17a⁵




***N*-(1*S*,2*S*)-(+)-*trans*-2-hydroxy-1-indanyl]-1,2-benzisoselenazol-3(2*H*)-one 17⁵**




2,2'-Diselenobis(*N*-cyclohexylbenzamide) 18a¹




N-cyclohexyl-1,2-benzisoselenazol-3(2*H*)-one 18²




2,2'-Diselenobis(*N*-phenylbenzamide) 19a⁶




N-phenyl-1,2-benzisoselenazol-3(2*H*)-one 19⁶



2,2'-Diselenobis(*N*-(4-methoxyphenyl)benzamide) 20a⁶



N-(4-methoxyphenyl)-1,2-benzisoselenazol-3(2*H*)-one 20⁶



V. References

1. A. J. Pacuła, K. B. Kaczor, A. Wojtowicz, J. Antosiewicz, A. Janecka, A. Długosz, T. Janecki, J. Scianowski, *Bioorg. Med. Chem.*, **2017**, *25*, 126–131.
2. A. J. Pacuła, J. Ścianowski and K. B. Aleksandrak, *RSC Adv.* **2014**, *4*, 48959–48962.
3. M. Pietka-Ottlik, P. Potaczek, E. Piasecki, J. Mlochowski, *Molecules* **2010**, *15*, 8214-8228.
4. M. Obieziurska, A. J. Pacuła, U. Juhas, J. Antosiewicz, J. Ścianowski, *Catalysts* **2018**, *8*, 493-507.
5. A. Laskowska, A. J. Pacuła-Miszewska, A. Długosz-Pokorska, A. Janecka, A. Wojtczak, J. Scianowski, *Materials*, **2022**, *15*, 2068-2082.
6. A. J. Pacuła, M. Obieziurska, J. Ścianowski, K. B. Kaczor, J. Antosiewicz, *Arkivoc*, **2018**, *3*, 144-155.