

Sulfonamides linked to sulfonate esters via hydrazone functionality: Synthesis, human carbonic anhydrase inhibitory activities, and molecular modeling studies

M. İhsan Han^a, Miyase Gözde Gündüz^b, Gökçe Alçı^c, Simone Giovannuzzi^d, Dönay Yuvalı^e, Claudiu T. Supuran^{d*}, Şengül Dilem Doğan^{f*}

^a Department of Pharmaceutical Chemistry, Faculty of Pharmacy, Erciyes University, 38039, Kayseri, Turkey

^b Department of Pharmaceutical Chemistry, Faculty of Pharmacy, Hacettepe University, 06100, Ankara, Turkey

^c Drug Application and Research Center, Erciyes University, 38039 Kayseri, Turkey

^d Neurofarba Department, Universita degli Studi di Firenze, Sezione di Scienze Farmaceutiche, Via Ugo Schiff 6, 50019 Sesto Fiorentino, Florence, Italy

^e Faculty of Pharmacy, Deparment of Analytical Chemistry, Erciyes University, Kayseri, 38039, Turkey

^f Department of Basic Sciences, Faculty of Pharmacy, Erciyes University, 38039, Kayseri, Turkey


* Corresponding authors:


Prof. Dr. Claudiu T. Supuran, Universita degli Studi di Firenze, Sezione di Scienze Farmaceutiche,
e-mail: claudiu.supuran@unifi.it

Dr. Şengül Dilem Doğan, Department of Basic Sciences, Faculty of Pharmacy, Erciyes University, 38039, Kayseri, Turkey, e-mail: dogandilem@gmail.com



Content	Page
¹ H NMR, ¹³ C NMR, HRMS, and FT-IR spectra of SH1-SH14	2-29
Binding mode of SH3 in the active site of hCA I and II	30

¹H NMR, ¹³C NMR, HRMS, and FT-IR spectra of **SH1**








User Spectra





¹H NMR, ¹³C NMR, HRMS, and FT-IR spectra of SH2




User Spectra





¹H NMR, ¹³C NMR, HRMS, and FT-IR spectra of **SH3**




User Spectra





¹H NMR, ¹³C NMR, HRMS, and FT-IR spectra of **SH4**




User Spectra





¹H NMR, ¹³C NMR, HRMS, and FT-IR spectra of **SH5**





User Spectra





¹H NMR, ¹³C NMR, HRMS, and FT-IR spectra of **SH6**




User Spectra





¹H NMR, ¹³C NMR, HRMS, and FT-IR spectra of **SH7**




User Spectra





¹H NMR, ¹³C NMR, HRMS, and FT-IR spectra of **SH8**




User Spectra





¹H NMR, ¹³C NMR, HRMS, and FT-IR spectra of **SH9**




User Spectra





¹H NMR, ¹³C NMR, HRMS, and FT-IR spectra of **SH10**




User Spectra





¹H NMR, ¹³C NMR, HRMS, and FT-IR spectra of **SH11**




User Spectra





¹H NMR, ¹³C NMR, HRMS, and FT-IR spectra of **SH12**




User Spectra





¹H NMR, ¹³C NMR, HRMS, and FT-IR spectra of **SH13**




User Spectra




¹H NMR, ¹³C NMR, HRMS, and FT-IR spectra of **SH14**



User Spectra





Binding mode of **SH3** in the active site of hCA I (A) and II (B). Ligand is represented as green color stick and balls, protein backbone as a white thin tube, hydrogen bonds as purple dashes, and important amino acid residues as gray and black sticks, for hCA I and II, respectively. The zinc ion is shown as a faded blue sphere.