

Supplementary Information

Ptxplatin: An multifunctional Pt(IV) antitumor prodrug

*Ran Zhang,^{a,b} Yueyue Zhang,^a Liumei Tang,^a Yixing Xu,^a Hao Li,^{a,b} Xueping Jiang,^a
Xiangdong Xin^a, and Zhongzheng Gui^{*a,b}*

^a School of Biotechnology, Jiangsu University of Science and Technology, Zhenjiang,
Jiangsu 212100, People's Republic of China

^b Sericultural Research Institute, Chinese Academy of Agricultural Sciences,
Zhenjiang, Jiangsu 212100, People's Republic of China

Table of Contents	Pages
Fig. S1. HR-MS spectrum of 1 .	S4
Fig. S2. HR-MS spectrum of 2 .	S4
Fig. S3. Structural characterization of PTX, 3 and 4 .	S5
Fig. S4. ¹³ C-NMR spectrum of compound 3 .	S5
Fig. S5. HR-MS spectrum of compound 3 .	S6
Fig. S6. HPLC characterization of compound 3 .	S6
Fig. S7. ¹³ C-NMR spectrum of compound 4 .	S7
Fig. S8. HR-MS spectrum of compound 4 .	S7
Fig. S9. HPLC characterization of compound 4 .	S8
Fig. S10. ¹ H-NMR spectrum of compound 5 .	S8
Fig. S11. ¹³ C-NMR spectrum of compound 5 .	S9
Fig. S12. HR-MS spectrum of compound 5 .	S9
Fig. S13. HPLC characterization of compound 5 .	S10
Fig. S14. ¹ H-NMR spectrum of compound 6 .	S10
Fig. S15. ¹³ C-NMR spectrum of compound 6 .	S11
Fig. S16. HR-MS spectrum of compound 6 .	S11
Fig. S17. HPLC characterization of compound 6 .	S12
Fig. S18. ¹ H-NMR spectrum of compound 7 .	S12
Fig. S19. ¹³ C-NMR spectrum of compound 7 .	S13
Fig. S20. HR-MS spectrum of compound 7 .	S13
Fig. S21. HPLC characterization of compound 7 .	S14
Fig. S22. ¹ H-NMR spectrum of compound 8 .	S14
Fig. S23. ¹³ C-NMR spectrum of compound 8 .	S15
Fig. S24. HR-MS spectrum of compound 8 .	S15
Fig. S25. HPLC characterization of compound 8 .	S16

Fig. S26. ¹ H-NMR spectrum of compound 9 .	S16
Fig. S27. ¹³ C-NMR spectrum of compound 9 .	S17
Fig. S28. HR-MS spectrum of compound 9 .	S17
Fig. S29. HPLC characterization of compound 9 .	S18
Fig. S30. ¹ H-NMR spectrum of compound 10 .	S18
Fig. S31. ¹³ C-NMR spectrum of compound 10 .	S19
Fig. S32. HR-MS spectrum of compound 10 .	S19
Fig. S33. HPLC characterization of compound 10 .	S20
Fig. S34. ¹ H-NMR spectrum of compound 11 .	S20
Fig. S35. ¹³ C-NMR spectrum of compound 11 .	S21
Fig. S36. HR-MS spectrum of compound 11 .	S21
Fig. S37. HPLC characterization of compound 11 .	S22
Fig. S38. ¹ H-NMR spectrum of compound 12 .	S22
Fig. S39. ¹³ C-NMR spectrum of compound 12 .	S23
Fig. S40. HR-MS spectrum of compound 12 .	S23
Fig. S41. HPLC characterization of compound 12 .	S24
Fig. S42. The expressions of MMP-2 in MCF-7 cells.	S24
Fig. S43. Cellular accumulation of Pt compounds in MCF-7 cells.	S25
Fig. S44. The results of HR-MS for cell extract after treated with 5 .	S25




Fig. S1. HR-MS spectrum of compound 1.




Fig. S2. HR-MS spectrum of compound 2.




Fig. S3. Structural characterization of PTX, **3** and **4**. ^1H NMR spectra of (a) PTX in CDCl_3 , (b) compound **3** in CDCl_3 , (c) PTX in $\text{DMSO-}d_6$, (d) compound **4** in $\text{DMSO-}d_6$.




Fig. S4. ^{13}C NMR spectrum of compound **3** in CDCl_3




Fig. S5. HR-MS spectrum of compound 3.




Fig. S6. HPLC characterization of compound 3.




Fig. S7. ^{13}C NMR spectrum of compound **4** in $\text{DMSO-}d_6$.




Fig. S8. HR-MS spectrum of compound **4**.




Fig. S9. HPLC characterization of compound 4.




Fig. S10. ¹H-NMR spectrum of compound 5 in DMSO-*d*₆.




Fig. S11. ^{13}C -NMR spectrum of compound **5** in $\text{DMSO-}d_6$.




Fig. S12. HR-MS spectrum (positive mode) of compound **5**.




Fig. S13. HPLC characterization of compound **5**.




Fig. S14. ¹H-NMR spectrum of compound **6** in DMSO-*d*₆.




Fig. S15. ^{13}C -NMR spectrum of compound **6** in DMSO- d_6 .




Fig. S16. HR-MS spectrum (positive mode) of compound **6**.




Fig. S17. HPLC characterization of compound **6**.




Fig. S18. ¹H-NMR spectrum of compound **7** in DMSO-*d*₆.




Fig. S19. ^{13}C -NMR spectrum of compound 7 in DMSO- d_6 .




Fig. S20. HR-MS spectrum (positive mode) of compound 7.




Fig. S21. HPLC characterization of compound 7.




Fig. S22. ¹H-NMR spectrum of compound 8 in DMSO-*d*₆.




Fig. S23. ^{13}C -NMR spectrum of compound **8** in $\text{DMSO-}d_6$.




Fig. S24. HR-MS spectrum (positive mode) of compound **8**.




Fig. S25. HPLC characterization of compound **8**.




Fig. S26. ¹H-NMR spectrum of compound **9** in DMSO-*d*₆.





Fig. S27. ^{13}C -NMR spectrum of compound **9** in $\text{DMSO-}d_6$.



S28. HR-MS spectrum (positive mode) of compound **9**.




Fig. S31. ^{13}C -NMR spectrum of compound **10** in DMSO- d_6 .




Fig. S32. HR-MS spectrum (positive mode) of compound **10**.




Fig. S33. HPLC characterization of compound 10.




Fig. S34. ¹H-NMR spectrum of compound 11 in DMSO-*d*₆.




Fig. S35. ^{13}C -NMR spectrum of compound **11** in $\text{DMSO-}d_6$.




Fig. S36. HR-MS spectrum (positive mode) of compound **11**.




Fig. S37. HPLC characterization of compound 11.




Fig. S38. ¹H-NMR spectrum of compound 12 in DMSO-*d*₆.




Fig. S39. ^{13}C -NMR spectrum of compound 12 in $\text{DMSO-}d_6$.




Fig. S40. HR-MS spectrum (positive mode) of compound 12.




Fig. S41. HPLC characterization of compound **12**.




Fig. S42. Western blotting analysis of MMP-2 in MCF-7 cells after treatment with various compounds (5 μ M) for 24 h, Mixture stands for a combination of CDDP (5 μ M) and PTX (5 μ M).




Fig. S43. Cellular accumulation of the indicated compounds at 10 μ M for 9 h treatment.




Fig. S44 The results of HPLC and HR-MS analysis for MCF-7 cells extract after treated with **5** for 4 h, using untreated cells extract as controls.