

Supporting information

Novel one-pot synthesis, characterization, and antifungal and antibacterial activity of a small library of 2-aryloxy-1,4-naphthoquinone derivatives

Chaves-Carballo, K.^a, Lamoureux, G.V.^a, Perez, A.L.^a, Bella Cruz, A.,^b Cechinel Filho, V.^b

^a Centro de Investigaciones en Productos Naturales and Escuela de Química, Universidad de Costa Rica, San Pedr2060, San José, Costa Rica.

^b Núcleo de Investigações Químico-Farmacêuticas (NIQFAR), CCS, Universidade do Vale do Itajaí (UNIVALI), Itajaí, SC, Brazil.

Table of contents

2-(2-methoxyphenoxy)-1,4-naphthoquinone 34	3
2-(3-methoxyphenoxy)-1,4-naphthoquinone 35	6
2-(4-methoxyphenoxy)-1,4-naphthoquinone 36	9
2-(3-nitrophenoxy)-1,4-naphthoquinone 37	12
2-(2-bromophenoxy)-1,4-naphthoquinone 38	15
2-(4-bromophenoxy)-1,4-naphthoquinone 39	18
2-(2,4-dibromophenoxy)-1,4-naphthoquinone 40	21
2-(2,4,6-tribromophenoxy)-1,4-naphthoquinone 41	24
2-(3,4-dichlorophenoxy)-1,4-naphthoquinone 42	27
2-(4-tritylphenoxy)-1,4-naphthoquinone 43	30
2-(4-phenylphenoxy)-1,4-naphthoquinone 44	33
2-(4-tert-butylphenoxy)-1,4-naphthoquinone 45	36
2-(2-chlorophenoxy)-1,4-naphthoquinone 46	39
2-(3-chlorophenoxy)-1,4-naphthoquinone 47	42
2-(4-chlorophenoxy)-1,4-naphthoquinone 48	45
2-phenoxy-1,4-naphthoquinone 49	48
2-(2-nitrophenoxy)-1,4-naphthoquinone 50	51
2-(2,4,5-trichlorophenoxy)-1,4-naphthoquinone 51	54
2-(2,4,6-trichlorophenoxy)-1,4-naphthoquinone 52	57


2-(4-chloro-3,5-dimethylphenoxy)-1,4-naphthoquinone 53	60
2-(2-isopropyl-5-methylphenoxy)-1,4-naphthoquinone 54	63
2-(5-chloro-2-(2,4-dichlorophenoxy)phenoxy)-1,4-naphthoquinone 55	67
2-(4-allyl-2-methoxyphenoxy)-1,4-naphthoquinone 56	70
2-(4-[2-(4-hydroxyphenyl)propan-2-yl]phenoxy)-1,4-naphthoquinone 57	74
2-(4-[2-(4-hydroxyphenyl)cyclohexyl]phenoxy)-1,4-naphthoquinone 58	77
2-(2-methylphenoxy)-1,4-naphthoquinone 59	80
2-(3-methylphenoxy)-1,4-naphthoquinone 60	83
2-(4-methylphenoxy)-1,4-naphthoquinone 61	86
2-(4-nitrophenoxy)-1,4-naphthoquinone 62	89
2-(2,4-dichlorophenoxy)-1,4-naphthoquinone 63	92

Synthesis procedures and characterization of 1,4-naphthoquinone phenolic ether derivatives


2-(2-methoxyphenoxy)-1,4-naphthoquinone **34**

34 was obtained according to general procedure I 2.1.1 and purified by hot extraction-filtration with MTBE, cold base wash (MTBE as solvent and NaOH 0.5 mol/L as base) and recrystallization with isooctane. Yield 141 mg (68%), yellow crystals, mp = (151.0 – 153.0) °C. ^1H NMR (600 MHz, CDCl_3): δ (ppm) = 8.21 (1H, d, $J = 9$ Hz, ArH), 8.06 (1H, d, $J = 9$ Hz, ArH), 7.75 (2H, m, ArH), 7.28 (1H, t, $J_{\text{ortho}} = 8$ Hz, ArH), 7.13 (1H, d, $J_{\text{ortho}} = 9$ Hz, ArH), 7.03 (1H, t, $J_{\text{ortho}} = 8$ Hz, ArH), 7.00 (1H, d, $J_{\text{ortho}} = 8$ Hz, ArH), 5.85 (1H, s, ArH), 3.80 (3H, s, CH_3). ^{13}C NMR (150 MHz, CDCl_3): δ (ppm) = 185.2, 179.7, 159.5, 150.7, 140.9, 134.2, 133.4, 132.1, 131.2, 127.7, 126.7, 126.1, 122.4, 121.4, 113.0, 112.7, 55.7. IR (KBr pellet) cm^{-1} : 3068, 2958, 1683, 1652, 1611, 1499, 1261, 1205. UV-Vis (MeOH) nm: 201, 243 (max), 271, 331. TOF-MS: m/z $[\text{M}+\text{H}]^+$ calc for $\text{C}_{17}\text{H}_{13}\text{O}_4$: 281.0814; found: 281.0819. Purity: 99.9%. MIC (μM): 892 (*S. aureus*), > 3000 (*E. coli*, *C. albicans*).


2-(2-methoxyphenoxy)-1,4-naphthoquinone **34** ^1H -NMR




2-(2-methoxyphenoxy)-1,4-naphthoquinone **34** ^{13}C -NMR




2-(2-methoxyphenoxy)-1,4-naphthoquinone **34** UV-Vis (MeOH)



2-(2-methoxyphenoxy)-1,4-naphthoquinone **34** IR (KBr pellet)




2-(2-methoxyphenoxy)-1,4-naphthoquinone **34** HPLC chromatogram (274 nm, 100% MeCN)




2-(3-methoxyphenoxy)-1,4-naphthoquinone **35**

35 was obtained according to general procedure I 2.1.1 and purified by hot extraction-filtration with MTBE, cold base wash (MTBE as solvent and NaOH 0.5 mol/L as base) and recrystallization with isooctane. Yield 109 mg (52%), orange crystals, mp = (84.0 – 87.0) °C. ^1H NMR (600 MHz, CDCl_3): δ (ppm) = 8.23 (1H, d, J = 7.5 Hz, ArH), 8.09 (1H, d, J = 7.5 Hz, ArH), 7.80 (2H, m, ArH), 7.38 (1H, t, J_{ortho} = 7.5 Hz, ArH), 6.88 (1H, d, J_{ortho} = 9 Hz, ArH), 6.76 (1H, d, J_{ortho} = 8 Hz, ArH), 6.71 (1H, s, ArH), 6.05 (1H, s, ArH), 3.84 (3H, s, CH_3). ^{13}C NMR (150 MHz, CDCl_3): δ (ppm) = 185.0, 179.9, 161.2, 160.4, 153.6, 134.4, 133.5, 132.0, 131.1, 130.8, 126.8, 126.2, 113.5, 113.0, 112.3, 55.5. UV-Vis (MeOH) nm: 211, 244 (max), 270, 331. IR (KBr pellet) cm^{-1} : 3053, 2960, 1678, 1653, 1611, 1584, 1264, 1204. TOF-MS: m/z $[\text{M}+\text{H}]^+$ calc for $\text{C}_{17}\text{H}_{13}\text{O}_4$: 281.0814; found: 281.0812. Purity: 95.9%. MIC (μM): 223 (*S. aureus*), 446 (*C. albicans*), >3000 (*E. coli*).


2-(3-methoxyphenoxy)-1,4-naphthoquinone **35** ^1H NMR




2-(3-methoxyphenoxy)-1,4-naphthoquinone **35** ^{13}C RMN




2-(3-methoxyphenoxy)-1,4-naphthoquinone **35** UV-Vis (MeOH)



2-(3-methoxyphenoxy)-1,4-naphthoquinone 35 IR (KBr pellet)




2-(3-methoxyphenoxy)-1,4-naphthoquinone 35 HPLC chromatogram (286 nm, 40% H₂O/MeOH)




2-(4-methoxyphenoxy)-1,4-naphthoquinone **36**

36 was obtained according to general procedure I 2.1.1 and purified by hot extraction-filtration with MTBE, chromatography column (CH₂Cl₂ as eluent, isocratically) and recrystallization with isooctane. Yield 201 mg (96%), yellow crystals, mp = (133.0 – 134.0) °C. ¹H NMR (600 MHz, CDCl₃): δ (ppm) = 8.20 (1H, d, *J* = 8 Hz, ArH), 8.06 (1H, d, *J* = 8 Hz, ArH), 7.76 (2H, m, ArH), 7.05 (2H, d, *J*_{ortho} = 8 Hz, ArH), 6.96 (2H, d, *J*_{ortho} = 8 Hz, ArH), 5.95 (1H, s, ArH), 3.83 (3H, s, CH₃). ¹³C NMR (150 MHz, CDCl₃): δ (ppm) = 185.0, 179.9, 161.2, 160.4, 153.6, 132.0, 131.1, 126.8, 126.2, 121.1, 117.6, 113.0, 55.5. IR (KBr pellet) cm⁻¹: 3073, 2923, 1680, 1650, 1619, 1503, 1259, 1219. UV-Vis (MeOH) nm: 204, 243 (max), 274, 333. TOF-MS: *m/z* [M+H]⁺ calc for C₁₇H₁₃O₄: 281.0814; found: 281.0817. Purity: 99.3%. MIC (μM): 1784 (*S. aureus*), >3000 (*C. albicans*), >3000 (*E. coli*).


2-(4-methoxyphenoxy)-1,4-naphthoquinone **36** ¹H NMR




2-(4-methoxyphenoxy)-1,4-naphthoquinone **36** ^{13}C NMR




2-(4-methoxyphenoxy)-1,4-naphthoquinone **36** UV-Vis (MeOH)



2-(4-methoxyphenoxy)-1,4-naphthoquinone 36 IR (KBr pellet)




2-(4-methoxyphenoxy)-1,4-naphthoquinone 36 HPLC chromatogram (270 nm, 100% MeCN)




2-(3-nitrophenoxy)-1,4-naphthoquinone **37**

37 was obtained according to general procedure I 2.1.1 and purified by hot extraction-filtration with 50% n-heptane/toluene and recrystallization with n-heptane. Yield 173 mg (58%), yellow crystals, mp = (166.0 – 167.0) °C. ^1H NMR (600 MHz, CDCl_3): δ (ppm) = 8.21 (2H, d, $J = 6$ Hz, ArH), 8.10 (1H, dd, $J = 1.2$ Hz, $J = 6.9$ Hz, ArH), 8.04 (1H, t, $J_{\text{meta}} = 1.2$ Hz, ArH), 7.80 (2H, m, ArH), 7.68 (1H, t, $J_{\text{ortho}} = 8.1$ Hz, ArH), 7.52 (1H, ddd, $J_{\text{meta}} = 1.2$ Hz, $J_{\text{meta}} = 2.3$ Hz, $J_{\text{ortho}} = 8.1$ Hz, ArH), 6.05 (1H, s, ArH). ^{13}C NMR (150 MHz, CDCl_3): δ (ppm) = 184.4, 179.3, 159.2, 153.5, 149.4, 134.7, 133.9, 131.8, 131.2, 130.9, 127.1, 127.0, 126.4, 121.3, 116.4, 115.0. IR (KBr pellet) cm^{-1} : 3085, 1676, 1654, 1608, 1527, 1353, 1261, 1222. UV-Vis (MeOH) nm: 209 (max), 241, 272, 331. TOF-MS: m/z $[\text{M}+\text{H}]^+$ calc for $\text{C}_{16}\text{H}_{10}\text{NO}_5$: 296.0559; found: 296.0558. Purity: 99.0%. MIC (μM): 21.2 (*S. aureus*), > 3000 (*C. albicans*), > 3000 (*E. coli*).


2-(3-nitrophenoxy)-1,4-naphthoquinone **37** ^1H NMR




2-(3-nitrophenoxy)-1,4-naphthoquinone 37 ^{13}C NMR




2-(3-nitrophenoxy)-1,4-naphthoquinone 37 UV-Vis (MeOH)



2-(3-nitrophenoxy)-1,4-naphthoquinone 37 IR (KBr pellet)




2-(3-nitrophenoxy)-1,4-naphthoquinone 37 HPLC chromatogram (268 nm, 100% MeCN)




2-(2-bromophenoxy)-1,4-naphthoquinone **38**

38 was obtained according to general procedure I 2.1.1 and purified by hot extraction-filtration with MTBE, column chromatography (1% triethylamine/benzene as eluent, isocratically) and recrystallization with isooctane. Yield 287 mg (87%), yellow crystals, mp = (119.0 – 121.0) °C. ^1H NMR (600 MHz, CDCl_3): δ (ppm) = 8.23 (1H, d, $J = 6.4$ Hz, ArH), 8.08 (1H, d, $J = 6.4$ Hz, ArH), 7.78 (2H, m, ArH), 7.69 (1H, d, $J_{\text{ortho}} = 8.6$ Hz, ArH), 7.42 (1H, t, $J_{\text{ortho}} = 8.6$ Hz, ArH), 7.22 (1H, t, $J_{\text{ortho}} = 8.6$ Hz, ArH), 7.21 (1H, d, $J_{\text{ortho}} = 6.4$ Hz, ArH), 5.83 (1H, s, ArH). ^{13}C NMR (150 MHz, CDCl_3): δ (ppm) = 184.8, 179.3, 158.7, 149.6, 134.4, 134.3, 133.6, 131.9, 131.0, 129.4, 128.1, 126.8, 126.2, 123.0, 115.6, 113.5. IR (KBr pellet) cm^{-1} : 3068, 1684, 1654, 1612, 1595, 1263, 1225, 660. UV-Vis (MeOH) nm: 203, 209 (max), 215, 249, 271, 333. TOF-MS: m/z $[\text{M}+\text{H}]^+$ calc for $\text{C}_{16}\text{H}_{10}\text{O}_3\text{Br}$: 328.9813; found: 328.9806. Purity: 99.7%. MIC (μM): 760 (*S. aureus*), 190 (*C. albicans*), >3000 (*E. coli*).


2-(2-bromophenoxy)-1,4-naphthoquinone **38** ^1H NMR




2-(2-bromophenoxy)-1,4-naphthoquinone 38 ^{13}C NMR




2-(2-bromophenoxy)-1,4-naphthoquinone 38 UV-Vis (MeOH)



2-(2-bromophenoxy)-1,4-naphthoquinone **38** IR (KBr pellet)




2-(2-bromophenoxy)-1,4-naphthoquinone **38** HPLC chromatogram (270 nm, 100% MeCN)




2-(4-bromophenoxy)-1,4-naphthoquinone **39**

39 was obtained according to general procedure I 2.1.1 and purified by hot extraction-filtration with MTBE and recrystallization with isooctane. Yield 246 mg (75%), yellow needles, mp = (134.0 – 135.0) °C. ^1H NMR (600 MHz, CDCl_3): δ (ppm) = 8.20 (1H, dd, $J = 2$ Hz, $J = 6$ Hz, ArH), 8.07 (1H, dd, $J = 6$ Hz, ArH), 7.77 (2H, m, ArH), 7.59 (2H, d, $J_{\text{ortho}} = 8$ Hz, ArH), 7.04 (2H, d, $J_{\text{ortho}} = 8$ Hz, ArH), 5.98 (1H, s, ArH). ^{13}C NMR (150 MHz, CDCl_3): δ (ppm) = 184.7, 179.7, 160.0, 151.8, 134.6, 133.7, 133.5, 131.9, 131.0, 126.8, 126.3, 122.9, 119.8, 113.7. IR (KBr pellet) cm^{-1} : 3095, 1684, 1652, 1612, 1479, 1263, 1232, 717. UV-Vis (MeOH) nm: 221, 249 (max), 269, 335. TOF-MS: m/z $[\text{M}+\text{H}]^+$ calc for $\text{C}_{16}\text{H}_{10}\text{O}_3\text{Br}$: 328.9813; found: 328.9814. Purity: 99.7%. MIC (μM): >3000 (*S. aureus*), > 3000 (*C. albicans*), >3000 (*E. coli*).


2-(4-bromophenoxy)-1,4-naphthoquinone **39** ^1H NMR




2-(4-bromophenoxy)-1,4-naphthoquinone **39** ^{13}C NMR




2-(4-bromophenoxy)-1,4-naphthoquinone **39** UV-Vis (MeOH)



2-(4-bromophenoxy)-1,4-naphthoquinone **39 IR (KBr pellet)**




2-(4-bromophenoxy)-1,4-naphthoquinone **39 HPLC chromatogram (270 nm, 100% MeCN)**




2-(2,4-dibromophenoxy)-1,4-naphthoquinone **40**

40 was obtained according to general procedure I 2.1.1 and purified by hot extraction-filtration with MTBE and recrystallization with isooctane. Yield 251 mg (62%), yellow solid, mp = (133.0 – 134.0) °C. ^1H NMR (600 MHz, CDCl_3): δ (ppm) 8.21 (1H, dd, $J_{\text{ortho}} = 7.5$ Hz, ArH), 8.08 (1H, dd, $J_{\text{ortho}} = 7.5$ Hz, ArH), 7.84 (1H, d, $J_{\text{meta}} = 1.9$ Hz, ArH), 7.78 (2H, m, ArH), 7.54 (1H, dd, $J_{\text{meta}} = 1.9$ Hz, $J_{\text{ortho}} = 9.4$ Hz, ArH), 7.09 (1H, d, $J_{\text{ortho}} = 7.5$ Hz, ArH), 5.84 (1H, s, ArH). ^{13}C NMR (150 MHz, CDCl_3): δ (ppm) = 184.5, 179.1, 158.4, 149.0, 136.7, 134.6, 133.7, 132.5, 131.9, 131.0, 126.9, 126.3, 124.1, 120.4, 116.7, 113.8. IR (KBr pellet) cm^{-1} : 3086, 1686, 1652, 1612, 1465, 1260, 1236, 783, 717. UV-Vis (MeOH) nm: 221, 249 (max), 270, 333. TOF-MS: m/z $[\text{M}+\text{H}]^+$ calc for $\text{C}_{16}\text{H}_9\text{O}_3\text{Br}_2$: 406.8918; found: 406.8923. Purity: 98.9%. MIC (μM): >2000 (*S. aureus*), >2000 (*C. albicans*), >2000 (*E. coli*).


2-(2,4-dibromophenoxy)-1,4-naphthoquinone **40** ^1H NMR




2-(2,4-dibromophenoxy)-1,4-naphthoquinone **40** ^{13}C NMR




2-(2,4-dibromophenoxy)-1,4-naphthoquinone **40** UV-Vis (MeOH)



2-(2,4-dibromophenoxy)-1,4-naphthoquinone **40** IR (KBr pellet)




2-(2,4-dibromophenoxy)-1,4-naphthoquinone **40** HPLC chromatogram (270 nm, 100% MeCN)




2-(2,4,6-tribromophenoxy)-1,4-naphthoquinone **41**

41 was obtained according to general procedure I 2.1.1 and purified by hot extraction-filtration with CH₂Cl₂, column chromatography (50% benzene/cyclohexane to dissolve the solid and benzene as eluent, isocratically) and recrystallization with methanol. Yield 316 mg (65%), yellow solid, mp = (170.0 – 172.0) °C. ¹H NMR (600 MHz, CDCl₃): δ (ppm) 8.23 (1H, m, ArH), 8.09 (1H, m, ArH), 7.79 (2H, m, ArH), 7.36 (2H, s, ArH), 5.82 (1H, s, ArH). ¹³C NMR (150 MHz, CDCl₃): δ (ppm) = 184.4, 178.5, 156.4, 148.9, 134.6, 134.2, 133.8, 131.9, 131.0, 126.9, 126.4, 117.8, 113.4, 112.7. IR (KBr pellet) cm⁻¹: 3067, 1684, 1654, 1618, 1593, 1237, 1176, 714. UV-Vis (MeCN) nm: 231, 271 (max). TOF-MS: m/z [M+H]⁺ calc for C₁₆H₈O₃Br₃: 484.8024; found: 484.8034. Purity: 97.3%. MIC (μM): >2000 (*S. aureus*), >2000 (*C. albicans*), >2000 (*E. coli*).


2-(2,4,6-tribromophenoxy)-1,4-naphthoquinone **41** ¹H NMR




2-(2,4,6-tribromophenoxy)-1,4-naphthoquinone **41** ^{13}C NMR




2-(2,4,6-tribromophenoxy)-1,4-naphthoquinone **41** UV-Vis (MeCN)



2-(2,4,6-tribromophenoxy)-1,4-naphthoquinone **41** IR (KBr pellet)




2-(2,4,6-tribromophenoxy)-1,4-naphthoquinone **41** HPLC chromatogram (270 nm, 100% MeCN)




2-(3,4-dichlorophenoxy)-1,4-naphthoquinone **42**

42 was obtained according to general procedure I 2.1.1 and purified by hot extraction-filtration with MTBE and recrystallization with isooctane. Yield 160 mg (50%), yellow needles, mp = (162.0 – 163.0) °C. ^1H NMR (600 MHz, CDCl_3): δ (ppm) 8.19 (1H, m, ArH), 8.08 (1H, m, ArH), 7.78 (2H, m, ArH), 7.54 (1H, d, $J_{ortho} = 8$ Hz, ArH), 7.29 (1H, d, $J_{meta} = 2$ Hz, ArH), 7.03 (1H, dd, $J_{meta} = 2$ Hz, $J_{ortho} = 8$ Hz, ArH), 6.03 (1H, s, ArH). ^{13}C NMR (150 MHz, CDCl_3): δ (ppm) = 184.6, 179.4, 159.5, 151.5, 134.6, 134.1, 133.8, 131.8, 131.8, 130.9, 130.7, 126.9, 126.3, 123.2, 120.5, 114.6. IR (KBr pellet) cm^{-1} : 3069, 1683, 1643, 1624, 1595, 1466, 1265, 1216, 986. UV-Vis (MeOH) nm: 207 (max), 241, 276, 331. TOF-MS: m/z $[\text{M}+\text{H}]^+$ calc for $\text{C}_{16}\text{H}_9\text{O}_3\text{Cl}_2$: 318.9929; found: 318.9930. Purity: 98.7%. MIC (μM): >3000 (*S. aureus*), >3000 (*C. albicans*), >3000 (*E. coli*).


2-(3,4-dichlorophenoxy)-1,4-naphthoquinone **42** ^1H NMR




2-(3,4-dichlorophenoxy)-1,4-naphthoquinone **42** ^{13}C NMR




2-(3,4-dichlorophenoxy)-1,4-naphthoquinone **42** UV-Vis (MeOH)



2-(3,4-dichlorophenoxy)-1,4-naphthoquinone **42** IR (KBr pellet)




2-(3,4-dichlorophenoxy)-1,4-naphthoquinone **42** HPLC chromatogram (270 nm, 100% MeCN)




2-(4-tritylphenoxy)-1,4-naphthoquinone **43**

43 was obtained according to general procedure I 2.1.1, but with lesser amount of reagents (0.40 mmol of 2-bromo-1,4-naphthoquinone, 0.40 mmol of 4-tritylphenol and 0.60 mmol of dried Cs_2CO_3), and purified by hot extraction-filtration with CH_2Cl_2 and column chromatography (1% triethylamine/benzene as eluent, isocratically). Yield 163 mg (82%), beige solid, mp = (256.0 – 257.0) °C. ^1H NMR (600 MHz, CDCl_3): δ (ppm) 8.20 (1H, dd, $J = 2.0$ Hz, $J = 7.7$ Hz, ArH), 8.08 (1H, dd, $J = 2.0$ Hz, $J = 7.7$ Hz, ArH), 7.76 (2H, m, ArH), 7.30 (6H, t, ArH), 7.28 (3H, d, ArH), 7.22 (6H, t, ArH), 7.21 (2H, d, ArH), 7.02 (2H, d, $J_{\text{ortho}} = 8.3$ Hz, ArH), 6.07 (1H, s, ArH). ^{13}C NMR (150 MHz, CDCl_3): δ (ppm) = 185.1, 179.9, 160.3, 150.5, 146.3, 145.4, 134.4, 133.5, 133.2, 132.0, 131.1, 131.0, 127.4, 126.8, 126.3, 126.3, 120.0, 113.3, 64.7. IR (KBr pellet) cm^{-1} : 3059, 1681, 1655, 1600, 1496, 1261, 1225. UV-Vis (MeCN) nm: 218 (max), 245, 269, 331. TOF-MS: m/z $[\text{M}+\text{H}]^+$ calc for $\text{C}_{35}\text{H}_{25}\text{O}_3$: 493.1804; found: 493.1801. Purity: 96.0%. MIC (μM): >2000 (*S. aureus*), >2000 (*C. albicans*), >2000 (*E. coli*).


2-(4-tritylphenoxy)-1,4-naphthoquinone **43** ^1H NMR




2-(4-tritylphenoxy)-1,4-naphthoquinone **43** ^{13}C NMR




2-(4-tritylphenoxy)-1,4-naphthoquinone **43** UV-Vis (MeCN)



2-(4-tritylphenoxy)-1,4-naphthoquinone **43** IR (KBr pellet)




2-(4-tritylphenoxy)-1,4-naphthoquinone **43** HPLC chromatogram (270 nm, 100% MeCN)




2-(4-phenylphenoxy)-1,4-naphthoquinone **44**

44 was obtained according to general procedure I 2.1.1 and purified by column chromatography (benzene as eluent, isocratically) and recrystallization with methanol. Yield 53 mg (16%), beige needles, mp = (199.0 – 200.0) °C. ^1H NMR (600 MHz, CDCl_3): δ (ppm) 8.22 (1H, d, J = 8 Hz, ArH), 8.08 (1H, d, J = 8 Hz, ArH), 7.77 (2H, m, ArH), 7.67 (2H, d, J_{ortho} = 8 Hz, ArH), 7.59 (2H, d, J_{ortho} = 8 Hz, ArH), 7.47 (2H, t, J_{ortho} = 8 Hz, ArH), 7.38 (1H, t, J_{ortho} = 8 Hz, ArH), 7.21 (2H, d, J_{ortho} = 8 Hz, ArH), 6.06 (1H, s, ArH). ^{13}C NMR (150 MHz, CDCl_3): δ (ppm) 185.0, 179.9, 160.5, 152.0, 139.9, 134.5, 133.6, 132.0, 131.1, 129.1, 128.9, 127.7, 127.2, 126.8, 126.3, 121.4, 113.5. IR (KBr pellet) cm^{-1} : 3036, 1683, 1651, 1615, 1593, 1263, 1226. UV-Vis (MeOH) nm: 212, 247 (max), 332. TOF-MS: m/z $[\text{M}+\text{H}]^+$ calc for $\text{C}_{22}\text{H}_{15}\text{O}_3$: 327.1021; found: 327.1021. Purity: 99.4%. MIC (μM): >3000 (*S. aureus*), >3000 (*C. albicans*), >3000 (*E. coli*).


2-(4-phenylphenoxy)-1,4-naphthoquinone **44** ^1H NMR




2-(4-phenylphenoxy)-1,4-naphthoquinone **44** ^{13}C NMR




2-(4-phenylphenoxy)-1,4-naphthoquinone **44** UV-Vis (MeOH)



2-(4-phenylphenoxy)-1,4-naphthoquinone **44** IR (KBr pellet)




2-(4-phenylphenoxy)-1,4-naphthoquinone **44** HPLC chromatogram (270 nm, 100% MeCN)




2-(4-tert-butylphenoxy)-1,4-naphthoquinone **45**

45 was obtained according to general procedure I 2.1.1 and purified by hot extraction-filtration with MTBE, column chromatography (benzene as eluent, isocratically) and recrystallization with isooctane. Yield 171 mg (56%), yellow needles, mp = (151.0 – 152.0) °C. ^1H NMR (600 MHz, CDCl_3): δ (ppm) 8.22 (1H, d, J = 6 Hz, ArH), 8.06 (1H, d, J = 6 Hz, ArH), 7.77 (2H, m, ArH), 7.46 (2H, d, J_{ortho} = 7 Hz, ArH), 7.05 (2H, d, J_{ortho} = 9 Hz, ArH), 5.97 (1H, s, ArH), 1.35 (9H, s, CH_3). ^{13}C NMR (150 MHz, CDCl_3): δ (ppm) 185.1, 180.0, 160.8, 150.2, 149.7, 134.4, 133.5, 132.0, 131.1, 127.2, 126.8, 126.2, 120.4, 113.2, 34.6, 31.4. IR (KBr pellet) cm^{-1} : 3061, 2958, 1678, 1657, 1598, 1504, 1262, 1213. UV-Vis (MeCN) nm: 201, 249 (max), 270, 331. TOF-MS: m/z $[\text{M}+\text{H}]^+$ calc for $\text{C}_{20}\text{H}_{19}\text{O}_3$: 307.1334; found: 307.1335. Purity: 98.8%. MIC (μM): >3000 (*S. aureus*), >3000 (*C. albicans*), >3000 (*E. coli*).


2-(4-tert-butylphenoxy)-1,4-naphthoquinone **45** ^1H NMR




2-(4- *tert*-butylphenoxy)-1,4-naphthoquinone **45** ^{13}C NMR




2-(4- *tert*-butylphenoxy)-1,4-naphthoquinone **45** UV-Vis (MeCN)



2-(4- *tert*-butylphenoxy)-1,4-naphthoquinone **45** IR (KBr pellet)




2-(4- *tert*-butylphenoxy)-1,4-naphthoquinone **45** HPLC chromatogram (270 nm, 100% MeCN)




2-(2-chlorophenoxy)-1,4-naphthoquinone **46**

46 was obtained according to general procedure I 2.1.1 and purified by hot extraction-filtration with MTBE, column chromatography (gradient elution starting with 15% MTBE/cyclohexane to 50% MTBE/cyclohexane) and recrystallization with isooctane. Yield 124 mg (58%), yellow solid, mp = (95.0 – 96.0) °C. ^1H NMR (600 MHz, CDCl_3): δ (ppm) 8.22 (1H, d, $J = 7$ Hz, ArH), 8.07 (1H, d, $J = 7$ Hz, ArH), 7.77 (2H, m, ArH), 7.52 (1H, dd, $J_{\text{meta}} = 3$ Hz, $J_{\text{ortho}} = 8$ Hz, ArH), 7.37 (1H, td, $J_{\text{meta}} = 3$ Hz, $J_{\text{ortho}} = 8$ Hz, ArH), 7.29 (1H, td, $J_{\text{meta}} = 3$ Hz, $J_{\text{ortho}} = 8$ Hz, ArH), 7.22 (1H, dd, $J_{\text{meta}} = 3$ Hz, $J_{\text{ortho}} = 8$ Hz, ArH), 5.83 (1H, s, ArH). ^{13}C NMR (150 MHz, CDCl_3): δ (ppm) = 184.8, 179.3, 158.8, 148.5, 134.5, 133.7, 132.0, 131.3, 131.1, 128.7, 127.9, 126.9, 126.7, 126.3, 123.1, 113.4. IR (KBr pellet) cm^{-1} : 3068, 1683, 1653, 1611, 1499, 1261, 1205, 976. UV-Vis (MeOH) nm: 203 (max), 243, 270, 332. TOF-MS: m/z $[\text{M}+\text{H}]^+$ calc for $\text{C}_{16}\text{H}_{10}\text{O}_3\text{Cl}$: 285.0318; found: 285.0320. Purity: 97.9%. MIC (μM): 220 (*S. aureus*), 220 (*C. albicans*), >3000 (*E. coli*).


2-(2-chlorophenoxy)-1,4-naphthoquinone **46** ^1H NMR




2-(2-chlorophenoxy)-1,4-naphthoquinone **46** ^{13}C NMR




2-(2-chlorophenoxy)-1,4-naphthoquinone **46** UV-Vis (MeOH)



2-(2-chlorophenoxy)-1,4-naphthoquinone **46** IR (KBr pellet)




2-(2-chlorophenoxy)-1,4-naphthoquinone **46** HPLC chromatogram (286 nm, 40% H₂O/MeOH)




2-(3-chlorophenoxy)-1,4-naphthoquinone **47**

47 was obtained according to general procedure I 2.1.1 and purified by hot extraction-filtration with MTBE, column chromatography (gradient elution starting with 10% MTBE/cyclohexane to 20% MTBE/cyclohexane) and recrystallization with isooctane. Yield 103 mg (48%), yellow solid, mp = (103.0 – 105.0) °C. ^1H NMR (600 MHz, CDCl_3): δ (ppm) 8.20 (1H, d, $J = 7$ Hz, ArH), 8.08 (1H, d, $J = 7$ Hz, ArH), 7.78 (2H, m, ArH), 7.40 (1H, t, $J_{\text{ortho}} = 8$ Hz, ArH), 7.31 (1H, d, $J_{\text{ortho}} = 8$ Hz, ArH), 7.17 (1H, s, ArH), 7.06 (1H, d, $J_{\text{ortho}} = 8$ Hz, ArH), 6.00 (1H, s, ArH). ^{13}C NMR (150 MHz, CDCl_3): δ (ppm) = 184.8, 179.6, 159.9, 153.3, 135.7, 134.6, 133.7, 131.9, 131.9, 131.0, 127.0, 126.9, 126.3, 121.7, 119.4, 113.9. IR (KBr pellet) cm^{-1} : 3071, 1681, 1651, 1614, 1585, 1262, 1222, 982. UV-Vis (MeOH) nm: 204 (max), 248, 273, 332. TOF-MS: m/z $[\text{M}+\text{H}]^+$ calc for $\text{C}_{16}\text{H}_{10}\text{O}_3\text{Cl}$: 285.0318; found: 285.0323. Purity: 92.3%. MIC (μM): 878 (*S. aureus*), 220 (*C. albicans*), >3000 (*E. coli*).


2-(3-chlorophenoxy)-1,4-naphthoquinone **47** ^1H NMR




2-(3-chlorophenoxy)-1,4-naphthoquinone **47** ^{13}C NMR




2-(3-chlorophenoxy)-1,4-naphthoquinone **47** UV-Vis (MeOH)



2-(3-chlorophenoxy)-1,4-naphthoquinone **47** IR (KBr pellet)




2-(3-chlorophenoxy)-1,4-naphthoquinone **47** HPLC chromatogram (286 nm, 40% H₂O/MeOH)




2-(4-chlorophenoxy)-1,4-naphthoquinone **48**

48 was obtained according to general procedure I 2.1.1 and purified by hot extraction-filtration with MTBE, column chromatography (benzene as eluent, isocratically) and recrystallization with isooctane. Yield 162 mg (76%), yellow solid, mp = (131.0 – 133.0) °C. ^1H NMR (600 MHz, CDCl_3): δ (ppm) 8.23 (1H, dd, $J = 2$ Hz, $J = 7$ Hz, ArH), 8.11 (1H, dd, $J = 2$ Hz, $J = 7$ Hz, ArH), 7.81 (2H, m, ArH), 7.47 (2H, d, $J_{ortho} = 6$ Hz, ArH), 7.12 (2H, d, $J_{ortho} = 6$ Hz, ArH), 6.00 (1H, s, ArH). ^{13}C NMR (150 MHz, CDCl_3): δ (ppm) = 184.7, 179.7, 160.1, 151.2, 134.6, 133.6, 132.1, 131.9, 131.1, 130.5, 126.8, 126.3, 122.5, 113.6. IR (KBr pellet) cm^{-1} : 3049, 1684, 1654, 1625, 1482, 1238, 981. UV-Vis (MeOH) nm: 204, 221, 243 (max), 271, 330. TOF-MS: m/z $[\text{M}+\text{H}]^+$ calc for $\text{C}_{16}\text{H}_{10}\text{O}_3\text{Cl}$: 285.0318; found: 285.0321. Purity: 99.6%. MIC (μM): 439 (*S. aureus*), 439 (*C. albicans*), >3000 (*E. coli*).


2-(4-chlorophenoxy)-1,4-naphthoquinone **48** ^1H NMR




2-(4-chlorophenoxy)-1,4-naphthoquinone **48** ^{13}C NMR




2-(4-chlorophenoxy)-1,4-naphthoquinone **48** UV-Vis (MeOH)



2-(4-chlorophenoxy)-1,4-naphthoquinone **48** IR (KBr pellet)




2-(4-chlorophenoxy)-1,4-naphthoquinone **48** HPLC chromatogram (270 nm, 100% MeCN)




2-phenoxy-1,4-naphthoquinone **49**

49 was obtained according to general procedure I 2.1.1 and purified by hot extraction-filtration with MTBE, cold base wash with MTBE as solvent and NaOH 0.5 mol/L and recrystallization with isooctane. Yield 144 mg (77%), yellow-orange crystals, mp = (94.0 – 102.0) °C. ^1H NMR (600 MHz, CDCl_3): δ (ppm) = 8.20 (1H, dd, $J = 3.4$ Hz, $J = 6.8$ Hz, ArH), 8.07 (1H, dd, $J = 3.4$ Hz, $J = 6.8$ Hz, ArH), 7.76 (2H, m, ArH), 7.47 (2H, t, ArH), 7.32 (1H, tt, $J_{\text{meta}} = 1.3$ Hz, $J_{\text{ortho}} = 7.5$ Hz, ArH), 7.14 (2H, d, $J_{\text{ortho}} = 8.8$ Hz, ArH), 5.96 (1H, s, ArH). ^{13}C NMR (150 MHz, CDCl_3): δ (ppm) = 185.0, 179.9, 160.5, 152.7, 134.5, 133.5, 132.0, 131.1, 130.4, 126.8, 126.6, 126.2, 121.1, 113.4. IR (KBr pellet) cm^{-1} : 3067, 1684, 1654, 1614, 1586, 1262, 1213. UV-Vis (MeOH) nm: 210, 244 (max), 270, 331. TOF-MS: m/z $[\text{M}+\text{H}]^+$ calc for $\text{C}_{16}\text{H}_{11}\text{O}_3$: 251.0708; found: 251.0703. Purity: 98.6%. MIC (μM): 250 (*S. aureus*), 250 (*C. albicans*), >3000 (*E. coli*).


2-phenoxy-1,4-naphthoquinone **49** ^1H NMR




2-phenoxy-1,4-naphthoquinone **49** ^{13}C NMR




2-phenoxy-1,4-naphthoquinone **49** UV-Vis (MeOH)



2-phenoxy-1,4-naphthoquinone **49** IR (KBr pellet)




2-phenoxy-1,4-naphthoquinone **49** HPLC chromatogram (270 nm, 100% MeCN)




2-(2-nitrophenoxy)-1,4-naphthoquinone **50**

50 was obtained according to general procedure I 2.1.1 and purified by hot extraction-filtration with 25% n-heptane/isooctane and recrystallization with isooctane. Yield 89 mg (30%), yellow crystals, mp = (115.0 – 118.0) °C. ^1H NMR (600 MHz, CDCl_3): δ (ppm) = 8.22 (1H, m, ArH), 8.18 (1H, dd, $J = 2.6$ Hz, $J = 6.9$ Hz, ArH), 8.08 (1H, m, ArH), 7.78 (2H, m, ArH), 7.75 (1H, m, ArH), 7.51 (1H, m, ArH), 7.33 (1H, dd, $J_{\text{meta}} = 1.4$ Hz, $J_{\text{ortho}} = 8.2$ Hz, ArH), 5.95 (1H, s, ArH). ^{13}C NMR (150 MHz, CDCl_3): δ (ppm) = 184.5, 178.8, 159.4, 146.1, 141.6, 135.4, 134.6, 133.8, 131.9, 131.0, 127.4, 127.0, 126.7, 126.4, 124.4, 114.2. IR (KBr pellet) cm^{-1} : 3072, 1677, 1655, 1602, 1529, 1349, 1264, 1231. UV-Vis (MeOH) nm: 204, 218, 252 (max), 273, 334. TOF-MS: m/z $[\text{M}+\text{H}]^+$ calc for $\text{C}_{16}\text{H}_{10}\text{NO}_5$: 296.0559; found: 296.0562. Purity: non measured. MIC (μM): 13 (*S. aureus*), 85 (*C. albicans*), 1693 (*E. coli*).


2-(2-nitrophenoxy)-1,4-naphthoquinone **50** ^1H NMR




2-(2-nitrophenoxy)-1,4-naphthoquinone **50** ^{13}C NMR



2-(2-nitrophenoxy)-1,4-naphthoquinone **50** UV-Vis (MeOH)




2-(2-nitrophenoxy)-1,4-naphthoquinone **50** IR (KBr pellet)




2-(2,4,5-trichlorophenoxy)-1,4-naphthoquinone **51**

51 was obtained according to general procedure I 2.1.1 and purified by hot extraction-filtration and recrystallization, both with n-heptane. Yield 92 mg (26%), yellow needles, mp = (158.0 – 159.0) °C. ^1H NMR (600 MHz, CDCl_3): δ (ppm) 8.21 (1H, dd, $J = 2.5$ Hz, $J = 7.6$ Hz, ArH), 8.09 (1H, dd, $J = 2.5$ Hz, $J = 7.6$ Hz, ArH), 7.79 (2H, m, ArH), 7.65 (1H, s, ArH), 7.35 (1H, s, ArH), 5.93 (1H, s, ArH). ^{13}C NMR (150 MHz, CDCl_3): δ (ppm) = 184.4, 178.8, 158.1, 147.5, 134.7, 133.9, 132.4, 132.0, 131.8, 131.5, 131.0, 126.9, 126.4, 125.8, 124.4, 114.1. IR (KBr pellet) cm^{-1} : 3077, 1686, 1655, 1624, 1466, 1264, 1180, 755. UV-Vis (MeOH) nm: 217, 249 (max), 269, 333. TOF-MS: m/z $[\text{M}+\text{H}]^+$ calc for $\text{C}_{16}\text{H}_8\text{O}_3\text{Cl}_3$: 352.9539; found: 352.9547. Purity: 98.7%. MIC (μM): >2000 (*S. aureus*), >2000 (*C. albicans*), >2000 (*E. coli*).


2-(2,4,5-trichlorophenoxy)-1,4-naphthoquinone **51** ^1H NMR




2-(2,4,5-trichlorophenoxy)-1,4-naphthoquinone **51** ^{13}C NMR




2-(2,4,5-trichlorophenoxy)-1,4-naphthoquinone **51** UV-Vis (MeOH)



2-(2,4,5-trichlorophenoxy)-1,4-naphthoquinone **51 IR (KBr pellet)**




2-(2,4,5-trichlorophenoxy)-1,4-naphthoquinone **51 HPLC chromatogram (274 nm, 100% MeCN)**




2-(2,4,6-trichlorophenoxy)-1,4-naphthoquinone **52**

52 was obtained according to general procedure I 2.1.1 and purified by hot extraction-filtration with MTBE and recrystallization with isooctane. Yield 155 mg (44%), yellow needles, mp = (163.0 – 164.0) °C. ^1H NMR (600 MHz, CDCl_3): δ (ppm) 8.23 (1H, m, ArH), 8.08 (1H, m, ArH), 7.78 (2H, m, ArH), 7.46 (2H, s, ArH), 5.82 (1H, s, ArH). ^{13}C NMR (150 MHz, CDCl_3): δ (ppm) = 184.4, 178.5, 156.8, 143.9, 134.6, 133.8, 132.9, 131.9, 131.0, 129.4, 129.3, 126.9, 126.4, 113.2. IR (KBr pellet) cm^{-1} : 3070, 1684, 1654, 1618, 1444, 1260, 1243, 972. UV-Vis (MeCN) nm: 246 (max), 270, 335. TOF-MS: m/z $[\text{M}+\text{H}]^+$ calc for $\text{C}_{16}\text{H}_8\text{O}_3\text{Cl}_3$: 352.9537; found: 352.9539. Purity: 99.3%. MIC (μM): >2000 (*S. aureus*), >2000 (*C. albicans*), >2000 (*E. coli*).


2-(2,4,6-trichlorophenoxy)-1,4-naphthoquinone **52** ^1H NMR




2-(2,4,6-trichlorophenoxy)-1,4-naphthoquinone **52** ^{13}C NMR




2-(2,4,6-trichlorophenoxy)-1,4-naphthoquinone **52** UV-Vis (MeCN)



2-(2,4,6-trichlorophenoxy)-1,4-naphthoquinone 52 IR (KBr pellet)




2-(2,4,6-trichlorophenoxy)-1,4-naphthoquinone 52 HPLC chromatogram (270 nm, 100% MeCN)




2-(4-chloro-3,5-dimethylphenoxy)-1,4-naphthoquinone **53**

53 was obtained according to general procedure I 2.1.1 and purified by hot extraction-filtration with isooctane, cold base wash with NaOH 0.5 mol/L and CCl₄ as solvent, and recrystallization with isooctane. Yield 164 mg (53%), yellow-orange crystals, mp = (144.0 – 145.5) °C. ¹H NMR (600 MHz, CDCl₃): δ (ppm) = 8.20 (1H, dd, *J*_{ortho} = 7.3 Hz, ArH), 8.07 (1H, dd, *J*_{ortho} = 7.3 Hz, ArH), 7.76 (2H, m, ArH), 6.88 (2H, s, ArH), 5.98 (1H, s, ArH), 2.40 (6H, s, CH₃). ¹³C NMR (150 MHz, CDCl₃): δ (ppm) = 185.0, 179.8, 160.4, 150.3, 138.7, 134.5, 133.6, 132.4, 131.9, 131.1, 126.8, 126.2, 120.7, 113.5, 20.9. IR (KBr pellet) cm⁻¹: 3068, 2953, 2931, 1686, 1648, 1615, 1596, 1259, 1200, 777. UV-Vis (MeOH) nm: 201, 210 (max), 248, 273, 331. TOF-MS: *m/z* [M+H]⁺ calc for C₁₈H₁₄O₃Cl: 313.0631; found: 313.0631. Purity: 98.0%. MIC (μM): >3000 (*S. aureus*), >3000 (*C. albicans*), >3000 (*E. coli*).


2-(4-chloro-3,5-dimethylphenoxy)-1,4-naphthoquinone **53** ¹H NMR




2-(4-chloro-3,5-dimethylphenoxy)-1,4-naphthoquinone **53** ^{13}C NMR




2-(4-chloro-3,5-dimethylphenoxy)-1,4-naphthoquinone **53** UV-Vis (MeOH)



2-(4-chloro-3,5-dimethylphenoxy)-1,4-naphthoquinone 53 IR (KBr pellet)




2-(4-chloro-3,5-dimethylphenoxy)-1,4-naphthoquinone 53 HPLC chromatogram (270 nm, 100% MeCN)




2-(2-isopropyl-5-methylphenoxy)-1,4-naphthoquinone **54**

54 was obtained according to general procedure I 2.1.1 and purified by hot extraction-filtration with isooctane, cold base wash with NaOH 0.5 mol/L and CCl₄ as solvent, and recrystallization with isooctane. Yield 178 mg (58%), yellow-orange pellets, mp = (97.0 – 99.0) °C. ¹H NMR (600 MHz, CDCl₃): δ (ppm) = 8.21 (1H, m, ArH), 8.07 (1H, m, ArH), 7.76 (2H, m, ArH), 7.27 (1H, d, *J*_{ortho} = 7.4 Hz, ArH), 7.09 (1H, d, *J*_{ortho} = 8.4 Hz, ArH), 6.81 (1H, d, ArH), 5.93 (1H, s, ArH), 3.00 (1H, m, CH), 2.35 (3H, s, CH₃), 1.20 (6H, d, -CH(CH₃)₂). ¹³C NMR (150 MHz, CDCl₃): δ (ppm) = 185.1, 179.9, 160.6, 149.6, 137.6, 137.2, 134.4, 133.5, 132.0, 131.2, 127.8, 127.4, 126.0, 126.2, 121.5, 113.0, 26.7, 23.3, 20.8. IR (KBr pellet) cm⁻¹: 3071, 2953, 1685, 1654, 1607, 1577, 1262, 1194. UV-Vis (MeOH) nm: 218, 249 (max), 272, 332. TOF-MS: m/z [M+H]⁺ calc for C₂₀H₁₉O₃: 307.1334; found: 307.1339. Purity: 96.2%. MIC (μM): >3000 (*S. aureus*), >3000 (*C. albicans*), >3000 (*E. coli*).


2-(2-isopropyl-5-methylphenoxy)-1,4-naphthoquinone **54** ¹H NMR




2-(2-isopropyl-5-methylphenoxy)-1,4-naphthoquinone **54** ^{13}C NMR




2-(2-isopropyl-5-methylphenoxy)-1,4-naphthoquinone **54** UV-Vis (MeOH)



2-(2-isopropyl-5-methylphenoxy)-1,4-naphthoquinone **54** IR (KBr pellet)




2-(2-isopropyl-5-methylphenoxy)-1,4-naphthoquinone **54** HPLC chromatogram (286 nm, 40% H₂O/MeOH)




2-(5-chloro-2-(2,4-dichlorophenoxy)phenoxy)-1,4-naphthoquinone **55**

55 was obtained according to general procedure I 2.1.1 and purified by hot extraction-filtration with CH₂Cl₂, column chromatography (gradient elution from 100% benzene to 15% diethyl ether/benzene) and cold base wash with NaOH 0.5 mol/L and CH₂Cl₂ as solvent. Yield 178 mg (40%), ochre solid, mp = (116.0 – 119.0) °C. ¹H NMR (600 MHz, CDCl₃): δ (ppm) = 8.15 (1H, dd, *J* = 1.6 Hz, *J* = 7.1 Hz, ArH), 8.07 (1H, dd, *J* = 1.6 Hz, *J* = 7.1 Hz, ArH), 7.76 (2H, ddd, *J*_{meta} = 1.6 Hz, *J*_{ortho} = 7.9 Hz, *J*_{ortho} = 7.9 Hz, ArH), 7.36 (1H, d, *J*_{meta} = 1.9 Hz, ArH), 7.28 (1H, d, *J*_{meta} = 1.9 Hz, ArH), 7.22 (1H, dd, *J*_{meta} = 1.9 Hz, *J*_{ortho} = 8.4 Hz, ArH), 7.19 (1H, dd, *J*_{meta} = 1.9 Hz, *J*_{ortho} = 8.4 Hz, ArH), 6.91 (1H, d, *J*_{ortho} = 8.4 Hz, ArH), 6.82 (1H, d, *J*_{ortho} = 8.4 Hz, ArH), 6.07 (1H, s, ArH). ¹³C NMR (150 MHz, CDCl₃): δ (ppm) = 184.7, 179.0, 158.4, 149.6, 146.4, 142.6, 134.4, 133.6, 131.9, 131.0, 130.6, 130.5, 129.5, 128.4, 127.6, 126.8, 126.6, 126.3, 123.7, 121.4, 119.5, 114.1. IR (KBr pellet) cm⁻¹: 3067, 1684, 1655, 1619, 1595, 1472, 1260, 1218, 980. UV-Vis (MeCN) nm: 271 (max), 335. TOF-MS: *m/z* [M+H]⁺ calc for C₂₂H₁₂O₄Cl₃: 444.9801; found: 444.9804. Purity: 99.1%. MIC (μM): 0.34 (*S. aureus*), >2000 (*C. albicans*), 0.56 (*E. coli*).


2-(5-chloro-2-(2,4-dichlorophenoxy)phenoxy)-1,4-naphthoquinone **55** ¹H NMR




2-(5-chloro-2-(2,4-dichlorophenoxy)phenoxy)-1,4-naphthoquinone **55** ^{13}C NMR




2-(5-chloro-2-(2,4-dichlorophenoxy)phenoxy)-1,4-naphthoquinone **55** UV-Vis (MeCN)



2-(5-chloro-2-(2,4-dichlorophenoxy)phenoxy)-1,4-naphthoquinone 55 IR (KBr pellet)




2-(5-chloro-2-(2,4-dichlorophenoxy)phenoxy)-1,4-naphthoquinone 55 HPLC chromatogram (270 nm, 100% MeCN)




2-(4-allyl-2-methoxyphenoxy)-1,4-naphthoquinone **56**

56 was obtained according to general procedure I 2.1.1, except that 3.0 g of whole cloves (*Syzygium aromaticum*) were used to give approx. 1.25 mmol of eugenol. The purification included hot extraction-filtration with diethyl ether and recrystallization with isooctane. Yield 35 mg (11%), ochre solid, mp = (75.0 – 77.0) °C. ¹H NMR (600 MHz, CDCl₃): δ (ppm) = 8.20 (1H, d, *J* = 4.8 Hz, ArH), 8.10 (1H, d, *J* = 4.8 Hz, ArH), 7.80 (2H, m, ArH), 7.04 (1H, d, *J*_{ortho} = 4.8 Hz, ArH), 6.85 (1H, s, ArH), 6.82 (1H, dd, *J*_{ortho} = 4.8 Hz, ArH), 5.98 (1H, m, CH), 5.85 (1H, s, ArH), 5.14 (2H, d, *J* = 5.7 Hz, =CH₂), 3.78 (3H, s, CH₃), 3.41 (2H, d, *J* = 6.3 Hz, -CH₂). ¹³C NMR (150 MHz, CDCl₃): δ (ppm) = 185.3, 179.8, 159.8, 150.6, 140.0, 139.3, 136.8, 134.2, 133.4, 132.2, 131.3, 126.7, 126.1, 122.2, 121.3, 116.4, 113.4, 112.8, 55.8, 40.1. IR (KBr pellet) cm⁻¹: 3068, 2938, 1683, 1650, 1617, 1596, 1503, 1262, 1210. UV-Vis (MeOH) nm: 219, 249 (max), 271, 332. TOF-MS: *m/z* [M+H]⁺ calc for C₂₀H₁₇O₄: 321.1127; found: 321.1122. Purity: 97.1%. MIC (μM): 1561 (*S. aureus*), >3000 (*C. albicans*), >3000 (*E. coli*).


2-(4-allyl-2-methoxyphenoxy)-1,4-naphthoquinone **56** ¹H NMR




2-(4-allyl-2-methoxyphenoxy)-1,4-naphthoquinone **56** ^{13}C NMR




2-(4-allyl-2-methoxyphenoxy)-1,4-naphthoquinone **56** UV-Vis (MeOH)



2-(4-allyl-2-methoxyphenoxy)-1,4-naphthoquinone **56** IR (KBr pellet)




2-(4-allyl-2-methoxyphenoxy)-1,4-naphthoquinone **56** HPLC chromatogram (270 nm, 100% MeCN)




2-(4-[2-(4-hydroxyphenyl)propan-2-yl]phenoxy)-1,4-naphthoquinone **57**

57 was obtained according to general procedure I 2.1.1, but using 474 mg (2.0 mmol) of 2-bromo-1,4-naphthoquinone, 228 mg (1.0 mmol) of bisphenol A and 975 mg (3.0 mmol) of dried Cs_2CO_3 . The product was purified by hot extraction-filtration with MTBE and recrystallization with toluene. Yield 304 mg (56%), yellow-to-orange solid, mp = (199.0 – 201.0) °C. ^1H NMR (600 MHz, CDCl_3): δ (ppm) = 8.20 (2H, d, J = 7 Hz, ArH), 8.06 (2H, d, J = 7 Hz, ArH), 7.76 (4H, m, ArH), 7.33 (4H, d, J_{ortho} = 9 Hz, ArH), 7.07 (4H, d, J_{ortho} = 9 Hz, ArH), 6.01 (2H, s, ArH), 1.73 (6H, s, CH_3). ^{13}C NMR (150 MHz, CDCl_3): δ (ppm) = 185.1, 179.7, 160.5, 150.6, 148.6, 134.4, 133.5, 131.9, 131.1, 128.7, 126.8, 126.2, 120.7, 113.3, 42.7, 31.0. IR (KBr pellet) cm^{-1} : 3066, 2974, 1684, 1650, 1597, 1498, 1263, 1216. UV-Vis (MeCN) nm: 251 (max), 271, 334. TOF-MS: m/z $[\text{M}+\text{H}]^+$ calc for $\text{C}_{35}\text{H}_{25}\text{O}_6$: 541.1651; found: 541.1664. Purity: 95.1%. MIC (μM): 925 (*S. aureus*), >1850 (*C. albicans*), >1850 (*E. coli*).


2-(4-[2-(4-hydroxyphenyl)propan-2-yl]phenoxy)-1,4-naphthoquinone **57** ^1H NMR




2-(4-[2-(4-hydroxyphenyl)propan-2-yl]phenoxy)-1,4-naphthoquinone **57** ^{13}C NMR




2-(4-[2-(4-hydroxyphenyl)propan-2-yl]phenoxy)-1,4-naphthoquinone **57** UV-Vis (MeCN)



2-(4-[2-(4-hydroxyphenyl)propan-2-yl]phenoxy)-1,4-naphthoquinone 57 IR (KBr pellet)




2-(4-[2-(4-hydroxyphenyl)propan-2-yl]phenoxy)-1,4-naphthoquinone 57 HPLC chromatogram (270 nm, 100% MeCN)




2-(4-[2-(4-hydroxyphenyl)cyclohexyl]phenoxy)-1,4-naphthoquinone **58**

58 was obtained according to general procedure I 2.1.1, but using 237 mg (1.0 mmol) of 2-bromo-1,4-naphthoquinone, 200 mg (0.75 mmol) of bisphenol Z and 590 mg (1.6 mmol) of dried Cs₂CO₃. The product was purified by hot extraction-filtration with MTBE and recrystallization with benzene. Yield 251 mg (92%), yellow solid, mp = (248.0 – 250.0) °C. ¹H NMR (600 MHz, CDCl₃): δ (ppm) = 8.20 (2H, d, *J* = 7 Hz, ArH), 8.06 (2H, d, *J* = 7 Hz, ArH), 7.76 (4H, m, ArH), 7.36 (4H, d, *J*_{ortho} = 6 Hz; ArH), 7.07 (4H, d, *J*_{ortho} = 8 Hz, ArH), 6.01 (2H, s, ArH), 2.30 (4H, s, -C(CH₂CH₂)₂CH₂), 1.59 (2H, m, -C(CH₂CH₂)₂CH₂), 1.56 (4H, m, -C(CH₂CH₂)₂CH₂). ¹³C NMR (150 MHz, CDCl₃): δ (ppm) = 185.1, 179.9, 160.5, 150.4, 146.6, 134.4, 133.5, 132.0, 131.1, 128.3, 126.8, 126.2, 120.8, 113.3, 46.1, 37.5, 26.2, 22.8. IR (KBr pellet) cm⁻¹: 3058, 2931, 1681, 1652, 1597, 1501, 1262, 1225. UV-Vis (MeCN) nm: 270 (max), 330. TOF-MS: *m/z* [M+H]⁺ calc for C₃₈H₂₉O₆: 581.1964; found: 581.1965. Purity: 93.6%. MIC (μM): (*S. aureus*), (*C. albicans*), (*E. coli*).


2-(4-[2-(4-hydroxyphenyl)cyclohexyl]phenoxy)-1,4-naphthoquinone **58** ¹H NMR




2-(4-[2-(4-hydroxyphenyl)cyclohexyl]phenoxy)-1,4-naphthoquinone **58** ^{13}C NMR




2-(4-[2-(4-hydroxyphenyl)cyclohexyl]phenoxy)-1,4-naphthoquinone **58** UV-Vis (MeCN)



2-(4-[2-(4-hydroxyphenyl)cyclohexyl]phenoxy)-1,4-naphthoquinone **58** IR (KBr pellet)




2-(4-[2-(4-hydroxyphenyl)cyclohexyl]phenoxy)-1,4-naphthoquinone **58** HPLC chromatogram (270 nm, 100% MeCN)




2-(2-methylphenoxy)-1,4-naphthoquinone **59**

59 was obtained according to general procedure II 2.1.2 and purified by hot extraction-filtration with MTBE and column chromatography (1% triethylamine/benzene as eluent, isocratically). Yield 70 mg (27%), reddish-brown solid, mp = (79.0 – 83.0) °C. ^1H NMR (600 MHz, CDCl_3): δ (ppm) = 8.22 (1H, d, J = 6.8 Hz, ArH), 8.07 (1H, d, J = 6.8 Hz, ArH), 7.77 (2H, m, ArH), 7.31 (1H, d, J_{ortho} = 8 Hz, ArH), 7.28 (1H, td, J_{meta} = 2 Hz, J_{ortho} = 8 Hz, ArH), 7.22 (1H, td, J_{meta} = 2 Hz, J_{ortho} = 8 Hz, ArH), 7.05 (1H, d, J_{ortho} = 8 Hz, ArH), 5.83 (1H, s, ArH), 2.21 (3H, s, $-\text{CH}_3$). ^{13}C NMR (150 MHz, CDCl_3): δ (ppm) = 185.0, 179.9, 159.7, 150.8, 134.4, 133.5, 132.0, 131.2, 130.0, 127.9, 126.8, 126.8, 126.2, 121.2, 112.8, 15.8. IR (KBr pellet) cm^{-1} : 3068, 2927, 1685, 1655, 1613, 1595, 1260, 1226. UV-Vis (MeOH) nm: 213, 249 (max), 268, 332. TOF-MS: m/z $[\text{M}+\text{H}]^+$ calc for $\text{C}_{17}\text{H}_{13}\text{O}_3$: 265.0865; found: 265.0865. Purity: 96.6%.


2-(2-methylphenoxy)-1,4-naphthoquinone **59** ^1H NMR




2-(2-methylphenoxy)-1,4-naphthoquinone **59** ^{13}C NMR




2-(2-methylphenoxy)-1,4-naphthoquinone **59** UV-Vis (MeOH)



2-(2-methylphenoxy)-1,4-naphthoquinone **59** IR (KBr pellet)




2-(2-methylphenoxy)-1,4-naphthoquinone **59** HPLC chromatogram (286 nm, 40% H₂O/MeOH)




2-(3-methylphenoxy)-1,4-naphthoquinone **60**

60 was obtained according to general procedure II 2.1.2 and purified by hot extraction-filtration with MTBE and column chromatography (1% triethylamine/benzene as eluent, isocratically). Yield 181 mg (69%), yellow solid, mp = (97.0 – 102.0) °C. ^1H NMR (600 MHz, CDCl_3): δ (ppm) = 8.21 (1H, dd, $J = 2.1$ Hz, $J = 6.4$ Hz, ArH), 8.07 (1H, dd, $J = 2.1$ Hz, $J = 6.4$ Hz, ArH), 7.77 (2H, m, ArH), 7.33 (1H, t, $J_{\text{ortho}} = 6.4$ Hz, ArH), 7.12 (1H, d, $J_{\text{ortho}} = 6.4$ Hz, ArH), 6.95 (1H, d, ArH), 6.93 (2H, d, $J_{\text{ortho}} = 8.6$ Hz, ArH), 5.97 (1H, s, ArH), 2.39 (3H, s, $-\text{CH}_3$). ^{13}C NMR (150 MHz, CDCl_3): δ (ppm) = 185.1, 180.0, 160.6, 152.7, 140.8, 134.4, 133.5, 132.0, 131.2, 130.1, 127.4, 126.8, 126.2, 121.6, 118.0, 113.4, 21.3. IR (KBr pellet) cm^{-1} : 3068, 2927, 1685, 1648, 1607, 1594, 1261, 1245. UV-Vis (MeOH) nm: 212, 248 (max), 270, 333. TOF-MS: m/z $[\text{M}+\text{H}]^+$ calc for $\text{C}_{17}\text{H}_{13}\text{O}_3$: 265.0865; found: 265.0863. Purity: 98.7%. MIC (μM): 473 (*S. aureus*), 237 (*C. albicans*), >3000 (*E. coli*).


2-(3-methylphenoxy)-1,4-naphthoquinone **60** ^1H NMR




2-(3-methylphenoxy)-1,4-naphthoquinone **60** ^{13}C NMR




2-(3-methylphenoxy)-1,4-naphthoquinone **60** UV-Vis (MeOH)



2-(3-methylphenoxy)-1,4-naphthoquinone **60** IR (KBr pellet)




2-(3-methylphenoxy)-1,4-naphthoquinone **60** HPLC chromatogram (274 nm, 100% MeCN)




2-(4-methylphenoxy)-1,4-naphthoquinone **61**

61 was obtained according to general procedure II 2.1.2 and purified by hot extraction-filtration with MTBE and column chromatography (1% triethylamine/benzene as eluent, isocratically). Yield 155 mg (59%), yellow solid, mp = (100.0 – 102.0) °C. ^1H NMR (600 MHz, CDCl_3): δ (ppm) = 8.20 (1H, d, J = 5 Hz, ArH), 8.06 (1H, d, J = 5 Hz, ArH), 7.76 (2H, m, ArH), 7.25 (2H, d, J_{ortho} = 7.5 Hz, ArH), 7.01 (2H, d, J_{ortho} = 7.5 Hz, ArH), 5.98 (1H, s, ArH), 2.40 (3H, s, $-\text{CH}_3$). ^{13}C NMR (150 MHz, CDCl_3): δ (ppm) = 185.4, 180.0, 160.7, 150.4, 136.4, 134.4, 133.5, 132.0, 131.2, 130.8, 126.8, 126.2, 120.8, 113.2, 20.9. IR (KBr pellet) cm^{-1} : 3072, 2922, 2865, 1679, 1651, 1618, 1506, 1261, 1231. UV-Vis (MeOH) nm: 205, 249 (max), 271, 330. TOF-MS: m/z $[\text{M}+\text{H}]^+$ calc for $\text{C}_{17}\text{H}_{13}\text{O}_3$: 265.0865; found: 265.0866. Purity: 97.9%. MIC (μM): 237 (*S. aureus*), 473 (*C. albicans*), >3000 (*E. coli*).


2-(4-methylphenoxy)-1,4-naphthoquinone **61** ^1H NMR




2-(4-methylphenoxy)-1,4-naphthoquinone **61** ^{13}C NMR




2-(4-methylphenoxy)-1,4-naphthoquinone **61** UV-Vis (MeOH)



2-(4-methylphenoxy)-1,4-naphthoquinone **61 IR (KBr pellet)**




2-(4-methylphenoxy)-1,4-naphthoquinone **61 HPLC chromatogram (286 nm, 40% H₂O/MeOH)**




2-(4-nitrophenoxy)-1,4-naphthoquinone **62**

62 was obtained according to general procedure III 2.1.3 and purified by cold base wash (MTBE as solvent and K_2CO_3 sat as base), column chromatography (benzene as eluent, isocratically) and recrystallization with isooctane. Yield 71 mg (24%), brown to yellow crystals, mp = (190.0 – 192.0) °C. 1H NMR (600 MHz, $CDCl_3$): δ (ppm) = 8.36 (2H, d, $J_{ortho} = 7.5$ Hz, ArH), 8.20 (1H, d, $J_{8-7} = 7.5$ Hz, ArH), 8.11 (1H, d, $J_{5-6} = 7.5$ Hz, ArH), 7.81 (2H, m, ArH), 7.30 (2H, d, $J_{ortho} = 7.5$ Hz, ArH), 6.18 (1H, s, ArH). ^{13}C NMR (150 MHz, $CDCl_3$): δ (ppm) = 184.4, 179.2, 158.6, 158.1, 145.4, 134.8, 133.9, 131.7, 130.9, 126.9, 126.4, 126.2, 121.1, 116.2. IR (KBr pellet) cm^{-1} : 3050, 1686, 1646, 1616, 1587, 1514, 1340, 1229, 886. UV-Vis (MeOH) nm: 203, 250, 269 (max). TOF-MS: m/z $[M+H]^+$ calc for $C_{16}H_{10}NO_5$: 296.0559; found: 296.0592. Purity: 97.1%. MIC (μM): 212 (*S. aureus*), 3387 (*C. albicans*), >3000 (*E. coli*).


2-(4-nitrophenoxy)-1,4-naphthoquinone **62** 1H NMR




2-(4-nitrophenoxy)-1,4-naphthoquinone **62** ^{13}C NMR




2-(4-nitrophenoxy)-1,4-naphthoquinone **62** UV-Vis (MeOH)



2-(4-nitrophenoxy)-1,4-naphthoquinone **62** IR (KBr pellet)




2-(4-nitrophenoxy)-1,4-naphthoquinone **62** HPLC chromatogram (270 nm, 100% MeCN)




2-(2,4-dichlorophenoxy)-1,4-naphthoquinone **63**

63 was obtained according to general procedure III 2.1.3 and purified by hot extraction-filtration and recrystallization, both with isooctane. Yield 177 mg (56%), yellow needle crystals, mp = (127.0 – 130.0) °C. ^1H NMR (600 MHz, CDCl_3): δ (ppm) = 8.21 (1H, dd, $J = 2.5$ Hz, $J = 6.7$ Hz, ArH), 8.08 (1H, dd, $J = 2.5$ Hz, $J = 6.7$ Hz, ArH), 7.78 (2H, m, ArH), 7.54 (1H, d, $J_{\text{meta}} = 2.3$ Hz, ArH), 7.35 (1H, dd, $J_{\text{meta}} = 3.0$ Hz, $J_{\text{ortho}} = 8.3$ Hz, ArH), 7.16 (1H, d, $J_{\text{ortho}} = 8.3$ Hz, ArH), 5.84 (1H, s, ArH). ^{13}C NMR (150 MHz, CDCl_3): δ (ppm) = 184.6, 179.1, 158.5, 147.2, 134.6, 133.7, 132.9, 131.9, 131.1, 131.0, 128.9, 127.6, 126.9, 126.4, 123.9, 113.6. IR (KBr pellet) cm^{-1} : 3094, 1685, 1653, 1613, 1474, 1262, 1242, 982. UV-Vis (MeOH) nm: 219, 249(max), 268, 333. TOF-MS: m/z $[\text{M}+\text{H}]^+$ calc for $\text{C}_{16}\text{H}_9\text{O}_3\text{Cl}_2$: 318.9929; found: 318.9928. Purity: 98.3%. MIC (μM): 49 (*S. aureus*), >3000 (*C. albicans*), >3000 (*E. coli*).


2-(2,4-dichlorophenoxy)-1,4-naphthoquinone **63** ^1H NMR




2-(2,4-dichlorophenoxy)-1,4-naphthoquinone **63** ^{13}C NMR




2-(2,4-dichlorophenoxy)-1,4-naphthoquinone **63** UV-Vis (MeOH)



2-(2,4-dichlorophenoxy)-1,4-naphthoquinone 63 IR (KBr pellet)



2-(2,4-dichlorophenoxy)-1,4-naphthoquinone 63 HPLC chromatogram (270 nm, 100% MeCN)



In order to avoid security-related warning messages when switching to secured connection, you may want either to:

- confirm the exception on the next page, or
- import our [CA key](#) in your web browser

Click [here](#) to proceed.

[Click2Drug](#)

[SwissDock](#)

[SwissParam](#)

[SwissSidechain](#)

[SwissBioisostere](#)

[SwissTargetPrediction](#)

[SwissADME](#)

[SwissSimilarity](#)

[About us](#)

[Swiss Institute of Bioinformatics](#)

SwissADME

- [Disclaimer](#)
- [Help](#)
- [FAQ](#)
- [Home](#)


This website allows you to compute physicochemical descriptors as well as to predict ADME parameters, pharmacokinetic properties, druglike nature and medicinal chemistry friendliness of one or multiple small molecules to support drug discovery.

The main article describing the web service and its underlying methodologies is [SwissADME: a free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules. *Sci. Rep.* \(2017\) 7:42717.](#)

For details about development and validation of iLOG, please refer to this article: [iLOGP: a simple, robust, and efficient description of *n*-octanol/water partition coefficient for drug design using the GB/SA approach. *J. Chem. Inf. Model.* \(2014\) 54\(12\):3284-3301.](#)

For details about development and validation of the BOILED-Egg, please refer to this article: [A BOILED-Egg to predict gastrointestinal absorption and brain penetration of small molecules. *ChemMedChem* \(2016\) 11\(11\):1117-1121.](#)

Developed and maintained by the [Molecular Modeling Group](#) of the SIB | Swiss Institute of Bioinformatics.



Enter a list of SMILES here:

```
Brc3ccc(OC2=CC(=O)c1ccccc1C2=O)c(Br)c3
```

Fill with an example

Clear



Run!

Show BOILED-Egg

Retrieve data:


 Powered by
ChemAxon

Molecule 1



SMILES
S Brc1ccc(c(c1)Br)OC1=CC(=O)c2c(C1=O)cccc2

Physicochemical Properties

Formula	C ₁₆ H ₈ Br ₂ O ₃
Molecular weight	408.04 g/mol
Num. heavy atoms	21
Num. arom. heavy atoms	12
Fraction Csp ³	0.00
Num. rotatable bonds	2
Num. H-bond acceptors	3
Num. H-bond donors	0
Molar Refractivity	85.66
TPSA	

Water Solubility

Log *S* (ESOL)

ESOL: Topological method implemented from
Delaney JS. 2004 J. Chem. Inf. Model.

-5.36

Solubility
Class

1.77e-03 mg/ml ; 4.33e-06 mol/l

Solubility class: Log *S* scale

Insoluble < -10 < Poorly Moderately soluble
< -6 < Moderately < -4
< Soluble < -2 Very < 0
< Highly

Log *S* (Ali)

Ali: Topological method implemented from
Ali J. et al. 2012 J. Chem. Inf. Model.

-4.91

Solubility
Class

4.98e-03 mg/ml ; 1.22e-05 mol/l

Solubility class: Log *S*

Topological Polar Surface Area: Calculated from Ertl P. et al. 2000 J. Med. Chem.	43.37 Å ²	<u>scale</u> Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Moderately soluble
Log <i>P</i>_{o/w} (iLOGP) ⓘ	Lipophilicity	Log <i>S</i> (SILICOS-IT) ⓘ	
iLOGP: in-house physics-based method implemented from Daina A et al. 2014 J. Chem. Inf. Model.	2.89	SILICOS-IT: Fragmental method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos-it.com	-6.91
Log <i>P</i>_{o/w} (XLOGP3) ⓘ		Solubility Class ⓘ	5.01e-05 mg/ml ; 1.23e-07 mol/l
XLOGP3: Atomistic and knowledge-based method calculated by XLOGP program, version 3.2.2, courtesy of CCBG, Shanghai Institute of Organic Chemistry	4.29	Solubility class: Log <i>S</i> scale Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Poorly soluble
Log <i>P</i>_{o/w} (WLOGP) ⓘ			Pharmacokinetics
WLOGP: Atomistic method implemented from Wildman SA and Crippen GM. 1999 J. Chem. Inf. Model.	4.55	GI absorption ⓘ Gastrointestinal absorption: according to the white of the BOILED-Egg	High
Log <i>P</i>_{o/w} (MLOGP) ⓘ		BBB permeant ⓘ	
MLOGP: Topological method implemented from Moriguchi I. et al. 1992 Chem. Pharm. Bull. Moriguchi I. et al. 1994 Chem. Pharm. Bull. Lipinski PA. et al. 2001 Adv. Drug. Deliv. Rev.	2.93	BBB permeation: according to the yolk of the BOILED-Egg	Yes
Log <i>P</i>_{o/w} (SILICOS-IT) ⓘ		P-gp substrate ⓘ	
SILICOS-IT: Hybrid fragmental/topological method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos-it.com	4.68	P-glycoprotein substrate: SVM model built on 1033 molecules (training set) and tested on 415 molecules (test set) 10-fold CV: ACC=0.72 / AUC=0.77 External: ACC=0.88 / AUC=0.94	No
Consensus Log <i>P</i>_{o/w} ⓘ		CYP1A2 inhibitor ⓘ	
Consensus Log <i>P</i>_{o/w}:		Cytochrome P450 1A2 inhibitor: SVM model built on 9145 molecules (training set) and tested on 3000 molecules (test set)	Yes

Average of all five predictions

3.87

10-fold CV: ACC=0.83 / AUC=0.90
External: ACC=0.84 / AUC=0.91

CYP2C19 inhibitor ⓘ

Cytochrome P450 2C19 inhibitor: SVM model built on 9272 molecules (training set) and tested on 3000 molecules (test set) Yes
10-fold CV: ACC=0.80 / AUC=0.86
External: ACC=0.80 / AUC=0.87

CYP2C9 inhibitor ⓘ

Cytochrome P450 2C9 inhibitor: SVM model built on 5940 molecules (training set) and tested on 2075 molecules (test set) Yes
10-fold CV: ACC=0.78 / AUC=0.85
External: ACC=0.71 / AUC=0.81

CYP2D6 inhibitor ⓘ

Cytochrome P450 2D6 inhibitor: SVM model built on 3664 molecules (training set) and tested on 1068 molecules (test set) No
10-fold CV: ACC=0.79 / AUC=0.85
External: ACC=0.81 / AUC=0.87

CYP3A4 inhibitor ⓘ

Cytochrome P450 3A4 inhibitor: SVM model built on 7518 molecules (training set) and tested on 2579 molecules (test set) No
10-fold CV: ACC=0.77 / AUC=0.85
External: ACC=0.78 / AUC=0.86

Log K_p (skin permeation) ⓘ

Skin permeation: QSPR model implemented from -5.74 cm/s

Potts RO and Guy RH.
1992 Pharm. Res.

Druglikeness

Lipinski ⓘ

Lipinski (Pfizer) filter:
implemented from

Lipinski CA. et al. 2001

Adv. Drug Deliv. Rev.

Yes; 0 violation

MW $\leq$ 500

MLOGP $\leq$ 4.15

N or O $\leq$ 10

NH or OH $\leq$ 5

Ghose ⓘ

Ghose filter:

implemented from

Ghose AK. et al. 1999 J.

Comb. Chem.

Yes

160 $\leq$ MW $\leq$ 480

-0.4 $\leq$ WLOGP $\leq$ 5.6

40 $\leq$ MR $\leq$ 130

20 $\leq$ atoms $\leq$ 70

Veber ⓘ

Veber (GSK) filter:

implemented from

Veber DF. et al. 2002 J.

Yes

Med. Chem.

Rotatable bonds $\leq$ 10

TPSA $\leq$ 140

Egan ⓘ

Egan (Pharmacia)

filter: implemented

from

Egan WJ. et al. 2000 J.

Yes

Med. Chem.

WLOGP $\leq$ 5.88

TPSA $\leq$ 131.6

Muegge ⓘ

Muegge (Bayer) filter:

implemented from

Muegge I. et al. 2001 J.

Med. Chem.

200 $\leq$ MW $\leq$ 600

-2 $\leq$ XLOGP $\leq$ 5

TPSA $\leq$ 150

Yes

Num. rings $\leq$ 7

Num. carbon $>$ 4

Num. heteroatoms $>$ 1

Num. rotatable bonds $\leq$

15

H-bond acc. $\leq$ 10

H-bond don. $\leq$ 5

In order to avoid security-related warning messages when switching to secured connection, you may want either to:

- confirm the exception on the next page, or
- import our [CA key](#) in your web browser

Click [here](#) to proceed.

[Click2Drug](#)

[SwissDock](#)

[SwissParam](#)

[SwissSidechain](#)

[SwissBioisostere](#)

[SwissTargetPrediction](#)

[SwissADME](#)

[SwissSimilarity](#)

[About us](#)

[Swiss Institute of Bioinformatics](#)

SwissADME

- [Disclaimer](#)
- [Help](#)
- [FAQ](#)
- [Home](#)


This website allows you to compute physicochemical descriptors as well as to predict ADME parameters, pharmacokinetic properties, druglike nature and medicinal chemistry friendliness of one or multiple small molecules to support drug discovery.

The main article describing the web service and its underlying methodologies is [SwissADME: a free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules. *Sci. Rep.* \(2017\) 7:42717.](#)

For details about development and validation of iLOG, please refer to this article: [iLOGP: a simple, robust, and efficient description of *n*-octanol/water partition coefficient for drug design using the GB/SA approach. *J. Chem. Inf. Model.* \(2014\) 54\(12\):3284-3301.](#)

For details about development and validation of the BOILED-Egg, please refer to this article: [A BOILED-Egg to predict gastrointestinal absorption and brain penetration of small molecules. *ChemMedChem* \(2016\) 11\(11\):1117-1121.](#)

Developed and maintained by the [Molecular Modeling Group](#) of the SIB | Swiss Institute of Bioinformatics.





Enter a list of SMILES here:

Clc3ccc(OC2=CC(=O)c1ccccc1C2=O)c(Cl)c3

Retrieve data:


 Powered by
ChemAxon

Molecule 1


 SMILES
S Clc1ccc(c(c1)Cl)OC1=CC(=O)c2c(C1=O)cccc2

Physicochemical Properties

Formula	C16H8Cl2O3
Molecular weight	319.14 g/mol
Num. heavy atoms	21
Num. arom. heavy atoms	12
Fraction Csp3	0.00
Num. rotatable bonds	2
Num. H-bond acceptors	3
Num. H-bond donors	0
Molar Refractivity	80.28
TPSA	

Water Solubility

Log S (ESOL)

ESOL: Topological method implemented from
Delaney JS. 2004 J. Chem. Inf. Model.

-4.73

 Solubility
Class

5.94e-03 mg/ml ; 1.86e-05 mol/l

Solubility class: Log S scale

Insoluble < -10 < Poorly Moderately soluble
< -6 < Moderately < -4
< Soluble < -2 Very < 0
< Highly

Log S (Ali)

Ali: Topological method implemented from
Ali J. et al. 2012 J. Chem. Inf. Model.

-4.78

 Solubility
Class

5.31e-03 mg/ml ; 1.66e-05 mol/l

Solubility class: Log S

In order to avoid security-related warning messages when switching to secured connection, you may want either to:

- confirm the exception on the next page, or
- import our [CA key](#) in your web browser

Click [here](#) to proceed.

[Click2Drug](#)

[SwissDock](#)

[SwissParam](#)

[SwissSidechain](#)

[SwissBioisostere](#)

[SwissTargetPrediction](#)

[SwissADME](#)

[SwissSimilarity](#)

[About us](#)

[Swiss Institute of Bioinformatics](#)

SwissADME

- [Disclaimer](#)
- [Help](#)
- [FAQ](#)
- [Home](#)

This website allows you to compute physicochemical descriptors as well as to predict ADME parameters, pharmacokinetic properties, druglike nature and medicinal chemistry friendliness of one or multiple small molecules to support drug discovery.

The main article describing the web service and its underlying methodologies is [SwissADME: a free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules. *Sci. Rep.* \(2017\) 7:42717.](#)

For details about development and validation of iLOG, please refer to this article: [iLOGP: a simple, robust, and efficient description of *n*-octanol/water partition coefficient for drug design using the GB/SA approach. *J. Chem. Inf. Model.* \(2014\) 54\(12\):3284-3301.](#)

For details about development and validation of the BOILED-Egg, please refer to this article: [A BOILED-Egg to predict gastrointestinal absorption and brain penetration of small molecules. *ChemMedChem* \(2016\) 11\(11\):1117-1121.](#)

Developed and maintained by the [Molecular Modeling Group](#) of the SIB | Swiss Institute of Bioinformatics.

In order to avoid security-related warning messages when switching to secured connection, you may want either to:

- confirm the exception on the next page, or
- import our [CA key](#) in your web browser

Click [here](#) to proceed.

[Click2Drug](#)

[SwissDock](#)

[SwissParam](#)

[SwissSidechain](#)

[SwissBioisostere](#)

[SwissTargetPrediction](#)

[SwissADME](#)

[SwissSimilarity](#)

[About us](#)

[Swiss Institute of Bioinformatics](#)

SwissADME

- [Disclaimer](#)
- [Help](#)
- [FAQ](#)
- [Home](#)


This website allows you to compute physicochemical descriptors as well as to predict ADME parameters, pharmacokinetic properties, druglike nature and medicinal chemistry friendliness of one or multiple small molecules to support drug discovery.

The main article describing the web service and its underlying methodologies is [SwissADME: a free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules. *Sci. Rep.* \(2017\) 7:42717.](#)

For details about development and validation of iLOG, please refer to this article: [iLOGP: a simple, robust, and efficient description of *n*-octanol/water partition coefficient for drug design using the GB/SA approach. *J. Chem. Inf. Model.* \(2014\) 54\(12\):3284-3301.](#)

For details about development and validation of the BOILED-Egg, please refer to this article: [A BOILED-Egg to predict gastrointestinal absorption and brain penetration of small molecules. *ChemMedChem* \(2016\) 11\(11\):1117-1121.](#)

Developed and maintained by the [Molecular Modeling Group](#) of the SIB | Swiss Institute of Bioinformatics.





Enter a list of SMILES here:

```
BrC3cc(Br)cc(Br)c3OC2=CC(=O)c1ccccc1C2=O
```


Retrieve data:


 Powered by
ChemAxon

Molecule 1



SMILES
S BrC1cc(Br)c(c(c1)Br)OC1=CC(=O)c2c(C1=O)cccc2

Physicochemical Properties

Formula	C16H7Br3O3
Molecular weight	486.94 g/mol
Num. heavy atoms	22
Num. arom. heavy atoms	12
Fraction Csp3	0.00
Num. rotatable bonds	2
Num. H-bond acceptors	3
Num. H-bond donors	0
Molar Refractivity	93.36
TPSA	

Water Solubility

Log S (ESOL)

ESOL: Topological method implemented from
Delaney JS. 2004 J. Chem. Inf. Model.

-6.27

Solubility
Class

2.63e-04 mg/ml ; 5.39e-07 mol/l

Solubility class: Log S scale

Insoluble < -10 < Poorly Poorly soluble
< -6 < Moderately < -4
< Soluble < -2 Very < 0
< Highly

Log S (Ali)

Ali: Topological method implemented from
Ali J. et al. 2012 J. Chem. Inf. Model.

-5.63

Solubility
Class

1.14e-03 mg/ml ; 2.35e-06 mol/l

Solubility class: Log S

In order to avoid security-related warning messages when switching to secured connection, you may want either to:

- confirm the exception on the next page, or
- import our [CA key](#) in your web browser

Click [here](#) to proceed.

[Click2Drug](#)

[SwissDock](#)

[SwissParam](#)

[SwissSidechain](#)

[SwissBioisostere](#)

[SwissTargetPrediction](#)

[SwissADME](#)

[SwissSimilarity](#)

[About us](#)

[Swiss Institute of Bioinformatics](#)

SwissADME

- [Disclaimer](#)
- [Help](#)
- [FAQ](#)
- [Home](#)

This website allows you to compute physicochemical descriptors as well as to predict ADME parameters, pharmacokinetic properties, druglike nature and medicinal chemistry friendliness of one or multiple small molecules to support drug discovery.

The main article describing the web service and its underlying methodologies is [SwissADME: a free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules. *Sci. Rep.* \(2017\) 7:42717.](#)

For details about development and validation of iLOG, please refer to this article: [iLOGP: a simple, robust, and efficient description of *n*-octanol/water partition coefficient for drug design using the GB/SA approach. *J. Chem. Inf. Model.* \(2014\) 54\(12\):3284-3301.](#)

For details about development and validation of the BOILED-Egg, please refer to this article: [A BOILED-Egg to predict gastrointestinal absorption and brain penetration of small molecules. *ChemMedChem* \(2016\) 11\(11\):1117-1121.](#)

Developed and maintained by the [Molecular Modeling Group](#) of the SIB | Swiss Institute of Bioinformatics.

In order to avoid security-related warning messages when switching to secured connection, you may want either to:

- confirm the exception on the next page, or
- import our [CA key](#) in your web browser

Click [here](#) to proceed.

[Click2Drug](#)

[SwissDock](#)

[SwissParam](#)

[SwissSidechain](#)

[SwissBioisostere](#)

[SwissTargetPrediction](#)

[SwissADME](#)

[SwissSimilarity](#)

[About us](#)

[Swiss Institute of Bioinformatics](#)

SwissADME

- [Disclaimer](#)
- [Help](#)
- [FAQ](#)
- [Home](#)


This website allows you to compute physicochemical descriptors as well as to predict ADME parameters, pharmacokinetic properties, druglike nature and medicinal chemistry friendliness of one or multiple small molecules to support drug discovery.

The main article describing the web service and its underlying methodologies is [SwissADME: a free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules. *Sci. Rep.* \(2017\) 7:42717.](#)

For details about development and validation of iLOG, please refer to this article: [iLOGP: a simple, robust, and efficient description of *n*-octanol/water partition coefficient for drug design using the GB/SA approach. *J. Chem. Inf. Model.* \(2014\) 54\(12\):3284-3301.](#)

For details about development and validation of the BOILED-Egg, please refer to this article: [A BOILED-Egg to predict gastrointestinal absorption and brain penetration of small molecules. *ChemMedChem* \(2016\) 11\(11\):1117-1121.](#)

Developed and maintained by the [Molecular Modeling Group](#) of the SIB | Swiss Institute of Bioinformatics.





Enter a list of SMILES here:

Clc1ccc(cc1Cl)OC3=CC(=O)c2ccccc2C3=O

Retrieve data:


 Powered by
ChemAxon

Molecule 1


 SMILES
O=C1C(=CC(=O)c2c1cccc2)Oc1ccc(c(c1)Cl)Cl

Physicochemical Properties

Formula	C16H8Cl2O3
Molecular weight	319.14 g/mol
Num. heavy atoms	21
Num. arom. heavy atoms	12
Fraction Csp3	0.00
Num. rotatable bonds	2
Num. H-bond acceptors	3
Num. H-bond donors	0
Molar Refractivity	80.28
TPSA	

Water Solubility

Log S (ESOL)

ESOL: Topological method implemented from
Delaney JS. 2004 J. Chem. Inf. Model.

-4.73

 Solubility
 Class

5.94e-03 mg/ml ; 1.86e-05 mol/l

Solubility class: Log S scale

Insoluble < -10 < Poorly Moderately soluble
< -6 < Moderately < -4
< Soluble < -2 Very < 0
< Highly

Log S (Ali)

Ali: Topological method implemented from
Ali J. et al. 2012 J. Chem. Inf. Model.

-4.78

 Solubility
 Class

5.31e-03 mg/ml ; 1.66e-05 mol/l

Solubility class: Log S

In order to avoid security-related warning messages when switching to secured connection, you may want either to:

- confirm the exception on the next page, or
- import our [CA key](#) in your web browser

Click [here](#) to proceed.

[Click2Drug](#)

[SwissDock](#)

[SwissParam](#)

[SwissSidechain](#)

[SwissBioisostere](#)

[SwissTargetPrediction](#)

[SwissADME](#)

[SwissSimilarity](#)

[About us](#)

[Swiss Institute of Bioinformatics](#)

SwissADME

- [Disclaimer](#)
- [Help](#)
- [FAQ](#)
- [Home](#)

This website allows you to compute physicochemical descriptors as well as to predict ADME parameters, pharmacokinetic properties, druglike nature and medicinal chemistry friendliness of one or multiple small molecules to support drug discovery.

The main article describing the web service and its underlying methodologies is [SwissADME: a free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules](#). *Sci. Rep.* (2017) 7:42717.

For details about development and validation of iLOG, please refer to this article: [iLOGP: a simple, robust, and efficient description of *n*-octanol/water partition coefficient for drug design using the GB/SA approach](#). *J. Chem. Inf. Model.* (2014) 54(12):3284-3301.

For details about development and validation of the BOILED-Egg, please refer to this article: [A BOILED-Egg to predict gastrointestinal absorption and brain penetration of small molecules](#). *ChemMedChem* (2016) 11(11):1117-1121.

Developed and maintained by the [Molecular Modeling Group](#) of the SIB | Swiss Institute of Bioinformatics.

In order to avoid security-related warning messages when switching to secured connection, you may want either to:

- confirm the exception on the next page, or
- import our [CA key](#) in your web browser

Click [here](#) to proceed.

[Click2Drug](#)

[SwissDock](#)

[SwissParam](#)

[SwissSidechain](#)

[SwissBioisostere](#)

[SwissTargetPrediction](#)

[SwissADME](#)

[SwissSimilarity](#)

[About us](#)

[Swiss Institute of Bioinformatics](#)

SwissADME

- [Disclaimer](#)
- [Help](#)
- [FAQ](#)
- [Home](#)


This website allows you to compute physicochemical descriptors as well as to predict ADME parameters, pharmacokinetic properties, druglike nature and medicinal chemistry friendliness of one or multiple small molecules to support drug discovery.

The main article describing the web service and its underlying methodologies is [SwissADME: a free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules](#). *Sci. Rep.* (2017) 7:42717.

For details about development and validation of iLOG, please refer to this article: [iLOGP: a simple, robust, and efficient description of *n*-octanol/water partition coefficient for drug design using the GB/SA approach](#). *J. Chem. Inf. Model.* (2014) 54(12):3284-3301.

For details about development and validation of the BOILED-Egg, please refer to this article: [A BOILED-Egg to predict gastrointestinal absorption and brain penetration of small molecules](#). *ChemMedChem* (2016) 11(11):1117-1121.

Developed and maintained by the [Molecular Modeling Group](#) of the SIB | Swiss Institute of Bioinformatics.





Enter a list of SMILES here:

Retrieve data:


 Powered by
ChemAxon

Molecule 1


 SMILES
S C=CCc1ccc(c(c1)OC)OC1=CC(=O)c2c(C1=O)cccc2

Physicochemical Properties

Formula	C20H16O4
Molecular weight	320.34 g/mol
Num. heavy atoms	24
Num. arom. heavy atoms	12
Fraction Csp3	0.10
Num. rotatable bonds	5
Num. H-bond acceptors	4
Num. H-bond donors	0
Molar Refractivity	90.85
TPSA	

Water Solubility

Log S (ESOL)

ESOL: Topological method implemented from
Delaney JS. 2004 J. Chem. Inf. Model.

-4.53

 Solubility
Class

9.43e-03 mg/ml ; 2.94e-05 mol/l

Solubility class: Log S scale

Insoluble < -10 < Poorly Moderately soluble
< -6 < Moderately < -4
< Soluble < -2 Very < 0
< Highly

Log S (Ali)

Ali: Topological method implemented from
Ali J. et al. 2012 J. Chem. Inf. Model.

-5.05

 Solubility
Class

2.89e-03 mg/ml ; 9.01e-06 mol/l

Solubility class: Log S

In order to avoid security-related warning messages when switching to secured connection, you may want either to:

- confirm the exception on the next page, or
- import our [CA key](#) in your web browser

Click [here](#) to proceed.

[Click2Drug](#)

[SwissDock](#)

[SwissParam](#)

[SwissSidechain](#)

[SwissBioisostere](#)

[SwissTargetPrediction](#)

[SwissADME](#)

[SwissSimilarity](#)

[About us](#)

[Swiss Institute of Bioinformatics](#)

SwissADME

- [Disclaimer](#)
- [Help](#)
- [FAQ](#)
- [Home](#)

This website allows you to compute physicochemical descriptors as well as to predict ADME parameters, pharmacokinetic properties, druglike nature and medicinal chemistry friendliness of one or multiple small molecules to support drug discovery.

The main article describing the web service and its underlying methodologies is [SwissADME: a free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules](#). *Sci. Rep.* (2017) 7:42717.

For details about development and validation of iLOG, please refer to this article: [iLOGP: a simple, robust, and efficient description of *n*-octanol/water partition coefficient for drug design using the GB/SA approach](#). *J. Chem. Inf. Model.* (2014) 54(12):3284-3301.

For details about development and validation of the BOILED-Egg, please refer to this article: [A BOILED-Egg to predict gastrointestinal absorption and brain penetration of small molecules](#). *ChemMedChem* (2016) 11(11):1117-1121.

Developed and maintained by the [Molecular Modeling Group](#) of the SIB | Swiss Institute of Bioinformatics.

In order to avoid security-related warning messages when switching to secured connection, you may want either to:

- confirm the exception on the next page, or
- import our [CA key](#) in your web browser

Click [here](#) to proceed.

[Click2Drug](#)

[SwissDock](#)

[SwissParam](#)

[SwissSidechain](#)

[SwissBioisostere](#)

[SwissTargetPrediction](#)

[SwissADME](#)

[SwissSimilarity](#)

[About us](#)

[Swiss Institute of Bioinformatics](#)

SwissADME

- [Disclaimer](#)
- [Help](#)
- [FAQ](#)
- [Home](#)


This website allows you to compute physicochemical descriptors as well as to predict ADME parameters, pharmacokinetic properties, druglike nature and medicinal chemistry friendliness of one or multiple small molecules to support drug discovery.

The main article describing the web service and its underlying methodologies is [SwissADME: a free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules](#). *Sci. Rep.* (2017) 7:42717.

For details about development and validation of iLOG, please refer to this article: [iLOGP: a simple, robust, and efficient description of *n*-octanol/water partition coefficient for drug design using the GB/SA approach](#). *J. Chem. Inf. Model.* (2014) 54(12):3284-3301.

For details about development and validation of the BOILED-Egg, please refer to this article: [A BOILED-Egg to predict gastrointestinal absorption and brain penetration of small molecules](#). *ChemMedChem* (2016) 11(11):1117-1121.

Developed and maintained by the [Molecular Modeling Group](#) of the SIB | Swiss Institute of Bioinformatics.


[Show BOILED-Egg](#)



Enter a list of SMILES here:

Cc1cc(ccc1)OC3=CC(=O)c2ccccc2C3=O
[Fill with an example](#)
[Clear](#)
[Run!](#)

Retrieve data:


 Powered by
 ChemAxon

Molecule 1


 SMILES
Cc1ccccc1OC1=CC(=O)c2c(C1=O)cccc2

Physicochemical Properties

Formula	C ₁₇ H ₁₂ O ₃
Molecular weight	264.28 g/mol
Num. heavy atoms	20
Num. arom. heavy atoms	12
Fraction Csp ³	0.06
Num. rotatable bonds	2
Num. H-bond acceptors	3
Num. H-bond donors	0
Molar Refractivity	75.22
TPSA	

Water Solubility

Log *S* (ESOL)

ESOL: Topological method implemented from Delaney JS. 2004 J. Chem. Inf. Model.

-3.85

 Solubility
 Class

3.73e-02 mg/ml ; 1.41e-04 mol/l

Solubility class: Log *S* scale

Insoluble < -10 < Poorly Soluble
 < -6 < Moderately < -4
 < Soluble < -2 Very < 0
 < Highly

Log *S* (Ali)

Ali: Topological method implemented from Ali J. et al. 2012 J. Chem. Inf. Model.

-3.86

 Solubility
 Class

3.69e-02 mg/ml ; 1.40e-04 mol/l

Solubility class: Log *S*

In order to avoid security-related warning messages when switching to secured connection, you may want either to:

- confirm the exception on the next page, or
- import our [CA key](#) in your web browser

Click [here](#) to proceed.

[Click2Drug](#)

[SwissDock](#)

[SwissParam](#)

[SwissSidechain](#)

[SwissBioisostere](#)

[SwissTargetPrediction](#)

[SwissADME](#)

[SwissSimilarity](#)

[About us](#)

[Swiss Institute of Bioinformatics](#)

SwissADME

- [Disclaimer](#)
- [Help](#)
- [FAQ](#)
- [Home](#)

This website allows you to compute physicochemical descriptors as well as to predict ADME parameters, pharmacokinetic properties, druglike nature and medicinal chemistry friendliness of one or multiple small molecules to support drug discovery.

The main article describing the web service and its underlying methodologies is [SwissADME: a free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules. *Sci. Rep.* \(2017\) 7:42717.](#)

For details about development and validation of iLOG, please refer to this article: [iLOGP: a simple, robust, and efficient description of *n*-octanol/water partition coefficient for drug design using the GB/SA approach. *J. Chem. Inf. Model.* \(2014\) 54\(12\):3284-3301.](#)

For details about development and validation of the BOILED-Egg, please refer to this article: [A BOILED-Egg to predict gastrointestinal absorption and brain penetration of small molecules. *ChemMedChem* \(2016\) 11\(11\):1117-1121.](#)

Developed and maintained by the [Molecular Modeling Group](#) of the SIB | Swiss Institute of Bioinformatics.

In order to avoid security-related warning messages when switching to secured connection, you may want either to:

- confirm the exception on the next page, or
- import our [CA key](#) in your web browser

Click [here](#) to proceed.

[Click2Drug](#)

[SwissDock](#)

[SwissParam](#)

[SwissSidechain](#)

[SwissBioisostere](#)

[SwissTargetPrediction](#)

[SwissADME](#)

[SwissSimilarity](#)

[About us](#)

[Swiss Institute of Bioinformatics](#)

SwissADME

- [Disclaimer](#)
- [Help](#)
- [FAQ](#)
- [Home](#)

This website allows you to compute physicochemical descriptors as well as to predict ADME parameters, pharmacokinetic properties, druglike nature and medicinal chemistry friendliness of one or multiple small molecules to support drug discovery.

The main article describing the web service and its underlying methodologies is [SwissADME: a free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules. *Sci. Rep.* \(2017\) 7:42717.](#)

For details about development and validation of iLOG, please refer to this article: [iLOGP: a simple, robust, and efficient description of *n*-octanol/water partition coefficient for drug design using the GB/SA approach. *J. Chem. Inf. Model.* \(2014\) 54\(12\):3284-3301.](#)

For details about development and validation of the BOILED-Egg, please refer to this article: [A BOILED-Egg to predict gastrointestinal absorption and brain penetration of small molecules. *ChemMedChem* \(2016\) 11\(11\):1117-1121.](#)

Developed and maintained by the [Molecular Modeling Group](#) of the SIB | Swiss Institute of Bioinformatics.

In order to avoid security-related warning messages when switching to secured connection, you may want either to:

- confirm the exception on the next page, or
- import our [CA key](#) in your web browser

Click [here](#) to proceed.

[Click2Drug](#)

[SwissDock](#)

[SwissParam](#)

[SwissSidechain](#)

[SwissBioisostere](#)

[SwissTargetPrediction](#)

[SwissADME](#)

[SwissSimilarity](#)

[About us](#)

[Swiss Institute of Bioinformatics](#)

SwissADME

- [Disclaimer](#)
- [Help](#)
- [FAQ](#)
- [Home](#)


This website allows you to compute physicochemical descriptors as well as to predict ADME parameters, pharmacokinetic properties, druglike nature and medicinal chemistry friendliness of one or multiple small molecules to support drug discovery.

The main article describing the web service and its underlying methodologies is [SwissADME: a free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules. *Sci. Rep.* \(2017\) 7:42717.](#)

For details about development and validation of iLOG, please refer to this article: [iLOGP: a simple, robust, and efficient description of *n*-octanol/water partition coefficient for drug design using the GB/SA approach. *J. Chem. Inf. Model.* \(2014\) 54\(12\):3284-3301.](#)

For details about development and validation of the BOILED-Egg, please refer to this article: [A BOILED-Egg to predict gastrointestinal absorption and brain penetration of small molecules. *ChemMedChem* \(2016\) 11\(11\):1117-1121.](#)

Developed and maintained by the [Molecular Modeling Group](#) of the SIB | Swiss Institute of Bioinformatics.



Enter a list of SMILES here:

```
Brc1ccccc1OC2=CC(=O)c1ccccc1C2=O
```

Fill with an example

Clear



Run!

Show BOILED-Egg

Retrieve data:


 Powered by
ChemAxon

Molecule 1


 SMILES
S Brc1ccccc1OC1=CC(=O)c2c(C1=O)cccc2

Physicochemical Properties

Formula	C16H9BrO3
Molecular weight	329.14 g/mol
Num. heavy atoms	20
Num. arom. heavy atoms	12
Fraction Csp3	0.00
Num. rotatable bonds	2
Num. H-bond acceptors	3
Num. H-bond donors	0
Molar Refractivity	77.96
TPSA	

Water Solubility

Log S (ESOL)

ESOL: Topological method implemented from
Delaney JS. 2004 J. Chem. Inf. Model.

-4.46

 Solubility
Class

1.14e-02 mg/ml ; 3.46e-05 mol/l

Solubility class: Log S scale

Insoluble < -10 < Poorly Moderately soluble
< -6 < Moderately < -4
< Soluble < -2 Very < 0
< Highly

Log S (Ali)

Ali: Topological method implemented from
Ali J. et al. 2012 J. Chem. Inf. Model.

-4.20

 Solubility
Class

2.09e-02 mg/ml ; 6.34e-05 mol/l

Solubility class: Log S

In order to avoid security-related warning messages when switching to secured connection, you may want either to:

- confirm the exception on the next page, or
- import our [CA key](#) in your web browser

Click [here](#) to proceed.

[Click2Drug](#)

[SwissDock](#)

[SwissParam](#)

[SwissSidechain](#)

[SwissBioisostere](#)

[SwissTargetPrediction](#)

[SwissADME](#)

[SwissSimilarity](#)

[About us](#)

[Swiss Institute of Bioinformatics](#)

SwissADME

- [Disclaimer](#)
- [Help](#)
- [FAQ](#)
- [Home](#)

This website allows you to compute physicochemical descriptors as well as to predict ADME parameters, pharmacokinetic properties, druglike nature and medicinal chemistry friendliness of one or multiple small molecules to support drug discovery.

The main article describing the web service and its underlying methodologies is [SwissADME: a free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules. *Sci. Rep.* \(2017\) 7:42717.](#)

For details about development and validation of iLOG, please refer to this article: [iLOGP: a simple, robust, and efficient description of *n*-octanol/water partition coefficient for drug design using the GB/SA approach. *J. Chem. Inf. Model.* \(2014\) 54\(12\):3284-3301.](#)

For details about development and validation of the BOILED-Egg, please refer to this article: [A BOILED-Egg to predict gastrointestinal absorption and brain penetration of small molecules. *ChemMedChem* \(2016\) 11\(11\):1117-1121.](#)

Developed and maintained by the [Molecular Modeling Group](#) of the SIB | Swiss Institute of Bioinformatics.

In order to avoid security-related warning messages when switching to secured connection, you may want either to:

- confirm the exception on the next page, or
- import our [CA key](#) in your web browser

Click [here](#) to proceed.

[Click2Drug](#)

[SwissDock](#)

[SwissParam](#)

[SwissSidechain](#)

[SwissBioisostere](#)

[SwissTargetPrediction](#)

[SwissADME](#)

[SwissSimilarity](#)

[About us](#)

[Swiss Institute of Bioinformatics](#)

SwissADME

- [Disclaimer](#)
- [Help](#)
- [FAQ](#)
- [Home](#)


This website allows you to compute physicochemical descriptors as well as to predict ADME parameters, pharmacokinetic properties, druglike nature and medicinal chemistry friendliness of one or multiple small molecules to support drug discovery.

The main article describing the web service and its underlying methodologies is [SwissADME: a free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules. *Sci. Rep.* \(2017\) 7:42717.](#)

For details about development and validation of iLOG, please refer to this article: [iLOGP: a simple, robust, and efficient description of *n*-octanol/water partition coefficient for drug design using the GB/SA approach. *J. Chem. Inf. Model.* \(2014\) 54\(12\):3284-3301.](#)

For details about development and validation of the BOILED-Egg, please refer to this article: [A BOILED-Egg to predict gastrointestinal absorption and brain penetration of small molecules. *ChemMedChem* \(2016\) 11\(11\):1117-1121.](#)

Developed and maintained by the [Molecular Modeling Group](#) of the SIB | Swiss Institute of Bioinformatics.



Enter a list of SMILES here:

Cc3ccccc3OC2=CC(=O)c1ccccc1C2=O

Fill with an example

Clear



Run!

Show BOILED-Egg

Retrieve data:


 Powered by
ChemAxon

Molecule 1


 SMILES
S Cc1ccccc1OC1=CC(=O)c2c(C1=O)cccc2

Physicochemical Properties

Formula	C17H12O3
Molecular weight	264.28 g/mol
Num. heavy atoms	20
Num. arom. heavy atoms	12
Fraction Csp3	0.06
Num. rotatable bonds	2
Num. H-bond acceptors	3
Num. H-bond donors	0
Molar Refractivity	75.22
TPSA	

Water Solubility

Log S (ESOL)

ESOL: Topological method implemented from
Delaney JS. 2004 J. Chem. Inf. Model.

-3.85

 Solubility
Class

3.73e-02 mg/ml ; 1.41e-04 mol/l

Solubility class: Log S scale

Insoluble < -10 < Poorly Soluble
< -6 < Moderately < -4
< Soluble < -2 Very < 0
< Highly

Log S (Ali)

Ali: Topological method implemented from
Ali J. et al. 2012 J. Chem. Inf. Model.

-3.86

 Solubility
Class

3.69e-02 mg/ml ; 1.40e-04 mol/l

Solubility class: Log S

In order to avoid security-related warning messages when switching to secured connection, you may want either to:

- confirm the exception on the next page, or
- import our [CA key](#) in your web browser

Click [here](#) to proceed.

[Click2Drug](#)

[SwissDock](#)

[SwissParam](#)

[SwissSidechain](#)

[SwissBioisostere](#)

[SwissTargetPrediction](#)

[SwissADME](#)

[SwissSimilarity](#)

[About us](#)

[Swiss Institute of Bioinformatics](#)

SwissADME

- [Disclaimer](#)
- [Help](#)
- [FAQ](#)
- [Home](#)

This website allows you to compute physicochemical descriptors as well as to predict ADME parameters, pharmacokinetic properties, druglike nature and medicinal chemistry friendliness of one or multiple small molecules to support drug discovery.

The main article describing the web service and its underlying methodologies is [SwissADME: a free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules. *Sci. Rep.* \(2017\) 7:42717.](#)

For details about development and validation of iLOGP, please refer to this article: [iLOGP: a simple, robust, and efficient description of *n*-octanol/water partition coefficient for drug design using the GB/SA approach. *J. Chem. Inf. Model.* \(2014\) 54\(12\):3284-3301.](#)

For details about development and validation of the BOILED-Egg, please refer to this article: [A BOILED-Egg to predict gastrointestinal absorption and brain penetration of small molecules. *ChemMedChem* \(2016\) 11\(11\):1117-1121.](#)

Developed and maintained by the [Molecular Modeling Group](#) of the SIB | Swiss Institute of Bioinformatics.

In order to avoid security-related warning messages when switching to secured connection, you may want either to:

- confirm the exception on the next page, or
- import our [CA key](#) in your web browser

Click [here](#) to proceed.

[Click2Drug](#)

[SwissDock](#)

[SwissParam](#)

[SwissSidechain](#)

[SwissBioisostere](#)

[SwissTargetPrediction](#)

[SwissADME](#)

[SwissSimilarity](#)

[About us](#)

[Swiss Institute of Bioinformatics](#)

SwissADME

- [Disclaimer](#)
- [Help](#)
- [FAQ](#)
- [Home](#)


This website allows you to compute physicochemical descriptors as well as to predict ADME parameters, pharmacokinetic properties, druglike nature and medicinal chemistry friendliness of one or multiple small molecules to support drug discovery.

The main article describing the web service and its underlying methodologies is [SwissADME: a free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules. *Sci. Rep.* \(2017\) 7:42717.](#)

For details about development and validation of iLOG, please refer to this article: [iLOGP: a simple, robust, and efficient description of *n*-octanol/water partition coefficient for drug design using the GB/SA approach. *J. Chem. Inf. Model.* \(2014\) 54\(12\):3284-3301.](#)

For details about development and validation of the BOILED-Egg, please refer to this article: [A BOILED-Egg to predict gastrointestinal absorption and brain penetration of small molecules. *ChemMedChem* \(2016\) 11\(11\):1117-1121.](#)

Developed and maintained by the [Molecular Modeling Group](#) of the SIB | Swiss Institute of Bioinformatics.


[Show BOILED-Egg](#)



Enter a list of SMILES here:

COc3ccccc3OC2=CC(=O)c1ccccc1C2=O
[Fill with an example](#)
[Clear](#)
[Run!](#)

Retrieve data:


 Powered by
ChemAxon

Molecule 1


 SMILES
S COc1ccccc1OC1=CC(=O)c2c(C1=O)cccc2

Physicochemical Properties

Formula	C17H12O4
Molecular weight	280.27 g/mol
Num. heavy atoms	21
Num. arom. heavy atoms	12
Fraction Csp3	0.06
Num. rotatable bonds	3
Num. H-bond acceptors	4
Num. H-bond donors	0
Molar Refractivity	76.75
TPSA	

Water Solubility

Log S (ESOL)

ESOL: Topological method implemented from
Delaney JS. 2004 J. Chem. Inf. Model.

-3.62

 Solubility
Class

6.77e-02 mg/ml ; 2.42e-04 mol/l

Solubility class: Log S scale

Insoluble < -10 < Poorly Soluble
< -6 < Moderately < -4
< Soluble < -2 Very < 0
< Highly

Log S (Ali)

Ali: Topological method implemented from
Ali J. et al. 2012 J. Chem. Inf. Model.

-3.64

 Solubility
Class

6.36e-02 mg/ml ; 2.27e-04 mol/l

Solubility class: Log S

In order to avoid security-related warning messages when switching to secured connection, you may want either to:

- confirm the exception on the next page, or
- import our [CA key](#) in your web browser

Click [here](#) to proceed.

[Click2Drug](#)

[SwissDock](#)

[SwissParam](#)

[SwissSidechain](#)

[SwissBioisostere](#)

[SwissTargetPrediction](#)

[SwissADME](#)

[SwissSimilarity](#)

[About us](#)

[Swiss Institute of Bioinformatics](#)

SwissADME

- [Disclaimer](#)
- [Help](#)
- [FAQ](#)
- [Home](#)

This website allows you to compute physicochemical descriptors as well as to predict ADME parameters, pharmacokinetic properties, druglike nature and medicinal chemistry friendliness of one or multiple small molecules to support drug discovery.

The main article describing the web service and its underlying methodologies is [SwissADME: a free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules](#). *Sci. Rep.* (2017) 7:42717.

For details about development and validation of iLOG, please refer to this article: [iLOGP: a simple, robust, and efficient description of *n*-octanol/water partition coefficient for drug design using the GB/SA approach](#). *J. Chem. Inf. Model.* (2014) 54(12):3284-3301.

For details about development and validation of the BOILED-Egg, please refer to this article: [A BOILED-Egg to predict gastrointestinal absorption and brain penetration of small molecules](#). *ChemMedChem* (2016) 11(11):1117-1121.

Developed and maintained by the [Molecular Modeling Group](#) of the SIB | Swiss Institute of Bioinformatics.

In order to avoid security-related warning messages when switching to secured connection, you may want either to:

- confirm the exception on the next page, or
- import our [CA key](#) in your web browser

Click [here](#) to proceed.

[Click2Drug](#)

[SwissDock](#)

[SwissParam](#)

[SwissSidechain](#)

[SwissBioisostere](#)

[SwissTargetPrediction](#)

[SwissADME](#)

[SwissSimilarity](#)

[About us](#)

[Swiss Institute of Bioinformatics](#)

SwissADME

- [Disclaimer](#)
- [Help](#)
- [FAQ](#)
- [Home](#)


This website allows you to compute physicochemical descriptors as well as to predict ADME parameters, pharmacokinetic properties, druglike nature and medicinal chemistry friendliness of one or multiple small molecules to support drug discovery.

The main article describing the web service and its underlying methodologies is [SwissADME: a free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules. *Sci. Rep.* \(2017\) 7:42717.](#)

For details about development and validation of iLOG, please refer to this article: [iLOGP: a simple, robust, and efficient description of *n*-octanol/water partition coefficient for drug design using the GB/SA approach. *J. Chem. Inf. Model.* \(2014\) 54\(12\):3284-3301.](#)

For details about development and validation of the BOILED-Egg, please refer to this article: [A BOILED-Egg to predict gastrointestinal absorption and brain penetration of small molecules. *ChemMedChem* \(2016\) 11\(11\):1117-1121.](#)

Developed and maintained by the [Molecular Modeling Group](#) of the SIB | Swiss Institute of Bioinformatics.



Enter a list of SMILES here:

```
BrC1ccc(cc1)OC3=CC(=O)c2ccccc2C3=O
```

Fill with an example

Clear



Run!

Show BOILED-Egg

Retrieve data:


 Powered by
ChemAxon

Molecule 1


 SMILES
S BrC1ccc(cc1)OC1=CC(=O)c2ccccc2C1=O

Physicochemical Properties

Formula	C16H9BrO3
Molecular weight	329.14 g/mol
Num. heavy atoms	20
Num. arom. heavy atoms	12
Fraction Csp3	0.00
Num. rotatable bonds	2
Num. H-bond acceptors	3
Num. H-bond donors	0
Molar Refractivity	77.96
TPSA	

Water Solubility

Log S (ESOL)

ESOL: Topological method implemented from
Delaney JS. 2004 J. Chem. Inf. Model.

-4.46

 Solubility
Class

1.14e-02 mg/ml ; 3.46e-05 mol/l

Solubility class: Log S scale

Insoluble < -10 < Poorly Moderately soluble
< -6 < Moderately < -4
< Soluble < -2 Very < 0
< Highly

Log S (Ali)

Ali: Topological method implemented from
Ali J. et al. 2012 J. Chem. Inf. Model.

-4.20

 Solubility
Class

2.09e-02 mg/ml ; 6.34e-05 mol/l

Solubility class: Log S

In order to avoid security-related warning messages when switching to secured connection, you may want either to:

- confirm the exception on the next page, or
- import our [CA key](#) in your web browser

Click [here](#) to proceed.

[Click2Drug](#)

[SwissDock](#)

[SwissParam](#)

[SwissSidechain](#)

[SwissBioisostere](#)

[SwissTargetPrediction](#)

[SwissADME](#)

[SwissSimilarity](#)

[About us](#)

[Swiss Institute of Bioinformatics](#)

SwissADME

- [Disclaimer](#)
- [Help](#)
- [FAQ](#)
- [Home](#)

This website allows you to compute physicochemical descriptors as well as to predict ADME parameters, pharmacokinetic properties, druglike nature and medicinal chemistry friendliness of one or multiple small molecules to support drug discovery.

The main article describing the web service and its underlying methodologies is [SwissADME: a free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules. *Sci. Rep.* \(2017\) 7:42717.](#)

For details about development and validation of iLOG, please refer to this article: [iLOGP: a simple, robust, and efficient description of *n*-octanol/water partition coefficient for drug design using the GB/SA approach. *J. Chem. Inf. Model.* \(2014\) 54\(12\):3284-3301.](#)

For details about development and validation of the BOILED-Egg, please refer to this article: [A BOILED-Egg to predict gastrointestinal absorption and brain penetration of small molecules. *ChemMedChem* \(2016\) 11\(11\):1117-1121.](#)

Developed and maintained by the [Molecular Modeling Group](#) of the SIB | Swiss Institute of Bioinformatics.

In order to avoid security-related warning messages when switching to secured connection, you may want either to:

- confirm the exception on the next page, or
- import our [CA key](#) in your web browser

Click [here](#) to proceed.

[Click2Drug](#)

[SwissDock](#)

[SwissParam](#)

[SwissSidechain](#)

[SwissBioisostere](#)

[SwissTargetPrediction](#)

[SwissADME](#)

[SwissSimilarity](#)

[About us](#)

[Swiss Institute of Bioinformatics](#)

SwissADME

- [Disclaimer](#)
- [Help](#)
- [FAQ](#)
- [Home](#)


This website allows you to compute physicochemical descriptors as well as to predict ADME parameters, pharmacokinetic properties, druglike nature and medicinal chemistry friendliness of one or multiple small molecules to support drug discovery.

The main article describing the web service and its underlying methodologies is [SwissADME: a free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules. *Sci. Rep.* \(2017\) 7:42717.](#)

For details about development and validation of iLOG, please refer to this article: [iLOGP: a simple, robust, and efficient description of *n*-octanol/water partition coefficient for drug design using the GB/SA approach. *J. Chem. Inf. Model.* \(2014\) 54\(12\):3284-3301.](#)

For details about development and validation of the BOILED-Egg, please refer to this article: [A BOILED-Egg to predict gastrointestinal absorption and brain penetration of small molecules. *ChemMedChem* \(2016\) 11\(11\):1117-1121.](#)

Developed and maintained by the [Molecular Modeling Group](#) of the SIB | Swiss Institute of Bioinformatics.



Enter a list of SMILES here:

Cc1ccc(cc1)OC3=CC(=O)c2ccccc2C3=O

Fill with an example

Clear



Run!

Show BOILED-Egg

Retrieve data:


 Powered by
ChemAxon

Molecule 1


 SMILES
S Cc1ccc(cc1)OC1=CC(=O)c2c(C1=O)cccc2

Physicochemical Properties

Formula	C17H12O3
Molecular weight	264.28 g/mol
Num. heavy atoms	20
Num. arom. heavy atoms	12
Fraction Csp3	0.06
Num. rotatable bonds	2
Num. H-bond acceptors	3
Num. H-bond donors	0
Molar Refractivity	75.22
TPSA	

Water Solubility

Log S (ESOL)

ESOL: Topological method implemented from
Delaney JS. 2004 J. Chem. Inf. Model.

-3.85

 Solubility
Class

3.73e-02 mg/ml ; 1.41e-04 mol/l

Solubility class: Log S scale

Insoluble < -10 < Poorly Soluble
< -6 < Moderately < -4
< Soluble < -2 Very < 0
< Highly

Log S (Ali)

Ali: Topological method implemented from
Ali J. et al. 2012 J. Chem. Inf. Model.

-3.86

 Solubility
Class

3.69e-02 mg/ml ; 1.40e-04 mol/l

Solubility class: Log S

In order to avoid security-related warning messages when switching to secured connection, you may want either to:

- confirm the exception on the next page, or
- import our [CA key](#) in your web browser

Click [here](#) to proceed.

[Click2Drug](#)

[SwissDock](#)

[SwissParam](#)

[SwissSidechain](#)

[SwissBioisostere](#)

[SwissTargetPrediction](#)

[SwissADME](#)

[SwissSimilarity](#)

[About us](#)

[Swiss Institute of Bioinformatics](#)

SwissADME

- [Disclaimer](#)
- [Help](#)
- [FAQ](#)
- [Home](#)

This website allows you to compute physicochemical descriptors as well as to predict ADME parameters, pharmacokinetic properties, druglike nature and medicinal chemistry friendliness of one or multiple small molecules to support drug discovery.

The main article describing the web service and its underlying methodologies is [SwissADME: a free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules. *Sci. Rep.* \(2017\) 7:42717.](#)

For details about development and validation of iLOG, please refer to this article: [iLOGP: a simple, robust, and efficient description of *n*-octanol/water partition coefficient for drug design using the GB/SA approach. *J. Chem. Inf. Model.* \(2014\) 54\(12\):3284-3301.](#)

For details about development and validation of the BOILED-Egg, please refer to this article: [A BOILED-Egg to predict gastrointestinal absorption and brain penetration of small molecules. *ChemMedChem* \(2016\) 11\(11\):1117-1121.](#)

Developed and maintained by the [Molecular Modeling Group](#) of the SIB | Swiss Institute of Bioinformatics.

In order to avoid security-related warning messages when switching to secured connection, you may want either to:

- confirm the exception on the next page, or
- import our [CA key](#) in your web browser

Click [here](#) to proceed.

[Click2Drug](#)

[SwissDock](#)

[SwissParam](#)

[SwissSidechain](#)

[SwissBioisostere](#)

[SwissTargetPrediction](#)

[SwissADME](#)

[SwissSimilarity](#)

[About us](#)

[Swiss Institute of Bioinformatics](#)

SwissADME

- [Disclaimer](#)
- [Help](#)
- [FAQ](#)
- [Home](#)


This website allows you to compute physicochemical descriptors as well as to predict ADME parameters, pharmacokinetic properties, druglike nature and medicinal chemistry friendliness of one or multiple small molecules to support drug discovery.

The main article describing the web service and its underlying methodologies is [SwissADME: a free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules. *Sci. Rep.* \(2017\) 7:42717.](#)

For details about development and validation of iLOG, please refer to this article: [iLOGP: a simple, robust, and efficient description of *n*-octanol/water partition coefficient for drug design using the GB/SA approach. *J. Chem. Inf. Model.* \(2014\) 54\(12\):3284-3301.](#)

For details about development and validation of the BOILED-Egg, please refer to this article: [A BOILED-Egg to predict gastrointestinal absorption and brain penetration of small molecules. *ChemMedChem* \(2016\) 11\(11\):1117-1121.](#)

Developed and maintained by the [Molecular Modeling Group](#) of the SIB | Swiss Institute of Bioinformatics.





Enter a list of SMILES here:

O=C6C=C(Oc1ccc(cc1)C(c2ccccc2)(c3ccccc3)c4ccccc4)C(=O)c5ccccc56

Retrieve data:


 Powered by
ChemAxon

Molecule 1



SMILE O=C1C(=CC(=O)c2c1ccccc2)Oc1ccc(cc1)C(c1ccccc1)(c1ccccc1)c1ccccc1

Physicochemical Properties

Formula	C35H24O3
Molecular weight	492.56 g/mol
Num. heavy atoms	38
Num. arom. heavy atoms	30
Fraction Csp3	0.03
Num. rotatable bonds	6
Num. H-bond acceptors	3
Num. H-bond donors	0
Molar Refractivity	148.57
TPSA	

Water Solubility

Log *S* (ESOL)

ESOL: Topological method implemented from Delaney JS. 2004 J. Chem. Inf. Model.

-8.09

Solubility Class

4.00e-06 mg/ml ; 8.12e-09 mol/l

Solubility class: Log *S* scale

Insoluble < -10 < Poorly Poorly soluble
< -6 < Moderately < -4
< Soluble < -2 Very < 0
< Highly

Log *S* (Ali)

Ali: Topological method implemented from Ali J. et al. 2012 J. Chem. Inf. Model.

-8.71

Solubility Class

9.57e-07 mg/ml ; 1.94e-09 mol/l

Solubility class: Log *S*

Topological Polar Surface Area: Calculated from Ertl P. et al. 2000 J. Med. Chem.	43.37 Å ²	<u>scale</u> Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Poorly soluble
Log $P_{o/w}$ (iLOGP) ⓘ	Lipophilicity	Log S (SILICOS-IT) ⓘ	
iLOGP: in-house physics-based method implemented from Daina A et al. 2014 J. Chem. Inf. Model.	4.29	SILICOS-IT: Fragmental method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos-it.com	-12.72
Log $P_{o/w}$ (XLOGP3) ⓘ		Solubility	9.36e-11 mg/ml ; 1.90e-13 mol/l
XLOGP3: Atomistic and knowledge-based method calculated by XLOGP program, version 3.2.2, courtesy of CCBG, Shanghai Institute of Organic Chemistry	7.95	Class ⓘ	
Log $P_{o/w}$ (WLOGP) ⓘ		Solubility class: Log S <u>scale</u> Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Insoluble
WLOGP: Atomistic method implemented from Wildman SA and Crippen GM. 1999 J. Chem. Inf. Model.	7.41		Pharmacokinetics
Log $P_{o/w}$ (MLOGP) ⓘ		GI absorption ⓘ	
MLOGP: Topological method implemented from Moriguchi I. et al. 1992 Chem. Pharm. Bull. Moriguchi I. et al. 1994 Chem. Pharm. Bull. Lipinski PA. et al. 2001 Adv. Drug. Deliv. Rev.	4.99	Gastrointestinal absorption: according to the white of the BOILED-Egg	Low
Log $P_{o/w}$ (SILICOS-IT) ⓘ		BBB permeant ⓘ	
SILICOS-IT: Hybrid fragmental/topological method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos-it.com	7.82	BBB permeation: according to the yolk of the BOILED-Egg	No
Consensus Log $P_{o/w}$ ⓘ		P-gp substrate ⓘ	
Consensus Log $P_{o/w}$:		P-glycoprotein substrate: SVM model built on 1033 molecules (training set) and tested on 415 molecules (test set) 10-fold CV: ACC=0.72 / AUC=0.77 External: ACC=0.88 / AUC=0.94	No
		CYP1A2 inhibitor ⓘ	
		Cytochrome P450 1A2 inhibitor: SVM model built on 9145 molecules (training set) and tested on 3000 molecules (test set)	Yes

Average of all five
predictions

6.49

10-fold CV: ACC=0.83 /
AUC=0.90
External: ACC=0.84 /
AUC=0.91

CYP2C19 inhibitor ⓘ

Cytochrome P450
2C19 inhibitor: SVM
model built on 9272
molecules (training set)
and tested on 3000
molecules (test set) No
10-fold CV: ACC=0.80 /
AUC=0.86
External: ACC=0.80 /
AUC=0.87

CYP2C9 inhibitor ⓘ

Cytochrome P450 2C9
inhibitor: SVM model
built on 5940 molecules
(training set)
and tested on 2075
molecules (test set) No
10-fold CV: ACC=0.78 /
AUC=0.85
External: ACC=0.71 /
AUC=0.81

CYP2D6 inhibitor ⓘ

Cytochrome P450 2D6
inhibitor: SVM model
built on 3664 molecules
(training set)
and tested on 1068
molecules (test set) No
10-fold CV: ACC=0.79 /
AUC=0.85
External: ACC=0.81 /
AUC=0.87

CYP3A4 inhibitor ⓘ

Cytochrome P450 3A4
inhibitor: SVM model
built on 7518 molecules
(training set)
and tested on 2579
molecules (test set) No
10-fold CV: ACC=0.77 /
AUC=0.85
External: ACC=0.78 /
AUC=0.86

Log K_p (skin
permeation) ⓘ

Skin permeation:
QSPR model
implemented from -3.66 cm/s

Potts RO and Guy RH.
1992 Pharm. Res.

Druglikeness

Lipinski

Lipinski (Pfizer) filter:
implemented from

Lipinski CA. et al. 2001

Adv. Drug Deliv. Rev.

Yes; 1 violation: MLOGP>4.15

MW ≤ 500

MLOGP ≤ 4.15

N or O ≤ 10

NH or OH ≤ 5

Ghose

Ghose filter:

implemented from

Ghose AK. et al. 1999 J.

Comb. Chem.

No; 3 violations: MW>480,
WLOGP>5.6, MR>130

160 ≤ MW ≤ 480

-0.4 ≤ WLOGP ≤ 5.6

40 ≤ MR ≤ 130

20 ≤ atoms ≤ 70

Veber

Veber (GSK) filter:

implemented from

Veber DF. et al. 2002 J.

Yes

Med. Chem.

Rotatable bonds ≤ 10

TPSA ≤ 140

Egan

Egan (Pharmacia)

filter: implemented

from

Egan WJ. et al. 2000 J.

No; 1 violation: WLOGP>5.88

Med. Chem.

WLOGP ≤ 5.88

TPSA ≤ 131.6

Muegge

Muegge (Bayer) filter:

implemented from

Muegge I. et al. 2001 J.

Med. Chem.

200 ≤ MW ≤ 600

-2 ≤ XLOGP ≤ 5

TPSA ≤ 150

No; 1 violation: XLOGP3>5

Num. rings ≤ 7

Num. carbon > 4

Num. heteroatoms > 1

Num. rotatable bonds ≤

15

H-bond acc. ≤ 10

H-bond don. ≤ 5

Bioavailability Score ⓘ


Abbott Bioavailability
Score: Probability of F 0.56
> 10% in rat
implemented from
Martin YC. 2005 J.
Med. Chem.

Medicinal Chemistry

PAINS ⓘ

1 alert: quinone_A ⓘ

Pan Assay Interference
Structures:
implemented from
Baell JB. & Holloway
GA. 2010 J. Med.
Chem.



Brenk ⓘ


Structural Alert:
implemented from 0 alert
Brenk R. et al. 2008
ChemMedChem

Leadlikeness ⓘ

Leadlikeness:
implemented from
Teague SJ. 1999 Angew. No; 2 violations: MW>350,
Chem. Int. Ed. XLOGP3>3.5
 $250 \leq MW \leq 350$
 $XLOGP \leq 3.5$
Num. rotatable bonds $\leq$
7

Synthetic accessibility ⓘ

Synthetic accessibility
score: from 1 (very
easy) to 10 (very
difficult)
based on 1024
fragmental contributions 4.50
(FP2) modulated by size
and complexity penalties,
trained on 12'782'590
molecules and tested on
40 external molecules
($r^2 = 0.94$)



Enter a list of SMILES here:

```
CC(C)(C)c1ccc(cc1)OC3=CC(=O)c2ccccc2C3=O
```

Fill with an example

Clear



Run!

Show BOILED-Egg

Retrieve data:


 Powered by
ChemAxon

Molecule 1



SMILES
S O=C1C(=CC(=O)c2c1cccc2)Oc1ccc(cc1)C(C)(C)C

Physicochemical Properties

Formula	C20H18O3
Molecular weight	306.36 g/mol
Num. heavy atoms	23
Num. arom. heavy atoms	12
Fraction Csp3	0.20
Num. rotatable bonds	3
Num. H-bond acceptors	3
Num. H-bond donors	0
Molar Refractivity	89.53
TPSA	

Water Solubility

Log S (ESOL)

ESOL: Topological method implemented from
Delaney JS. 2004 J. Chem. Inf. Model.

-4.81

Solubility
Class

4.71e-03 mg/ml ; 1.54e-05 mol/l

Solubility class: Log S scale

Insoluble < -10 < Poorly Moderately soluble
< -6 < Moderately < -4
< Soluble < -2 Very < 0
< Highly

Log S (Ali)

Ali: Topological method implemented from
Ali J. et al. 2012 J. Chem. Inf. Model.

-5.21

Solubility
Class

1.87e-03 mg/ml ; 6.10e-06 mol/l

Solubility class: Log S

Topological Polar Surface Area: Calculated from Ertl P. et al. 2000 J. Med. Chem.	43.37 Å ²	<u>scale</u> Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Moderately soluble
Log $P_{o/w}$ (iLOGP) ⓘ	Lipophilicity	Log S (SILICOS-IT) ⓘ	
iLOGP: in-house physics-based method implemented from Daina A et al. 2014 J. Chem. Inf. Model.	3.10	SILICOS-IT: Fragmental method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos-it.com	-6.49
Log $P_{o/w}$ (XLOGP3) ⓘ		Solubility Class ⓘ	9.94e-05 mg/ml ; 3.24e-07 mol/l
XLOGP3: Atomistic and knowledge-based method calculated by XLOGP program, version 3.2.2, courtesy of CCBG, Shanghai Institute of Organic Chemistry	4.58	Solubility class: Log S <u>scale</u> Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Poorly soluble
Log $P_{o/w}$ (WLOGP) ⓘ			Pharmacokinetics
WLOGP: Atomistic method implemented from Wildman SA and Crippen GM. 1999 J. Chem. Inf. Model.	4.33	GI absorption ⓘ	
Log $P_{o/w}$ (MLOGP) ⓘ		Gastrointestinal absorption: according to the white of the BOILED-Egg	High
MLOGP: Topological method implemented from Moriguchi I. et al. 1992 Chem. Pharm. Bull. Moriguchi I. et al. 1994 Chem. Pharm. Bull. Lipinski PA. et al. 2001 Adv. Drug. Deliv. Rev.	2.62	BBB permeant ⓘ	
Log $P_{o/w}$ (SILICOS-IT) ⓘ		BBB permeation: according to the yolk of the BOILED-Egg	Yes
SILICOS-IT: Hybrid fragmental/topological method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos-it.com	4.66	P-gp substrate ⓘ	
Consensus Log $P_{o/w}$ ⓘ		P-glycoprotein substrate: SVM model built on 1033 molecules (training set) and tested on 415 molecules (test set) 10-fold CV: ACC=0.72 / AUC=0.77 External: ACC=0.88 / AUC=0.94	No
Consensus Log $P_{o/w}$:		CYP1A2 inhibitor ⓘ	
		Cytochrome P450 1A2 inhibitor: SVM model built on 9145 molecules (training set) and tested on 3000 molecules (test set)	Yes

Average of all five predictions

3.86

10-fold CV: ACC=0.83 / AUC=0.90
External: ACC=0.84 / AUC=0.91

CYP2C19 inhibitor ⓘ

Cytochrome P450 2C19 inhibitor: SVM model built on 9272 molecules (training set) and tested on 3000 molecules (test set) Yes
10-fold CV: ACC=0.80 / AUC=0.86
External: ACC=0.80 / AUC=0.87

CYP2C9 inhibitor ⓘ

Cytochrome P450 2C9 inhibitor: SVM model built on 5940 molecules (training set) and tested on 2075 molecules (test set) Yes
10-fold CV: ACC=0.78 / AUC=0.85
External: ACC=0.71 / AUC=0.81

CYP2D6 inhibitor ⓘ

Cytochrome P450 2D6 inhibitor: SVM model built on 3664 molecules (training set) and tested on 1068 molecules (test set) No
10-fold CV: ACC=0.79 / AUC=0.85
External: ACC=0.81 / AUC=0.87

CYP3A4 inhibitor ⓘ

Cytochrome P450 3A4 inhibitor: SVM model built on 7518 molecules (training set) and tested on 2579 molecules (test set) Yes
10-fold CV: ACC=0.77 / AUC=0.85
External: ACC=0.78 / AUC=0.86

Log K_p (skin permeation) ⓘ

Skin permeation: QSPR model implemented from -4.92 cm/s

Potts RO and Guy RH.
1992 Pharm. Res.

Druglikeness

Lipinski ⓘ

Lipinski (Pfizer) filter:
implemented from
Lipinski CA. et al. 2001
Adv. Drug Deliv. Rev. Yes; 0 violation
MW $\leq$ 500
MLOGP $\leq$ 4.15
N or O $\leq$ 10
NH or OH $\leq$ 5

Ghose ⓘ

Ghose filter:
implemented from
Ghose AK. et al. 1999 J.
Comb. Chem. Yes
160 $\leq$ MW $\leq$ 480
-0.4 $\leq$ WLOGP $\leq$ 5.6
40 $\leq$ MR $\leq$ 130
20 $\leq$ atoms $\leq$ 70

Veber ⓘ

Veber (GSK) filter:
implemented from
Veber DF. et al. 2002 J. Yes
Med. Chem.
Rotatable bonds $\leq$ 10
TPSA $\leq$ 140

Egan ⓘ

Egan (Pharmacia)
filter: implemented
from
Egan WJ. et al. 2000 J. Yes
Med. Chem.
WLOGP $\leq$ 5.88
TPSA $\leq$ 131.6

Muegge ⓘ

Muegge (Bayer) filter:
implemented from
Muegge I. et al. 2001 J.
Med. Chem.
200 $\leq$ MW $\leq$ 600
-2 $\leq$ XLOGP $\leq$ 5
TPSA $\leq$ 150 Yes
Num. rings $\leq$ 7
Num. carbon $>$ 4
Num. heteroatoms $>$ 1
Num. rotatable bonds $\leq$
15
H-bond acc. $\leq$ 10
H-bond don. $\leq$ 5

Bioavailability Score ⓘ


Abbott Bioavailability
Score: Probability of F 0.56
> 10% in rat
implemented from
Martin YC. 2005 J.
Med. Chem.

Medicinal Chemistry

PAINS ⓘ

1 alert: quinone_A ⓘ

Pan Assay Interference
Structures:
implemented from
Baell JB. & Holloway
GA. 2010 J. Med.
Chem.



Brenk ⓘ

Structural Alert:
implemented from 0 alert
Brenk R. et al. 2008
ChemMedChem

Leadlikeness ⓘ

Leadlikeness:
implemented from
Teague SJ. 1999 Angew.
Chem. Int. Ed. No; 1 violation: XLOGP3>3.5
 $250 \leq MW \leq 350$
 $XLOGP \leq 3.5$
Num. rotatable bonds $\leq$
7

Synthetic accessibility ⓘ

Synthetic accessibility
score: from 1 (very
easy) to 10 (very
difficult)
based on 1024
fragmental contributions 3.22
(FP2) modulated by size
and complexity penalties,
trained on 12'782'590
molecules and tested on
40 external molecules
($r^2 = 0.94$)

Topological Polar Surface Area: <u>Calculated from Ertl P. et al. 2000 J. Med. Chem.</u>	43.37 Å ²	<u>scale</u> Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Soluble
Log $P_{o/w}$ (iLOGP) ⓘ	Lipophilicity	Log S (SILICOS-IT) ⓘ	
iLOGP: in-house physics-based method implemented from Daina A et al. 2014 J. Chem. Inf. Model.	2.50	SILICOS-IT: Fragmental method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT. http://www.silicos-it.com	-5.68
Log $P_{o/w}$ (XLOGP3) ⓘ		Solubility Class ⓘ	5.50e-04 mg/ml ; 2.08e-06 mol/l
XLOGP3: Atomistic and knowledge-based method calculated by XLOGP program, version 3.2.2, courtesy of CCBG, Shanghai Institute of Organic Chemistry	3.27	Solubility class: Log S scale Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Moderately soluble
Log $P_{o/w}$ (WLOGP) ⓘ			Pharmacokinetics
WLOGP: Atomistic method implemented from Wildman SA and Crippen GM. 1999 J. Chem. Inf. Model.	3.34	GI absorption ⓘ	
Log $P_{o/w}$ (MLOGP) ⓘ		Gastrointestinal absorption: according to the white of the BOILED-Egg	High
MLOGP: Topological method implemented from Moriguchi I. et al. 1992 Chem. Pharm. Bull. Moriguchi I. et al. 1994 Chem. Pharm. Bull. Lipinski PA. et al. 2001 Adv. Drug. Deliv. Rev.	1.92	BBB permeant ⓘ	
Log $P_{o/w}$ (SILICOS-IT) ⓘ		BBB permeation: according to the yolk of the BOILED-Egg	Yes
SILICOS-IT: Hybrid fragmental/topological method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT. http://www.silicos-it.com	3.85	P-gp substrate ⓘ	
Consensus Log $P_{o/w}$ ⓘ		P-glycoprotein substrate: SVM model built on 1033 molecules (training set) and tested on 415 molecules (test set) 10-fold CV: ACC=0.72 / AUC=0.77 External: ACC=0.88 / AUC=0.94	No
Consensus Log $P_{o/w}$:		CYP1A2 inhibitor ⓘ	
		Cytochrome P450 1A2 inhibitor: SVM model built on 9145 molecules (training set) and tested on 3000 molecules (test set)	Yes

Average of all five predictions

2.98

10-fold CV: ACC=0.83 / AUC=0.90
External: ACC=0.84 / AUC=0.91

CYP2C19 inhibitor ⓘ

Cytochrome P450 2C19 inhibitor: SVM model built on 9272 molecules (training set) and tested on 3000 molecules (test set) Yes
10-fold CV: ACC=0.80 / AUC=0.86
External: ACC=0.80 / AUC=0.87

CYP2C9 inhibitor ⓘ

Cytochrome P450 2C9 inhibitor: SVM model built on 5940 molecules (training set) and tested on 2075 molecules (test set) Yes
10-fold CV: ACC=0.78 / AUC=0.85
External: ACC=0.71 / AUC=0.81

CYP2D6 inhibitor ⓘ

Cytochrome P450 2D6 inhibitor: SVM model built on 3664 molecules (training set) and tested on 1068 molecules (test set) No
10-fold CV: ACC=0.79 / AUC=0.85
External: ACC=0.81 / AUC=0.87

CYP3A4 inhibitor ⓘ

Cytochrome P450 3A4 inhibitor: SVM model built on 7518 molecules (training set) and tested on 2579 molecules (test set) Yes
10-fold CV: ACC=0.77 / AUC=0.85
External: ACC=0.78 / AUC=0.86

Log K_p (skin permeation) ⓘ

Skin permeation: QSPR model implemented from -5.59 cm/s

Potts RO and Guy RH.
1992 Pharm. Res.

Druglikeness

Lipinski

Lipinski (Pfizer) filter:
implemented from
Lipinski CA. et al. 2001
Adv. Drug Deliv. Rev. Yes; 0 violation
MW $\leq$ 500
MLOGP $\leq$ 4.15
N or O $\leq$ 10
NH or OH $\leq$ 5

Ghose

Ghose filter:
implemented from
Ghose AK. et al. 1999 J.
Comb. Chem. Yes
160 $\leq$ MW $\leq$ 480
-0.4 $\leq$ WLOGP $\leq$ 5.6
40 $\leq$ MR $\leq$ 130
20 $\leq$ atoms $\leq$ 70

Veber

Veber (GSK) filter:
implemented from
Veber DF. et al. 2002 J. Yes
Med. Chem.
Rotatable bonds $\leq$ 10
TPSA $\leq$ 140

Egan

Egan (Pharmacia)
filter: implemented
from
Egan WJ. et al. 2000 J. Yes
Med. Chem.
WLOGP $\leq$ 5.88
TPSA $\leq$ 131.6

Muegge

Muegge (Bayer) filter:
implemented from
Muegge I. et al. 2001 J.
Med. Chem.
200 $\leq$ MW $\leq$ 600
-2 $\leq$ XLOGP $\leq$ 5
TPSA $\leq$ 150 Yes
Num. rings $\leq$ 7
Num. carbon $>$ 4
Num. heteroatoms $>$ 1
Num. rotatable bonds $\leq$
15
H-bond acc. $\leq$ 10
H-bond don. $\leq$ 5

Bioavailability Score ⓘ


Abbott Bioavailability
Score: Probability of F 0.56
> 10% in rat
implemented from
Martin YC. 2005 J.
Med. Chem.

Medicinal Chemistry

PAINS ⓘ

1 alert: quinone_A ⓘ

Pan Assay Interference
Structures:
implemented from
Baell JB. & Holloway
GA. 2010 J. Med.
Chem.



Brenk ⓘ


Structural Alert:
implemented from 0 alert
Brenk R. et al. 2008
ChemMedChem

Leadlikeness ⓘ

Leadlikeness:
implemented from
Teague SJ. 1999 Angew.
Chem. Int. Ed. Yes
 $250 \leq MW \leq 350$
 $XLOGP \leq 3.5$
Num. rotatable bonds $\leq$
7

Synthetic accessibility ⓘ

Synthetic accessibility
score: from 1 (very
easy) to 10 (very
difficult)
based on 1024
fragmental contributions 2.91
(FP2) modulated by size
and complexity penalties,
trained on 12'782'590
molecules and tested on
40 external molecules
($r^2 = 0.94$)



Enter a list of SMILES here:

Clc1ccc(cc1)OC3=CC(=O)c2ccccc2C3=O

Fill with an example

Clear



Run!

Show BOILED-Egg

Retrieve data:


 Powered by
ChemAxon

Molecule 1


 SMILES
S Clc1ccc(cc1)OC1=CC(=O)c2c(C1=O)cccc2

Physicochemical Properties

Formula	C16H9ClO3
Molecular weight	284.69 g/mol
Num. heavy atoms	20
Num. arom. heavy atoms	12
Fraction Csp3	0.00
Num. rotatable bonds	2
Num. H-bond acceptors	3
Num. H-bond donors	0
Molar Refractivity	75.27
TPSA	

Water Solubility

Log S (ESOL)

ESOL: Topological method implemented from
Delaney JS. 2004 J. Chem. Inf. Model.

-4.15

 Solubility
Class

2.03e-02 mg/ml ; 7.12e-05 mol/l

Solubility class: Log S scale

Insoluble < -10 < Poorly Moderately soluble
< -6 < Moderately < -4
< Soluble < -2 Very < 0
< Highly

Log S (Ali)

Ali: Topological method implemented from
Ali J. et al. 2012 J. Chem. Inf. Model.

-4.14

 Solubility
Class

2.08e-02 mg/ml ; 7.32e-05 mol/l

Solubility class: Log S

Topological Polar Surface Area: Calculated from Ertl P. et al. 2000 J. Med. Chem.	43.37 Å ²	<u>scale</u> Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Moderately soluble
Log <i>P</i>_{o/w} (iLOGP) ⓘ	Lipophilicity	Log <i>S</i> (SILICOS-IT) ⓘ	
iLOGP: in-house physics-based method implemented from Daina A et al. 2014 J. Chem. Inf. Model.	2.54	SILICOS-IT: Fragmental method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos-it.com	-5.90
Log <i>P</i>_{o/w} (XLOGP3) ⓘ		Solubility Class ⓘ	3.55e-04 mg/ml ; 1.25e-06 mol/l
XLOGP3: Atomistic and knowledge-based method calculated by XLOGP program, version 3.2.2, courtesy of CCBG, Shanghai Institute of Organic Chemistry	3.54	Solubility class: Log <i>S</i> <u>scale</u> Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Moderately soluble
Log <i>P</i>_{o/w} (WLOGP) ⓘ			Pharmacokinetics
WLOGP: Atomistic method implemented from Wildman SA and Crippen GM. 1999 J. Chem. Inf. Model.	3.68	GI absorption ⓘ	
Log <i>P</i>_{o/w} (MLOGP) ⓘ		Gastrointestinal absorption: according to the white of the BOILED-Egg	High
MLOGP: Topological method implemented from Moriguchi I. et al. 1992 Chem. Pharm. Bull. Moriguchi I. et al. 1994 Chem. Pharm. Bull. Lipinski PA. et al. 2001 Adv. Drug. Deliv. Rev.	2.19	BBB permeant ⓘ	
Log <i>P</i>_{o/w} (SILICOS-IT) ⓘ		BBB permeation: according to the yolk of the BOILED-Egg	Yes
SILICOS-IT: Hybrid fragmental/topological method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos-it.com	3.98	P-gp substrate ⓘ	
Consensus Log <i>P</i>_{o/w} ⓘ		P-glycoprotein substrate: SVM model built on 1033 molecules (training set) and tested on 415 molecules (test set) 10-fold CV: ACC=0.72 / AUC=0.77 External: ACC=0.88 / AUC=0.94	No
Consensus Log <i>P</i>_{o/w}:		CYP1A2 inhibitor ⓘ	
		Cytochrome P450 1A2 inhibitor: SVM model built on 9145 molecules (training set) and tested on 3000 molecules (test set)	Yes

Average of all five predictions

3.19

10-fold CV: ACC=0.83 / AUC=0.90
External: ACC=0.84 / AUC=0.91

CYP2C19 inhibitor ⓘ

Cytochrome P450 2C19 inhibitor: SVM model built on 9272 molecules (training set) and tested on 3000 molecules (test set) Yes
10-fold CV: ACC=0.80 / AUC=0.86
External: ACC=0.80 / AUC=0.87

CYP2C9 inhibitor ⓘ

Cytochrome P450 2C9 inhibitor: SVM model built on 5940 molecules (training set) and tested on 2075 molecules (test set) Yes
10-fold CV: ACC=0.78 / AUC=0.85
External: ACC=0.71 / AUC=0.81

CYP2D6 inhibitor ⓘ

Cytochrome P450 2D6 inhibitor: SVM model built on 3664 molecules (training set) and tested on 1068 molecules (test set) No
10-fold CV: ACC=0.79 / AUC=0.85
External: ACC=0.81 / AUC=0.87

CYP3A4 inhibitor ⓘ

Cytochrome P450 3A4 inhibitor: SVM model built on 7518 molecules (training set) and tested on 2579 molecules (test set) Yes
10-fold CV: ACC=0.77 / AUC=0.85
External: ACC=0.78 / AUC=0.86

Log K_p (skin permeation) ⓘ

Skin permeation: QSPR model implemented from -5.52 cm/s

Potts RO and Guy RH.
1992 Pharm. Res.

Druglikeness

Lipinski ⓘ

Lipinski (Pfizer) filter:
implemented from
Lipinski CA. et al. 2001
Adv. Drug Deliv. Rev. Yes; 0 violation
MW $\leq$ 500
MLOGP $\leq$ 4.15
N or O $\leq$ 10
NH or OH $\leq$ 5

Ghose ⓘ

Ghose filter:
implemented from
Ghose AK. et al. 1999 J.
Comb. Chem. Yes
160 $\leq$ MW $\leq$ 480
-0.4 $\leq$ WLOGP $\leq$ 5.6
40 $\leq$ MR $\leq$ 130
20 $\leq$ atoms $\leq$ 70

Veber ⓘ

Veber (GSK) filter:
implemented from
Veber DF. et al. 2002 J. Yes
Med. Chem.
Rotatable bonds $\leq$ 10
TPSA $\leq$ 140

Egan ⓘ

Egan (Pharmacia)
filter: implemented
from
Egan WJ. et al. 2000 J. Yes
Med. Chem.
WLOGP $\leq$ 5.88
TPSA $\leq$ 131.6

Muegge ⓘ

Muegge (Bayer) filter:
implemented from
Muegge I. et al. 2001 J.
Med. Chem.
200 $\leq$ MW $\leq$ 600
-2 $\leq$ XLOGP $\leq$ 5
TPSA $\leq$ 150 Yes
Num. rings $\leq$ 7
Num. carbon $>$ 4
Num. heteroatoms $>$ 1
Num. rotatable bonds $\leq$
15
H-bond acc. $\leq$ 10
H-bond don. $\leq$ 5

Bioavailability Score ⓘ


Abbott Bioavailability
Score: Probability of F 0.56
> 10% in rat
implemented from
Martin YC. 2005 J.
Med. Chem.

Medicinal Chemistry

PAINS ⓘ

1 alert: quinone_A ⓘ

Pan Assay Interference
Structures:
implemented from
Baell JB. & Holloway
GA. 2010 J. Med.
Chem.



Brenk ⓘ

Structural Alert:
implemented from 0 alert
Brenk R. et al. 2008
ChemMedChem

Leadlikeness ⓘ

Leadlikeness:
implemented from
Teague SJ. 1999 Angew.
Chem. Int. Ed. No; 1 violation: XLOGP3>3.5
 $250 \leq MW \leq 350$
 $XLOGP \leq 3.5$
Num. rotatable bonds $\leq$
7

Synthetic accessibility ⓘ

Synthetic accessibility
score: from 1 (very
easy) to 10 (very
difficult)
based on 1024
fragmental contributions 2.79
(FP2) modulated by size
and complexity penalties,
trained on 12'782'590
molecules and tested on
40 external molecules
($r^2 = 0.94$)

Topological Polar Surface Area: Calculated from Ertl P. et al. 2000 J. Med. Chem.	43.37 Å ²	<u>scale</u> Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Moderately soluble
Log $P_{o/w}$ (iLOGP) ⓘ		Log S (SILICOS-IT) ⓘ	
iLOGP: in-house physics-based method implemented from Daina A et al. 2014 J. Chem. Inf. Model.	2.65	SILICOS-IT: Fragmental method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos-it.com	-6.12
Log $P_{o/w}$ (XLOGP3) ⓘ		Solubility	2.51e-04 mg/ml ; 7.62e-07 mol/l
XLOGP3: Atomistic and knowledge-based method calculated by XLOGP program, version 3.2.2, courtesy of CCBG, Shanghai Institute of Organic Chemistry	3.60	Class ⓘ	
Log $P_{o/w}$ (WLOGP) ⓘ		Solubility class: Log S scale Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Poorly soluble
WLOGP: Atomistic method implemented from Wildman SA and Crippen GM. 1999 J. Chem. Inf. Model.	3.79		Pharmacokinetics
Log $P_{o/w}$ (MLOGP) ⓘ		GI absorption ⓘ	
MLOGP: Topological method implemented from Moriguchi I. et al. 1992 Chem. Pharm. Bull. Moriguchi I. et al. 1994 Chem. Pharm. Bull. Lipinski PA. et al. 2001 Adv. Drug. Deliv. Rev.	2.31	Gastrointestinal absorption: according to the white of the BOILED-Egg	High
Log $P_{o/w}$ (SILICOS-IT) ⓘ		BBB permeant ⓘ	
SILICOS-IT: Hybrid fragmental/topological method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos-it.com	4.01	BBB permeation: according to the yolk of the BOILED-Egg	Yes
Consensus Log $P_{o/w}$ ⓘ		P-gp substrate ⓘ	
Consensus Log $P_{o/w}$:		P-glycoprotein substrate: SVM model built on 1033 molecules (training set) and tested on 415 molecules (test set) 10-fold CV: ACC=0.72 / AUC=0.77 External: ACC=0.88 / AUC=0.94	No
		CYP1A2 inhibitor ⓘ	
		Cytochrome P450 1A2 inhibitor: SVM model built on 9145 molecules (training set) and tested on 3000 molecules (test set)	Yes

Average of all five predictions

3.27

10-fold CV: ACC=0.83 / AUC=0.90
External: ACC=0.84 / AUC=0.91

CYP2C19 inhibitor ⓘ

Cytochrome P450 2C19 inhibitor: SVM model built on 9272 molecules (training set) and tested on 3000 molecules (test set) Yes
10-fold CV: ACC=0.80 / AUC=0.86
External: ACC=0.80 / AUC=0.87

CYP2C9 inhibitor ⓘ

Cytochrome P450 2C9 inhibitor: SVM model built on 5940 molecules (training set) and tested on 2075 molecules (test set) Yes
10-fold CV: ACC=0.78 / AUC=0.85
External: ACC=0.71 / AUC=0.81

CYP2D6 inhibitor ⓘ

Cytochrome P450 2D6 inhibitor: SVM model built on 3664 molecules (training set) and tested on 1068 molecules (test set) No
10-fold CV: ACC=0.79 / AUC=0.85
External: ACC=0.81 / AUC=0.87

CYP3A4 inhibitor ⓘ

Cytochrome P450 3A4 inhibitor: SVM model built on 7518 molecules (training set) and tested on 2579 molecules (test set) Yes
10-fold CV: ACC=0.77 / AUC=0.85
External: ACC=0.78 / AUC=0.86

Log K_p (skin permeation) ⓘ

Skin permeation: QSPR model implemented from -5.75 cm/s

Potts RO and Guy RH.
1992 Pharm. Res.

Druglikeness

Lipinski

Lipinski (Pfizer) filter:
implemented from
Lipinski CA. et al. 2001
Adv. Drug Deliv. Rev. Yes; 0 violation
MW $\leq$ 500
MLOGP $\leq$ 4.15
N or O $\leq$ 10
NH or OH $\leq$ 5

Ghose

Ghose filter:
implemented from
Ghose AK. et al. 1999 J.
Comb. Chem. Yes
160 $\leq$ MW $\leq$ 480
-0.4 $\leq$ WLOGP $\leq$ 5.6
40 $\leq$ MR $\leq$ 130
20 $\leq$ atoms $\leq$ 70

Veber

Veber (GSK) filter:
implemented from
Veber DF. et al. 2002 J. Yes
Med. Chem.
Rotatable bonds $\leq$ 10
TPSA $\leq$ 140

Egan

Egan (Pharmacia)
filter: implemented
from
Egan WJ. et al. 2000 J. Yes
Med. Chem.
WLOGP $\leq$ 5.88
TPSA $\leq$ 131.6

Muegge

Muegge (Bayer) filter:
implemented from
Muegge I. et al. 2001 J.
Med. Chem.
200 $\leq$ MW $\leq$ 600
-2 $\leq$ XLOGP $\leq$ 5
TPSA $\leq$ 150 Yes
Num. rings $\leq$ 7
Num. carbon $>$ 4
Num. heteroatoms $>$ 1
Num. rotatable bonds $\leq$
15
H-bond acc. $\leq$ 10
H-bond don. $\leq$ 5

Bioavailability Score ⓘ


Abbott Bioavailability
Score: Probability of F 0.56
> 10% in rat
implemented from
Martin YC. 2005 J.
Med. Chem.

Medicinal Chemistry

PAINS ⓘ

1 alert: quinone_A ⓘ

Pan Assay Interference
Structures:
implemented from
Baell JB. & Holloway
GA. 2010 J. Med.
Chem.



Brenk ⓘ


Structural Alert:
implemented from 0 alert
Brenk R. et al. 2008
ChemMedChem

Leadlikeness ⓘ

Leadlikeness:
implemented from
Teague SJ. 1999 Angew.
Chem. Int. Ed. No; 1 violation: XLOGP3>3.5
 $250 \leq MW \leq 350$
 $XLOGP \leq 3.5$
Num. rotatable bonds $\leq$
7

Synthetic accessibility ⓘ

Synthetic accessibility
score: from 1 (very
easy) to 10 (very
difficult)
based on 1024
fragmental contributions 2.80
(FP2) modulated by size
and complexity penalties,
trained on 12'782'590
molecules and tested on
40 external molecules
($r^2 = 0.94$)



Enter a list of SMILES here:

O=C3C=C(Oc1ccccc1)C(=O)c2ccccc23

Fill with an example

Clear



Run!

Show BOILED-Egg

Retrieve data:


 Powered by
ChemAxon

Molecule 1


 SMILES
S O=C1C(=CC(=O)c2c1cccc2)Oc1ccccc1

Physicochemical Properties

Formula	C16H10O3
Molecular weight	250.25 g/mol
Num. heavy atoms	19
Num. arom. heavy atoms	12
Fraction Csp3	0.00
Num. rotatable bonds	2
Num. H-bond acceptors	3
Num. H-bond donors	0
Molar Refractivity	70.26
TPSA	

Water Solubility

Log S (ESOL)

ESOL: Topological method implemented from
Delaney JS. 2004 J. Chem. Inf. Model.

-3.56

 Solubility
Class

6.89e-02 mg/ml ; 2.75e-04 mol/l

Solubility class: Log S scale

Insoluble < -10 < Poorly Soluble
< -6 < Moderately < -4
< Soluble < -2 Very < 0
< Highly

Log S (Ali)

Ali: Topological method implemented from
Ali J. et al. 2012 J. Chem. Inf. Model.

-3.48

 Solubility
Class

8.25e-02 mg/ml ; 3.30e-04 mol/l

Solubility class: Log S

Topological Polar Surface Area: Calculated from Ertl P. et al. 2000 J. Med. Chem.	43.37 Å ²	<u>scale</u> Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Soluble
Log <i>P</i>_{o/w} (iLOGP) ⓘ	Lipophilicity	Log <i>S</i> (SILICOS-IT) ⓘ	
iLOGP: in-house physics-based method implemented from Daina A et al. 2014 J. Chem. Inf. Model.	2.25	SILICOS-IT: Fragmental method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos-it.com	-5.30
Log <i>P</i>_{o/w} (XLOGP3) ⓘ		Solubility Class ⓘ	1.26e-03 mg/ml ; 5.05e-06 mol/l
XLOGP3: Atomistic and knowledge-based method calculated by XLOGP program, version 3.2.2, courtesy of CCBG, Shanghai Institute of Organic Chemistry	2.91	Solubility class: Log <i>S</i> <u>scale</u> Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Moderately soluble
Log <i>P</i>_{o/w} (WLOGP) ⓘ			Pharmacokinetics
WLOGP: Atomistic method implemented from Wildman SA and Crippen GM. 1999 J. Chem. Inf. Model.	3.03	GI absorption ⓘ	
Log <i>P</i>_{o/w} (MLOGP) ⓘ		Gastrointestinal absorption: according to the white of the BOILED-Egg	High
MLOGP: Topological method implemented from Moriguchi I. et al. 1992 Chem. Pharm. Bull. Moriguchi I. et al. 1994 Chem. Pharm. Bull. Lipinski PA. et al. 2001 Adv. Drug. Deliv. Rev.	1.68	BBB permeant ⓘ	
Log <i>P</i>_{o/w} (SILICOS-IT) ⓘ		BBB permeation: according to the yolk of the BOILED-Egg	Yes
SILICOS-IT: Hybrid fragmental/topological method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos-it.com	3.36	P-gp substrate ⓘ	
Consensus Log <i>P</i>_{o/w} ⓘ		P-glycoprotein substrate: SVM model built on 1033 molecules (training set) and tested on 415 molecules (test set) 10-fold CV: ACC=0.72 / AUC=0.77 External: ACC=0.88 / AUC=0.94	No
Consensus Log <i>P</i>_{o/w}:		CYP1A2 inhibitor ⓘ	
		Cytochrome P450 1A2 inhibitor: SVM model built on 9145 molecules (training set) and tested on 3000 molecules (test set)	Yes

Average of all five
predictions

2.65

10-fold CV: ACC=0.83 /
AUC=0.90
External: ACC=0.84 /
AUC=0.91

CYP2C19 inhibitor ⓘ

Cytochrome P450
2C19 inhibitor: SVM
model built on 9272
molecules (training set)
and tested on 3000
molecules (test set) Yes
10-fold CV: ACC=0.80 /
AUC=0.86
External: ACC=0.80 /
AUC=0.87

CYP2C9 inhibitor ⓘ

Cytochrome P450 2C9
inhibitor: SVM model
built on 5940 molecules
(training set)
and tested on 2075
molecules (test set) No
10-fold CV: ACC=0.78 /
AUC=0.85
External: ACC=0.71 /
AUC=0.81

CYP2D6 inhibitor ⓘ

Cytochrome P450 2D6
inhibitor: SVM model
built on 3664 molecules
(training set)
and tested on 1068
molecules (test set) No
10-fold CV: ACC=0.79 /
AUC=0.85
External: ACC=0.81 /
AUC=0.87

CYP3A4 inhibitor ⓘ

Cytochrome P450 3A4
inhibitor: SVM model
built on 7518 molecules
(training set)
and tested on 2579
molecules (test set) No
10-fold CV: ACC=0.77 /
AUC=0.85
External: ACC=0.78 /
AUC=0.86

Log K_p (skin
permeation) ⓘ

Skin permeation:
QSPR model
implemented from -5.76 cm/s

Potts RO and Guy RH.
1992 Pharm. Res.

Druglikeness

Lipinski ⓘ

Lipinski (Pfizer) filter:
implemented from
Lipinski CA. et al. 2001
Adv. Drug Deliv. Rev. Yes; 0 violation
MW $\leq$ 500
MLOGP $\leq$ 4.15
N or O $\leq$ 10
NH or OH $\leq$ 5

Ghose ⓘ

Ghose filter:
implemented from
Ghose AK. et al. 1999 J.
Comb. Chem. Yes
160 $\leq$ MW $\leq$ 480
-0.4 $\leq$ WLOGP $\leq$ 5.6
40 $\leq$ MR $\leq$ 130
20 $\leq$ atoms $\leq$ 70

Veber ⓘ

Veber (GSK) filter:
implemented from
Veber DF. et al. 2002 J. Yes
Med. Chem.
Rotatable bonds $\leq$ 10
TPSA $\leq$ 140

Egan ⓘ

Egan (Pharmacia)
filter: implemented
from
Egan WJ. et al. 2000 J. Yes
Med. Chem.
WLOGP $\leq$ 5.88
TPSA $\leq$ 131.6

Muegge ⓘ

Muegge (Bayer) filter:
implemented from
Muegge I. et al. 2001 J.
Med. Chem.
200 $\leq$ MW $\leq$ 600
-2 $\leq$ XLOGP $\leq$ 5
TPSA $\leq$ 150 Yes
Num. rings $\leq$ 7
Num. carbon $>$ 4
Num. heteroatoms $>$ 1
Num. rotatable bonds $\leq$
15
H-bond acc. $\leq$ 10
H-bond don. $\leq$ 5

Bioavailability Score ⓘ


Abbott Bioavailability
Score: Probability of F 0.56
> 10% in rat
implemented from
Martin YC. 2005 J.
Med. Chem.

Medicinal Chemistry

PAINS ⓘ

1 alert: quinone_A ⓘ

Pan Assay Interference
Structures:
implemented from
Baell JB. & Holloway
GA. 2010 J. Med.
Chem.



Brenk ⓘ

Structural Alert:
implemented from 0 alert
Brenk R. et al. 2008
ChemMedChem

Leadlikeness ⓘ

Leadlikeness:
implemented from
Teague SJ. 1999 Angew.
Chem. Int. Ed. Yes
 $250 \leq MW \leq 350$
 $XLOGP \leq 3.5$
Num. rotatable bonds $\leq$
7

Synthetic accessibility ⓘ

Synthetic accessibility
score: from 1 (very
easy) to 10 (very
difficult)
based on 1024
fragmental contributions 2.81
(FP2) modulated by size
and complexity penalties,
trained on 12'782'590
molecules and tested on
40 external molecules
($r^2 = 0.94$)

Topological Polar Surface Area:	52.60 Å ²	<u>scale</u> Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Soluble
Calculated from Ertl P. et al. 2000 J. Med. Chem.			
	Lipophilicity	Log S (SILICOS-IT) ⓘ	
Log <i>P</i> _{o/w} (iLOGP) ⓘ		SILICOS-IT: Fragmental method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos-it.com	-5.42
<u>iLOGP: in-house physics-based method implemented from Daina A et al. 2014 J. Chem. Inf. Model.</u>	2.45		
Log <i>P</i> _{o/w} (XLOGP3) ⓘ		Solubility Class ⓘ	1.07e-03 mg/ml ; 3.83e-06 mol/l
<u>XLOGP3: Atomistic and knowledge-based method calculated by XLOGP program, version 3.2.2, courtesy of CCBG, Shanghai Institute of Organic Chemistry</u>	2.88	<u>Solubility class: Log S scale</u> Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Moderately soluble
Log <i>P</i> _{o/w} (WLOGP) ⓘ			Pharmacokinetics
<u>WLOGP: Atomistic method implemented from Wildman SA and Crippen GM. 1999 J. Chem. Inf. Model.</u>	3.04	GI absorption ⓘ <u>Gastrointestinal absorption: according to the white of the BOILED-Egg</u>	High
Log <i>P</i> _{o/w} (MLOGP) ⓘ		BBB permeant ⓘ	
<u>MLOGP: Topological method implemented from Moriguchi I. et al. 1992 Chem. Pharm. Bull. Moriguchi I. et al. 1994 Chem. Pharm. Bull. Lipinski PA. et al. 2001 Adv. Drug. Deliv. Rev.</u>	1.34	<u>BBB permeation: according to the yolk of the BOILED-Egg</u>	Yes
Log <i>P</i> _{o/w} (SILICOS-IT) ⓘ		P-gp substrate ⓘ	
<u>SILICOS-IT: Hybrid fragmental/topological method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos-it.com</u>	3.38	<u>P-glycoprotein substrate: SVM model built on 1033 molecules (training set) and tested on 415 molecules (test set) 10-fold CV: ACC=0.72 / AUC=0.77 External: ACC=0.88 / AUC=0.94</u>	No
Consensus Log <i>P</i> _{o/w} ⓘ		CYP1A2 inhibitor ⓘ	
<u>Consensus Log <i>P</i>_{o/w}:</u>		<u>Cytochrome P450 1A2 inhibitor: SVM model built on 9145 molecules (training set) and tested on 3000 molecules (test set)</u>	Yes

Average of all five predictions

2.62

10-fold CV: ACC=0.83 / AUC=0.90
External: ACC=0.84 / AUC=0.91

CYP2C19 inhibitor ⓘ

Cytochrome P450 2C19 inhibitor: SVM model built on 9272 molecules (training set) and tested on 3000 molecules (test set) Yes
10-fold CV: ACC=0.80 / AUC=0.86
External: ACC=0.80 / AUC=0.87

CYP2C9 inhibitor ⓘ

Cytochrome P450 2C9 inhibitor: SVM model built on 5940 molecules (training set) and tested on 2075 molecules (test set) Yes
10-fold CV: ACC=0.78 / AUC=0.85
External: ACC=0.71 / AUC=0.81

CYP2D6 inhibitor ⓘ

Cytochrome P450 2D6 inhibitor: SVM model built on 3664 molecules (training set) and tested on 1068 molecules (test set) No
10-fold CV: ACC=0.79 / AUC=0.85
External: ACC=0.81 / AUC=0.87

CYP3A4 inhibitor ⓘ

Cytochrome P450 3A4 inhibitor: SVM model built on 7518 molecules (training set) and tested on 2579 molecules (test set) Yes
10-fold CV: ACC=0.77 / AUC=0.85
External: ACC=0.78 / AUC=0.86

Log K_p (skin permeation) ⓘ

Skin permeation: QSPR model implemented from -5.96 cm/s

Potts RO and Guy RH.
1992 Pharm. Res.

Druglikeness

Lipinski ⓘ

Lipinski (Pfizer) filter:
implemented from
Lipinski CA. et al. 2001
Adv. Drug Deliv. Rev. Yes; 0 violation
MW $\leq$ 500
MLOGP $\leq$ 4.15
N or O $\leq$ 10
NH or OH $\leq$ 5

Ghose ⓘ

Ghose filter:
implemented from
Ghose AK. et al. 1999 J.
Comb. Chem. Yes
160 $\leq$ MW $\leq$ 480
-0.4 $\leq$ WLOGP $\leq$ 5.6
40 $\leq$ MR $\leq$ 130
20 $\leq$ atoms $\leq$ 70

Veber ⓘ

Veber (GSK) filter:
implemented from
Veber DF. et al. 2002 J. Yes
Med. Chem.
Rotatable bonds $\leq$ 10
TPSA $\leq$ 140

Egan ⓘ

Egan (Pharmacia)
filter: implemented
from
Egan WJ. et al. 2000 J. Yes
Med. Chem.
WLOGP $\leq$ 5.88
TPSA $\leq$ 131.6

Muegge ⓘ

Muegge (Bayer) filter:
implemented from
Muegge I. et al. 2001 J.
Med. Chem.
200 $\leq$ MW $\leq$ 600
-2 $\leq$ XLOGP $\leq$ 5
TPSA $\leq$ 150 Yes
Num. rings $\leq$ 7
Num. carbon $>$ 4
Num. heteroatoms $>$ 1
Num. rotatable bonds $\leq$
15
H-bond acc. $\leq$ 10
H-bond don. $\leq$ 5

Bioavailability Score ⓘ


Abbott Bioavailability
Score: Probability of F 0.56
> 10% in rat
implemented from
Martin YC. 2005 J.
Med. Chem.

Medicinal Chemistry

PAINS ⓘ

1 alert: quinone_A ⓘ

Pan Assay Interference
Structures:
implemented from
Baell JB. & Holloway
GA. 2010 J. Med.
Chem.



Brenk ⓘ


Structural Alert:
implemented from 0 alert
Brenk R. et al. 2008
ChemMedChem

Leadlikeness ⓘ

Leadlikeness:
implemented from
Teague SJ. 1999 Angew.
Chem. Int. Ed. Yes
 $250 \leq MW \leq 350$
 $XLOGP \leq 3.5$
Num. rotatable bonds $\leq$
7

Synthetic accessibility ⓘ

Synthetic accessibility
score: from 1 (very
easy) to 10 (very
difficult)
based on 1024
fragmental contributions 2.96
(FP2) modulated by size
and complexity penalties,
trained on 12'782'590
molecules and tested on
40 external molecules
($r^2 = 0.94$)



Enter a list of SMILES here:

```
[O-][N+](=O)c3ccccc3OC2=CC(=O)c1ccccc1C2=O
```

Fill with an example

Clear



Run!

Show BOILED-Egg

Retrieve data:


 Powered by
ChemAxon

Molecule 1


 SMILES
S O=C1C(=CC(=O)c2c1cccc2)Oc1ccccc1[N+](=O)[O-]

Physicochemical Properties

Formula	C16H9NO5
Molecular weight	295.25 g/mol
Num. heavy atoms	22
Num. arom. heavy atoms	12
Fraction Csp3	0.00
Num. rotatable bonds	3
Num. H-bond acceptors	5
Num. H-bond donors	0
Molar Refractivity	79.08
TPSA	

Water Solubility

Log S (ESOL)

ESOL: Topological method implemented from
Delaney JS. 2004 J. Chem. Inf. Model.

-3.60

 Solubility
Class

7.38e-02 mg/ml ; 2.50e-04 mol/l

Solubility class: Log S scale

Insoluble < -10 < Poorly Soluble
< -6 < Moderately < -4
< Soluble < -2 Very < 0
< Highly

Log S (Ali)

Ali: Topological method implemented from
Ali J. et al. 2012 J. Chem. Inf. Model.

-4.27

 Solubility
Class

1.59e-02 mg/ml ; 5.40e-05 mol/l

Solubility class: Log S

Topological Polar Surface Area: Calculated from Ertl P. et al. 2000 J. Med. Chem.	89.19 Å ²	<u>scale</u> Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Moderately soluble
Lipophilicity		Log S (SILICOS-IT) ⓘ	
Log $P_{o/w}$ (iLOGP) ⓘ		SILICOS-IT: Fragmental method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos-it.com	
iLOGP: in-house physics-based method implemented from Daina A et al. 2014 J. Chem. Inf. Model.	1.73		-4.66
Log $P_{o/w}$ (XLOGP3) ⓘ		Solubility Class ⓘ	6.40e-03 mg/ml ; 2.17e-05 mol/l
XLOGP3: Atomistic and knowledge-based method calculated by XLOGP program, version 3.2.2, courtesy of CCBG, Shanghai Institute of Organic Chemistry	2.74	Solubility class: Log S <u>scale</u> Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Moderately soluble
Log $P_{o/w}$ (WLOGP) ⓘ			Pharmacokinetics
WLOGP: Atomistic method implemented from Wildman SA and Crippen GM. 1999 J. Chem. Inf. Model.	2.94	GI absorption ⓘ	
Log $P_{o/w}$ (MLOGP) ⓘ		Gastrointestinal absorption: according to the white of the BOILED-Egg	High
MLOGP: Topological method implemented from Moriguchi I. et al. 1992 Chem. Pharm. Bull. Moriguchi I. et al. 1994 Chem. Pharm. Bull. Lipinski PA. et al. 2001 Adv. Drug. Deliv. Rev.	1.48	BBB permeant ⓘ	
Log $P_{o/w}$ (SILICOS-IT) ⓘ		BBB permeation: according to the yolk of the BOILED-Egg	No
SILICOS-IT: Hybrid fragmental/topological method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos-it.com	1.15	P-gp substrate ⓘ	
Consensus Log $P_{o/w}$ ⓘ		P-glycoprotein substrate: SVM model built on 1033 molecules (training set) and tested on 415 molecules (test set) 10-fold CV: ACC=0.72 / AUC=0.77 External: ACC=0.88 / AUC=0.94	No
Consensus Log $P_{o/w}$:		CYP1A2 inhibitor ⓘ	
		Cytochrome P450 1A2 inhibitor: SVM model built on 9145 molecules (training set) and tested on 3000 molecules (test set)	No

Average of all five predictions

2.01

10-fold CV: ACC=0.83 / AUC=0.90
External: ACC=0.84 / AUC=0.91

CYP2C19 inhibitor ⓘ

Cytochrome P450 2C19 inhibitor: SVM model built on 9272 molecules (training set) and tested on 3000 molecules (test set) Yes
10-fold CV: ACC=0.80 / AUC=0.86
External: ACC=0.80 / AUC=0.87

CYP2C9 inhibitor ⓘ

Cytochrome P450 2C9 inhibitor: SVM model built on 5940 molecules (training set) and tested on 2075 molecules (test set) Yes
10-fold CV: ACC=0.78 / AUC=0.85
External: ACC=0.71 / AUC=0.81

CYP2D6 inhibitor ⓘ

Cytochrome P450 2D6 inhibitor: SVM model built on 3664 molecules (training set) and tested on 1068 molecules (test set) No
10-fold CV: ACC=0.79 / AUC=0.85
External: ACC=0.81 / AUC=0.87

CYP3A4 inhibitor ⓘ

Cytochrome P450 3A4 inhibitor: SVM model built on 7518 molecules (training set) and tested on 2579 molecules (test set) No
10-fold CV: ACC=0.77 / AUC=0.85
External: ACC=0.78 / AUC=0.86

Log K_p (skin permeation) ⓘ

Skin permeation: QSPR model implemented from -6.16 cm/s

Potts RO and Guy RH.
1992 Pharm. Res.

Druglikeness

Lipinski ⓘ

Lipinski (Pfizer) filter:
implemented from
Lipinski CA. et al. 2001
Adv. Drug Deliv. Rev. Yes; 0 violation
MW $\leq$ 500
MLOGP $\leq$ 4.15
N or O $\leq$ 10
NH or OH $\leq$ 5

Ghose ⓘ

Ghose filter:
implemented from
Ghose AK. et al. 1999 J.
Comb. Chem. Yes
160 $\leq$ MW $\leq$ 480
-0.4 $\leq$ WLOGP $\leq$ 5.6
40 $\leq$ MR $\leq$ 130
20 $\leq$ atoms $\leq$ 70

Veber ⓘ

Veber (GSK) filter:
implemented from
Veber DF. et al. 2002 J. Yes
Med. Chem.
Rotatable bonds $\leq$ 10
TPSA $\leq$ 140

Egan ⓘ

Egan (Pharmacia)
filter: implemented
from
Egan WJ. et al. 2000 J. Yes
Med. Chem.
WLOGP $\leq$ 5.88
TPSA $\leq$ 131.6

Muegge ⓘ

Muegge (Bayer) filter:
implemented from
Muegge I. et al. 2001 J.
Med. Chem.
200 $\leq$ MW $\leq$ 600
-2 $\leq$ XLOGP $\leq$ 5
TPSA $\leq$ 150 Yes
Num. rings $\leq$ 7
Num. carbon $>$ 4
Num. heteroatoms $>$ 1
Num. rotatable bonds $\leq$
15
H-bond acc. $\leq$ 10
H-bond don. $\leq$ 5

Bioavailability Score ⓘ


Abbott Bioavailability Score: Probability of F
> 10% in rat
implemented from
Martin YC. 2005 J.
Med. Chem. 0.56

Medicinal Chemistry

PAINS ⓘ

1 alert: quinone_A ⓘ

Pan Assay Interference
Structures:
implemented from
Baell JB. & Holloway
GA. 2010 J. Med.
Chem.




quinone_A

2 alerts: nitro_group, oxygen-nitrogen_single_bond ⓘ

Brenk ⓘ

Structural Alert:
implemented from
Brenk R. et al. 2008
ChemMedChem



nitro_group



aliphatic, not in a ring

oxygen-nitrogen_single_bon

Leadlikeness ⓘ

Leadlikeness:
implemented from
Teague SJ. 1999 Angew.
Chem. Int. Ed. Yes
 $250 \leq MW \leq 350$
 $XLOGP \leq 3.5$
Num. rotatable bonds $\leq$
7

Synthetic accessibility ⓘ

Synthetic accessibility
score: from 1 (very
easy) to 10 (very
difficult)
based on 1024
fragmental contributions
(FP2) modulated by size 2.99
and complexity penalties,
trained on 12'782'590
molecules and tested on
40 external molecules
($r^2 = 0.94$)

Topological Polar Surface Area: Calculated from Ertl P. et al. 2000 J. Med. Chem.	43.37 Å ²	<u>scale</u> Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Soluble
Log <i>P</i>_{o/w} (iLOGP) ⓘ	Lipophilicity	Log <i>S</i> (SILICOS-IT) ⓘ	
iLOGP: in-house physics-based method implemented from Daina A et al. 2014 J. Chem. Inf. Model.	2.49	SILICOS-IT: Fragmental method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos-it.com	-5.68
Log <i>P</i>_{o/w} (XLOGP3) ⓘ		Solubility Class ⓘ	5.50e-04 mg/ml ; 2.08e-06 mol/l
XLOGP3: Atomistic and knowledge-based method calculated by XLOGP program, version 3.2.2, courtesy of CCBG, Shanghai Institute of Organic Chemistry	3.27	Solubility class: Log <i>S</i> scale Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Moderately soluble
Log <i>P</i>_{o/w} (WLOGP) ⓘ			Pharmacokinetics
WLOGP: Atomistic method implemented from Wildman SA and Crippen GM. 1999 J. Chem. Inf. Model.	3.34	GI absorption ⓘ	
Log <i>P</i>_{o/w} (MLOGP) ⓘ		Gastrointestinal absorption: according to the white of the BOILED-Egg	High
MLOGP: Topological method implemented from Moriguchi I. et al. 1992 Chem. Pharm. Bull. Moriguchi I. et al. 1994 Chem. Pharm. Bull. Lipinski PA. et al. 2001 Adv. Drug. Deliv. Rev.	1.92	BBB permeant ⓘ	
Log <i>P</i>_{o/w} (SILICOS-IT) ⓘ		BBB permeation: according to the yolk of the BOILED-Egg	Yes
SILICOS-IT: Hybrid fragmental/topological method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos-it.com	3.85	P-gp substrate ⓘ	
Consensus Log <i>P</i>_{o/w} ⓘ		P-glycoprotein substrate: SVM model built on 1033 molecules (training set) and tested on 415 molecules (test set) 10-fold CV: ACC=0.72 / AUC=0.77 External: ACC=0.88 / AUC=0.94	No
Consensus Log <i>P</i>_{o/w}:		CYP1A2 inhibitor ⓘ	
		Cytochrome P450 1A2 inhibitor: SVM model built on 9145 molecules (training set) and tested on 3000 molecules (test set)	Yes

Average of all five predictions

2.97

10-fold CV: ACC=0.83 / AUC=0.90
External: ACC=0.84 / AUC=0.91

CYP2C19 inhibitor ⓘ

Cytochrome P450 2C19 inhibitor: SVM model built on 9272 molecules (training set) and tested on 3000 molecules (test set) Yes
10-fold CV: ACC=0.80 / AUC=0.86
External: ACC=0.80 / AUC=0.87

CYP2C9 inhibitor ⓘ

Cytochrome P450 2C9 inhibitor: SVM model built on 5940 molecules (training set) and tested on 2075 molecules (test set) Yes
10-fold CV: ACC=0.78 / AUC=0.85
External: ACC=0.71 / AUC=0.81

CYP2D6 inhibitor ⓘ

Cytochrome P450 2D6 inhibitor: SVM model built on 3664 molecules (training set) and tested on 1068 molecules (test set) No
10-fold CV: ACC=0.79 / AUC=0.85
External: ACC=0.81 / AUC=0.87

CYP3A4 inhibitor ⓘ

Cytochrome P450 3A4 inhibitor: SVM model built on 7518 molecules (training set) and tested on 2579 molecules (test set) Yes
10-fold CV: ACC=0.77 / AUC=0.85
External: ACC=0.78 / AUC=0.86

Log K_p (skin permeation) ⓘ

Skin permeation: QSPR model implemented from -5.59 cm/s

Potts RO and Guy RH.
1992 Pharm. Res.

Druglikeness

Lipinski ⓘ

Lipinski (Pfizer) filter:
implemented from

Lipinski CA. et al. 2001

Adv. Drug Deliv. Rev.

Yes; 0 violation

MW $\leq$ 500

MLOGP $\leq$ 4.15

N or O $\leq$ 10

NH or OH $\leq$ 5

Ghose ⓘ

Ghose filter:

implemented from

Ghose AK. et al. 1999 J.

Comb. Chem.

Yes

160 $\leq$ MW $\leq$ 480

-0.4 $\leq$ WLOGP $\leq$ 5.6

40 $\leq$ MR $\leq$ 130

20 $\leq$ atoms $\leq$ 70

Veber ⓘ

Veber (GSK) filter:

implemented from

Veber DF. et al. 2002 J.

Yes

Med. Chem.

Rotatable bonds $\leq$ 10

TPSA $\leq$ 140

Egan ⓘ

Egan (Pharmacia)

filter: implemented

from

Egan WJ. et al. 2000 J.

Yes

Med. Chem.

WLOGP $\leq$ 5.88

TPSA $\leq$ 131.6

Muegge ⓘ

Muegge (Bayer) filter:

implemented from

Muegge I. et al. 2001 J.

Med. Chem.

200 $\leq$ MW $\leq$ 600

-2 $\leq$ XLOGP $\leq$ 5

TPSA $\leq$ 150

Yes

Num. rings $\leq$ 7

Num. carbon $>$ 4

Num. heteroatoms $>$ 1

Num. rotatable bonds $\leq$

15

H-bond acc. $\leq$ 10

H-bond don. $\leq$ 5


Bioavailability Score ⓘ

Abbott BioavailabilityScore: Probability of F 0.56
> 10% in ratimplemented from
Martin YC. 2005 J.
Med. Chem.

Medicinal Chemistry

PAINS ⓘ

1 alert: quinone_A ⓘ

Pan Assay InterferenceStructures:implemented from
Baell JB. & Holloway
GA. 2010 J. Med.
Chem.

Brenk ⓘ

Structural Alert:implemented from
Brenk R. et al. 2008
ChemMedChem

0 alert

Leadlikeness ⓘ


Leadlikeness:implemented from
Teague SJ. 1999 Angew.
Chem. Int. Ed.

Yes

 $250 \leq MW \leq 350$
 $XLOGP \leq 3.5$
Num. rotatable bonds $\leq$
7

Synthetic accessibility ⓘ

Synthetic accessibilityscore: from 1 (very
easy) to 10 (very
difficult)
based on 1024
fragmental contributions 2.99
(FP2) modulated by size
and complexity penalties,
trained on 12'782'590
molecules and tested on
40 external molecules
($r^2 = 0.94$)



Enter a list of SMILES here:

```
Clc3ccccc3OC2=CC(=O)c1ccccc1C2=O
```

Fill with an example

Clear



Run!

Show BOILED-Egg

Retrieve data:


 Powered by
ChemAxon

Molecule 1



SMILES
S Clc1ccccc1OC1=CC(=O)c2c(C1=O)cccc2

Physicochemical Properties

Formula	C16H9ClO3
Molecular weight	284.69 g/mol
Num. heavy atoms	20
Num. arom. heavy atoms	12
Fraction Csp3	0.00
Num. rotatable bonds	2
Num. H-bond acceptors	3
Num. H-bond donors	0
Molar Refractivity	75.27
TPSA	

Water Solubility

Log S (ESOL)

ESOL: Topological method implemented from Delaney JS. 2004 J. Chem. Inf. Model.

-4.15

Solubility
Class

2.03e-02 mg/ml ; 7.12e-05 mol/l

Solubility class: Log S scale

Insoluble < -10 < Poorly Moderately soluble
< -6 < Moderately < -4
< Soluble < -2 Very < 0
< Highly

Log S (Ali)

Ali: Topological method implemented from Ali J. et al. 2012 J. Chem. Inf. Model.

-4.14

Solubility
Class

2.08e-02 mg/ml ; 7.32e-05 mol/l

Solubility class: Log S

Topological Polar Surface Area: Calculated from Ertl P. et al. 2000 J. Med. Chem.	43.37 Å ²	<u>scale</u> Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Moderately soluble
Log <i>P</i>_{o/w} (iLOGP) ⓘ	Lipophilicity	Log <i>S</i> (SILICOS-IT) ⓘ	
iLOGP: in-house physics-based method implemented from Daina A et al. 2014 J. Chem. Inf. Model.	2.45	SILICOS-IT: Fragmental method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos- it.com	-5.90
Log <i>P</i>_{o/w} (XLOGP3) ⓘ		Solubility Class ⓘ	3.55e-04 mg/ml ; 1.25e-06 mol/l
XLOGP3: Atomistic and knowledge-based method calculated by XLOGP program, version 3.2.2, courtesy of CCBG, Shanghai Institute of Organic Chemistry	3.54	Solubility class: Log <i>S</i> scale Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Moderately soluble
Log <i>P</i>_{o/w} (WLOGP) ⓘ			Pharmacokinetics
WLOGP: Atomistic method implemented from Wildman SA and Crippen GM. 1999 J. Chem. Inf. Model.	3.68	GI absorption ⓘ	
Log <i>P</i>_{o/w} (MLOGP) ⓘ		Gastrointestinal absorption: according to the white of the BOILED-Egg	High
MLOGP: Topological method implemented from Moriguchi I. et al. 1992 Chem. Pharm. Bull. Moriguchi I. et al. 1994 Chem. Pharm. Bull. Lipinski PA. et al. 2001 Adv. Drug. Deliv. Rev.	2.19	BBB permeant ⓘ	
Log <i>P</i>_{o/w} (SILICOS-IT) ⓘ		BBB permeation: according to the yolk of the BOILED-Egg	Yes
SILICOS-IT: Hybrid fragmental/topological method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos- it.com	3.98	P-gp substrate ⓘ	
Consensus Log <i>P</i>_{o/w} ⓘ		P-glycoprotein substrate: SVM model built on 1033 molecules (training set) and tested on 415 molecules (test set) 10-fold CV: ACC=0.72 / AUC=0.77 External: ACC=0.88 / AUC=0.94	No
Consensus Log <i>P</i>_{o/w}:		CYP1A2 inhibitor ⓘ	
		Cytochrome P450 1A2 inhibitor: SVM model built on 9145 molecules (training set) and tested on 3000 molecules (test set)	Yes

Average of all five predictions

3.17

10-fold CV: ACC=0.83 / AUC=0.90
External: ACC=0.84 / AUC=0.91

CYP2C19 inhibitor ⓘ

Cytochrome P450 2C19 inhibitor: SVM model built on 9272 molecules (training set) and tested on 3000 molecules (test set) Yes
10-fold CV: ACC=0.80 / AUC=0.86
External: ACC=0.80 / AUC=0.87

CYP2C9 inhibitor ⓘ

Cytochrome P450 2C9 inhibitor: SVM model built on 5940 molecules (training set) and tested on 2075 molecules (test set) Yes
10-fold CV: ACC=0.78 / AUC=0.85
External: ACC=0.71 / AUC=0.81

CYP2D6 inhibitor ⓘ

Cytochrome P450 2D6 inhibitor: SVM model built on 3664 molecules (training set) and tested on 1068 molecules (test set) No
10-fold CV: ACC=0.79 / AUC=0.85
External: ACC=0.81 / AUC=0.87

CYP3A4 inhibitor ⓘ

Cytochrome P450 3A4 inhibitor: SVM model built on 7518 molecules (training set) and tested on 2579 molecules (test set) Yes
10-fold CV: ACC=0.77 / AUC=0.85
External: ACC=0.78 / AUC=0.86

Log K_p (skin permeation) ⓘ

Skin permeation: QSPR model implemented from -5.52 cm/s

Potts RO and Guy RH.
1992 Pharm. Res.

Druglikeness

Lipinski ⓘ

Lipinski (Pfizer) filter:
implemented from

Lipinski CA. et al. 2001

Adv. Drug Deliv. Rev.

Yes; 0 violation

MW $\leq$ 500

MLOGP $\leq$ 4.15

N or O $\leq$ 10

NH or OH $\leq$ 5

Ghose ⓘ

Ghose filter:

implemented from

Ghose AK. et al. 1999 J.

Comb. Chem.

Yes

160 $\leq$ MW $\leq$ 480

-0.4 $\leq$ WLOGP $\leq$ 5.6

40 $\leq$ MR $\leq$ 130

20 $\leq$ atoms $\leq$ 70

Veber ⓘ

Veber (GSK) filter:

implemented from

Veber DF. et al. 2002 J.

Yes

Med. Chem.

Rotatable bonds $\leq$ 10

TPSA $\leq$ 140

Egan ⓘ

Egan (Pharmacia)

filter: implemented

from

Egan WJ. et al. 2000 J.

Yes

Med. Chem.

WLOGP $\leq$ 5.88

TPSA $\leq$ 131.6

Muegge ⓘ

Muegge (Bayer) filter:

implemented from

Muegge I. et al. 2001 J.

Med. Chem.

200 $\leq$ MW $\leq$ 600

-2 $\leq$ XLOGP $\leq$ 5

TPSA $\leq$ 150

Yes

Num. rings $\leq$ 7

Num. carbon $>$ 4

Num. heteroatoms $>$ 1

Num. rotatable bonds $\leq$

15

H-bond acc. $\leq$ 10

H-bond don. $\leq$ 5

Bioavailability Score ⓘ


Abbott Bioavailability
Score: Probability of F 0.56
> 10% in rat
implemented from
Martin YC. 2005 J.
Med. Chem.

Medicinal Chemistry

PAINS ⓘ

1 alert: quinone_A ⓘ

Pan Assay Interference
Structures:
implemented from
Baell JB. & Holloway
GA. 2010 J. Med.
Chem.



Brenk ⓘ

Structural Alert:
implemented from 0 alert
Brenk R. et al. 2008
ChemMedChem

Leadlikeness ⓘ

Leadlikeness:
implemented from
Teague SJ. 1999 Angew.
Chem. Int. Ed. No; 1 violation: XLOGP3>3.5
 $250 \leq MW \leq 350$
 $XLOGP \leq 3.5$
Num. rotatable bonds $\leq$
7

Synthetic accessibility ⓘ

Synthetic accessibility
score: from 1 (very
easy) to 10 (very
difficult)
based on 1024
fragmental contributions 2.82
(FP2) modulated by size
and complexity penalties,
trained on 12'782'590
molecules and tested on
40 external molecules
($r^2 = 0.94$)

Topological Polar Surface Area:	43.37 Å ²	<u>scale</u> Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Moderately soluble
<u>Calculated from</u> <u>Ertl P. et al. 2000 J.</u> <u>Med. Chem.</u>			
	Lipophilicity	Log S (SILICOS-IT) ⓘ	
Log $P_{o/w}$ (iLOGP) ⓘ		SILICOS-IT: <u>Fragmental method</u> <u>calculated by</u> <u>FILTER-IT program,</u> <u>version 1.0.2, courtesy</u> <u>of SILICOS-IT,</u> <u>http://www.silicos-</u> <u>it.com</u>	-6.12
<u>iLOGP: in-house</u> <u>physics-based method</u> <u>implemented from</u> <u>Daina A et al. 2014 J.</u> <u>Chem. Inf. Model.</u>	2.56		
Log $P_{o/w}$ (XLOGP3) ⓘ		Solubility	2.51e-04 mg/ml ; 7.62e-07 mol/l
<u>XLOGP3: Atomistic</u> <u>and knowledge-based</u> <u>method calculated by</u> <u>XLOGP program,</u> <u>version 3.2.2, courtesy</u> <u>of CCBG, Shanghai</u> <u>Institute of Organic</u> <u>Chemistry</u>	3.60	Class ⓘ	
Log $P_{o/w}$ (WLOGP) ⓘ		<u>Solubility class: Log S</u> <u>scale</u> Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Poorly soluble
<u>WLOGP: Atomistic</u> <u>method implemented</u> <u>from</u> <u>Wildman SA and</u> <u>Crippen GM. 1999 J.</u> <u>Chem. Inf. Model.</u>	3.79		
Log $P_{o/w}$ (MLOGP) ⓘ			Pharmacokinetics
<u>MLOGP: Topological</u> <u>method implemented</u> <u>from</u> <u>Moriguchi I. et al. 1992</u> <u>Chem. Pharm. Bull.</u> <u>Moriguchi I. et al. 1994</u> <u>Chem. Pharm. Bull.</u> <u>Lipinski PA. et al. 2001</u> <u>Adv. Drug. Deliv. Rev.</u>	2.31	GI absorption ⓘ	
Log $P_{o/w}$ (SILICOS-IT) ⓘ		<u>Gastrointestinal</u> <u>absorption: according</u> <u>to the white of the</u> <u>BOILED-Egg</u>	High
<u>SILICOS-IT: Hybrid</u> <u>fragmental/topological</u> <u>method calculated by</u> <u>FILTER-IT program,</u> <u>version 1.0.2, courtesy</u> <u>of SILICOS-IT,</u> <u>http://www.silicos-</u> <u>it.com</u>	4.01	BBB permeant ⓘ	
Consensus Log $P_{o/w}$ ⓘ		<u>BBB permeation:</u> <u>according to the yolk of</u> <u>the BOILED-Egg</u>	Yes
<u>Consensus Log $P_{o/w}$:</u>		P-gp substrate ⓘ	
		<u>P-glycoprotein</u> <u>substrate: SVM model</u> <u>built on 1033 molecules</u> <u>(training set)</u> <u>and tested on 415</u> <u>molecules (test set)</u> <u>10-fold CV: ACC=0.72 /</u> <u>AUC=0.77</u> <u>External: ACC=0.88 /</u> <u>AUC=0.94</u>	No
		CYP1A2 inhibitor ⓘ	
		<u>Cytochrome P450 1A2</u> <u>inhibitor: SVM model</u> <u>built on 9145 molecules</u> <u>(training set)</u> <u>and tested on 3000</u> <u>molecules (test set)</u>	Yes

Average of all five predictions

3.26

10-fold CV: ACC=0.83 / AUC=0.90
External: ACC=0.84 / AUC=0.91

CYP2C19 inhibitor ⓘ

Cytochrome P450 2C19 inhibitor: SVM model built on 9272 molecules (training set) and tested on 3000 molecules (test set) Yes
10-fold CV: ACC=0.80 / AUC=0.86
External: ACC=0.80 / AUC=0.87

CYP2C9 inhibitor ⓘ

Cytochrome P450 2C9 inhibitor: SVM model built on 5940 molecules (training set) and tested on 2075 molecules (test set) Yes
10-fold CV: ACC=0.78 / AUC=0.85
External: ACC=0.71 / AUC=0.81

CYP2D6 inhibitor ⓘ

Cytochrome P450 2D6 inhibitor: SVM model built on 3664 molecules (training set) and tested on 1068 molecules (test set) No
10-fold CV: ACC=0.79 / AUC=0.85
External: ACC=0.81 / AUC=0.87

CYP3A4 inhibitor ⓘ

Cytochrome P450 3A4 inhibitor: SVM model built on 7518 molecules (training set) and tested on 2579 molecules (test set) Yes
10-fold CV: ACC=0.77 / AUC=0.85
External: ACC=0.78 / AUC=0.86

Log K_p (skin permeation) ⓘ

Skin permeation: QSPR model implemented from -5.75 cm/s

Potts RO and Guy RH.
1992 Pharm. Res.

Druglikeness

Lipinski ⓘ

Lipinski (Pfizer) filter:
implemented from
Lipinski CA. et al. 2001
Adv. Drug Deliv. Rev. Yes; 0 violation
MW $\leq$ 500
MLOGP $\leq$ 4.15
N or O $\leq$ 10
NH or OH $\leq$ 5

Ghose ⓘ

Ghose filter:
implemented from
Ghose AK. et al. 1999 J.
Comb. Chem. Yes
160 $\leq$ MW $\leq$ 480
-0.4 $\leq$ WLOGP $\leq$ 5.6
40 $\leq$ MR $\leq$ 130
20 $\leq$ atoms $\leq$ 70

Veber ⓘ

Veber (GSK) filter:
implemented from
Veber DF. et al. 2002 J. Yes
Med. Chem.
Rotatable bonds $\leq$ 10
TPSA $\leq$ 140

Egan ⓘ

Egan (Pharmacia)
filter: implemented
from
Egan WJ. et al. 2000 J. Yes
Med. Chem.
WLOGP $\leq$ 5.88
TPSA $\leq$ 131.6

Muegge ⓘ

Muegge (Bayer) filter:
implemented from
Muegge I. et al. 2001 J.
Med. Chem.
200 $\leq$ MW $\leq$ 600
-2 $\leq$ XLOGP $\leq$ 5
TPSA $\leq$ 150 Yes
Num. rings $\leq$ 7
Num. carbon $>$ 4
Num. heteroatoms $>$ 1
Num. rotatable bonds $\leq$
15
H-bond acc. $\leq$ 10
H-bond don. $\leq$ 5

Bioavailability Score ⓘ


Abbott Bioavailability
Score: Probability of F 0.56
> 10% in rat
implemented from
Martin YC. 2005 J.
Med. Chem.

Medicinal Chemistry

PAINS ⓘ

1 alert: quinone_A ⓘ

Pan Assay Interference
Structures:
implemented from
Baell JB. & Holloway
GA. 2010 J. Med.
Chem.



Brenk ⓘ


Structural Alert:
implemented from 0 alert
Brenk R. et al. 2008
ChemMedChem

Leadlikeness ⓘ

Leadlikeness:
implemented from
Teague SJ. 1999 Angew.
Chem. Int. Ed. No; 1 violation: XLOGP3>3.5
 $250 \leq MW \leq 350$
 $XLOGP \leq 3.5$
Num. rotatable bonds $\leq$
7

Synthetic accessibility ⓘ

Synthetic accessibility
score: from 1 (very
easy) to 10 (very
difficult)
based on 1024
fragmental contributions 2.81
(FP2) modulated by size
and complexity penalties,
trained on 12'782'590
molecules and tested on
40 external molecules
($r^2 = 0.94$)


[Show BOILED-Egg](#)



Enter a list of SMILES here:

COc1cccc(c1)OC3=CC(=O)c2ccccc2C3=O
[Fill with an example](#)
[Clear](#)
[Run!](#)

Retrieve data:


 Powered by
 ChemAxon

Molecule 1


 SMILES
COc1cccc(c1)OC1=CC(=O)c2c(C1=O)cccc2

Physicochemical Properties

Formula	C17H12O4
Molecular weight	280.27 g/mol
Num. heavy atoms	21
Num. arom. heavy atoms	12
Fraction Csp3	0.06
Num. rotatable bonds	3
Num. H-bond acceptors	4
Num. H-bond donors	0
Molar Refractivity	76.75
TPSA	

Water Solubility

Log S (ESOL)

ESOL: Topological method implemented from
Delaney JS. 2004 J. Chem. Inf. Model.

-3.62

 Solubility
 Class

6.77e-02 mg/ml ; 2.42e-04 mol/l

Solubility class: Log S scale

Insoluble < -10 < Poorly Soluble
< -6 < Moderately < -4
< Soluble < -2 Very < 0
< Highly

Log S (Ali)

Ali: Topological method implemented from
Ali J. et al. 2012 J. Chem. Inf. Model.

-3.64

 Solubility
 Class

6.36e-02 mg/ml ; 2.27e-04 mol/l

Solubility class: Log S

Topological Polar Surface Area:	52.60 Å ²	<u>scale</u> Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Soluble
Calculated from Ertl P. et al. 2000 J. Med. Chem.			
	Lipophilicity	Log S (SILICOS-IT) ⓘ	
Log <i>P</i> _{o/w} (iLOGP) ⓘ		SILICOS-IT: Fragmental method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos-it.com	-5.42
<u>iLOGP: in-house physics-based method implemented from Daina A et al. 2014 J. Chem. Inf. Model.</u>	2.56		
Log <i>P</i> _{o/w} (XLOGP3) ⓘ		Solubility Class ⓘ	1.07e-03 mg/ml ; 3.83e-06 mol/l
<u>XLOGP3: Atomistic and knowledge-based method calculated by XLOGP program, version 3.2.2, courtesy of CCBG, Shanghai Institute of Organic Chemistry</u>	2.88	<u>Solubility class: Log S scale</u> Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Moderately soluble
Log <i>P</i> _{o/w} (WLOGP) ⓘ			Pharmacokinetics
<u>WLOGP: Atomistic method implemented from Wildman SA and Crippen GM. 1999 J. Chem. Inf. Model.</u>	3.04	GI absorption ⓘ <u>Gastrointestinal absorption: according to the white of the BOILED-Egg</u>	High
Log <i>P</i> _{o/w} (MLOGP) ⓘ		BBB permeant ⓘ	
<u>MLOGP: Topological method implemented from Moriguchi I. et al. 1992 Chem. Pharm. Bull. Moriguchi I. et al. 1994 Chem. Pharm. Bull. Lipinski PA. et al. 2001 Adv. Drug. Deliv. Rev.</u>	1.34	<u>BBB permeation: according to the yolk of the BOILED-Egg</u>	Yes
Log <i>P</i> _{o/w} (SILICOS-IT) ⓘ		P-gp substrate ⓘ	
<u>SILICOS-IT: Hybrid fragmental/topological method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos-it.com</u>	3.38	<u>P-glycoprotein substrate: SVM model built on 1033 molecules (training set) and tested on 415 molecules (test set) 10-fold CV: ACC=0.72 / AUC=0.77 External: ACC=0.88 / AUC=0.94</u>	No
Consensus Log <i>P</i> _{o/w} ⓘ		CYP1A2 inhibitor ⓘ	
<u>Consensus Log <i>P</i>_{o/w}:</u>		<u>Cytochrome P450 1A2 inhibitor: SVM model built on 9145 molecules (training set) and tested on 3000 molecules (test set)</u>	Yes

Average of all five
predictions

2.64

10-fold CV: ACC=0.83 /
AUC=0.90
External: ACC=0.84 /
AUC=0.91

CYP2C19 inhibitor ⓘ

Cytochrome P450
2C19 inhibitor: SVM
model built on 9272
molecules (training set)
and tested on 3000
molecules (test set) Yes
10-fold CV: ACC=0.80 /
AUC=0.86
External: ACC=0.80 /
AUC=0.87

CYP2C9 inhibitor ⓘ

Cytochrome P450 2C9
inhibitor: SVM model
built on 5940 molecules
(training set)
and tested on 2075
molecules (test set) Yes
10-fold CV: ACC=0.78 /
AUC=0.85
External: ACC=0.71 /
AUC=0.81

CYP2D6 inhibitor ⓘ

Cytochrome P450 2D6
inhibitor: SVM model
built on 3664 molecules
(training set)
and tested on 1068
molecules (test set) No
10-fold CV: ACC=0.79 /
AUC=0.85
External: ACC=0.81 /
AUC=0.87

CYP3A4 inhibitor ⓘ

Cytochrome P450 3A4
inhibitor: SVM model
built on 7518 molecules
(training set)
and tested on 2579
molecules (test set) Yes
10-fold CV: ACC=0.77 /
AUC=0.85
External: ACC=0.78 /
AUC=0.86

Log K_p (skin
permeation) ⓘ

Skin permeation:
QSPR model -5.96 cm/s
implemented from

Potts RO and Guy RH.
1992 Pharm. Res.

Druglikeness

Lipinski ⓘ

Lipinski (Pfizer) filter:
implemented from

Lipinski CA. et al. 2001

Adv. Drug Deliv. Rev.

Yes; 0 violation

MW $\leq$ 500

MLOGP $\leq$ 4.15

N or O $\leq$ 10

NH or OH $\leq$ 5

Ghose ⓘ

Ghose filter:

implemented from

Ghose AK. et al. 1999 J.

Comb. Chem.

Yes

160 $\leq$ MW $\leq$ 480

-0.4 $\leq$ WLOGP $\leq$ 5.6

40 $\leq$ MR $\leq$ 130

20 $\leq$ atoms $\leq$ 70

Veber ⓘ

Veber (GSK) filter:

implemented from

Veber DF. et al. 2002 J.

Yes

Med. Chem.

Rotatable bonds $\leq$ 10

TPSA $\leq$ 140

Egan ⓘ

Egan (Pharmacia)

filter: implemented

from

Egan WJ. et al. 2000 J.

Yes

Med. Chem.

WLOGP $\leq$ 5.88

TPSA $\leq$ 131.6

Muegge ⓘ

Muegge (Bayer) filter:

implemented from

Muegge I. et al. 2001 J.

Med. Chem.

200 $\leq$ MW $\leq$ 600

-2 $\leq$ XLOGP $\leq$ 5

TPSA $\leq$ 150

Yes

Num. rings $\leq$ 7

Num. carbon $>$ 4

Num. heteroatoms $>$ 1

Num. rotatable bonds $\leq$

15

H-bond acc. $\leq$ 10

H-bond don. $\leq$ 5

Bioavailability Score ⓘ


Abbott Bioavailability
Score: Probability of F 0.56
> 10% in rat
implemented from
Martin YC. 2005 J.
Med. Chem.

Medicinal Chemistry

PAINS ⓘ

1 alert: quinone_A ⓘ

Pan Assay Interference
Structures:
implemented from
Baell JB. & Holloway
GA. 2010 J. Med.
Chem.



Brenk ⓘ


Structural Alert:
implemented from 0 alert
Brenk R. et al. 2008
ChemMedChem

Leadlikeness ⓘ

Leadlikeness:
implemented from
Teague SJ. 1999 Angew.
Chem. Int. Ed. Yes
 $250 \leq MW \leq 350$
 $XLOGP \leq 3.5$
Num. rotatable bonds $\leq$
7

Synthetic accessibility ⓘ

Synthetic accessibility
score: from 1 (very
easy) to 10 (very
difficult)
based on 1024
fragmental contributions 3.01
(FP2) modulated by size
and complexity penalties,
trained on 12'782'590
molecules and tested on
40 external molecules
($r^2 = 0.94$)





Enter a list of SMILES here:

Retrieve data:


 Powered by
ChemAxon

Molecule 1


 SMILES
O=C1C(=CC(=O)c2c1cccc2)Oc1cccc(c1)[N+](=O)[O-]

Physicochemical Properties

Formula	C16H9NO5
Molecular weight	295.25 g/mol
Num. heavy atoms	22
Num. arom. heavy atoms	12
Fraction Csp3	0.00
Num. rotatable bonds	3
Num. H-bond acceptors	5
Num. H-bond donors	0
Molar Refractivity	79.08
TPSA	

Water Solubility

Log S (ESOL)

ESOL: Topological method implemented from
Delaney JS. 2004 J. Chem. Inf. Model.

-3.60

 Solubility
 Class

7.38e-02 mg/ml ; 2.50e-04 mol/l

Solubility class: Log S scale

Insoluble < -10 < Poorly Soluble
< -6 < Moderately < -4
< Soluble < -2 Very < 0
< Highly

Log S (Ali)

Ali: Topological method implemented from
Ali J. et al. 2012 J. Chem. Inf. Model.

-4.27

 Solubility
 Class

1.59e-02 mg/ml ; 5.40e-05 mol/l

Solubility class: Log S

Topological Polar**Surface Area:**89.19 Å²

Calculated from

Ertl P. et al. 2000 J.

Med. Chem.

scaleInsoluble < -10 < Poorly

Moderately soluble

< -6 < Moderately < -4< Soluble < -2 Very < 0< Highly**Lipophilicity**Log $P_{o/w}$ (iLOGP) iLOGP: in-house
physics-based method
implemented from

2.01

Daina A et al. 2014 J.

Chem. Inf. Model.

Log S (SILICOS-IT) SILICOS-IT:Fragmental methodcalculated byFILTER-IT program,

-4.66

version 1.0.2, courtesyof SILICOS-IT,[\[Log \\$P_{o/w}\\$ \\(XLOGP3\\) !\\[\\]\\(e03857cdd33a5ff23dbb9f5eebaa4497_img.jpg\\)\]\(http://www.silicos-</u></div>
<div data-bbox=\)](http://www.silicos-</u></div>
<div data-bbox=)XLOGP3: Atomistic
and knowledge-based
method calculated by

2.74

XLOGP program,version 3.2.2, courtesyof CCBG, ShanghaiInstitute of OrganicChemistry

Solubility

6.40e-03 mg/ml ; 2.17e-05 mol/l

Class Solubility class: Log S scaleInsoluble < -10 < Poorly

Moderately soluble

< -6 < Moderately < -4< Soluble < -2 Very < 0< HighlyLog $P_{o/w}$ (WLOGP) WLOGP: Atomistic
method implemented
from

2.94

Wildman SA andCrippen GM. 1999 J.Chem. Inf. Model.GI absorption **Pharmacokinetics**Gastrointestinalabsorption: according

High

to the white of theBOILED-EggLog $P_{o/w}$ (MLOGP) MLOGP: Topological
method implemented
from

1.48

Moriguchi I. et al. 1992Chem. Pharm. Bull.Moriguchi I. et al. 1994Chem. Pharm. Bull.Lipinski PA. et al. 2001Adv. Drug. Deliv. Rev.BBB permeant BBB permeation:according to the yolk of

No

the BOILED-EggP-gp substrate P-glycoproteinsubstrate: SVM modelbuilt on 1033 molecules(training set)and tested on 415

No

molecules (test set)10-fold CV: ACC=0.72 /AUC=0.77External: ACC=0.88 /AUC=0.94Log $P_{o/w}$ (SILICOS-IT)SILICOS-IT: Hybrid
fragmental/topological
method calculated by

1.15

FILTER-IT program,version 1.0.2, courtesyof SILICOS-IT,[\[Consensus Log \\$P_{o/w}\\$!\\[\\]\\(a476851df51cd68478e3e6f64937c30c_img.jpg\\)\]\(http://www.silicos-</u></div>
<div data-bbox=\)](http://www.silicos-</u></div>
<div data-bbox=)Consensus Log $P_{o/w}$:CYP1A2 inhibitor Cytochrome P450 1A2inhibitor: SVM modelbuilt on 9145 molecules(training set)and tested on 3000

Yes

molecules (test set)

Average of all five predictions

2.06

10-fold CV: ACC=0.83 / AUC=0.90
External: ACC=0.84 / AUC=0.91

CYP2C19 inhibitor ⓘ

Cytochrome P450 2C19 inhibitor: SVM model built on 9272 molecules (training set) and tested on 3000 molecules (test set) Yes
10-fold CV: ACC=0.80 / AUC=0.86
External: ACC=0.80 / AUC=0.87

CYP2C9 inhibitor ⓘ

Cytochrome P450 2C9 inhibitor: SVM model built on 5940 molecules (training set) and tested on 2075 molecules (test set) Yes
10-fold CV: ACC=0.78 / AUC=0.85
External: ACC=0.71 / AUC=0.81

CYP2D6 inhibitor ⓘ

Cytochrome P450 2D6 inhibitor: SVM model built on 3664 molecules (training set) and tested on 1068 molecules (test set) No
10-fold CV: ACC=0.79 / AUC=0.85
External: ACC=0.81 / AUC=0.87

CYP3A4 inhibitor ⓘ

Cytochrome P450 3A4 inhibitor: SVM model built on 7518 molecules (training set) and tested on 2579 molecules (test set) No
10-fold CV: ACC=0.77 / AUC=0.85
External: ACC=0.78 / AUC=0.86

Log K_p (skin permeation) ⓘ

Skin permeation: QSPR model implemented from -6.16 cm/s

Potts RO and Guy RH.
1992 Pharm. Res.

Druglikeness

Lipinski ⓘ

Lipinski (Pfizer) filter:
implemented from
Lipinski CA. et al. 2001
Adv. Drug Deliv. Rev. Yes; 0 violation
MW $\leq$ 500
MLOGP $\leq$ 4.15
N or O $\leq$ 10
NH or OH $\leq$ 5

Ghose ⓘ

Ghose filter:
implemented from
Ghose AK. et al. 1999 J.
Comb. Chem. Yes
160 $\leq$ MW $\leq$ 480
-0.4 $\leq$ WLOGP $\leq$ 5.6
40 $\leq$ MR $\leq$ 130
20 $\leq$ atoms $\leq$ 70

Veber ⓘ

Veber (GSK) filter:
implemented from
Veber DF. et al. 2002 J. Yes
Med. Chem.
Rotatable bonds $\leq$ 10
TPSA $\leq$ 140

Egan ⓘ

Egan (Pharmacia)
filter: implemented
from
Egan WJ. et al. 2000 J. Yes
Med. Chem.
WLOGP $\leq$ 5.88
TPSA $\leq$ 131.6

Muegge ⓘ

Muegge (Bayer) filter:
implemented from
Muegge I. et al. 2001 J.
Med. Chem.
200 $\leq$ MW $\leq$ 600
-2 $\leq$ XLOGP $\leq$ 5
TPSA $\leq$ 150 Yes
Num. rings $\leq$ 7
Num. carbon $>$ 4
Num. heteroatoms $>$ 1
Num. rotatable bonds $\leq$
15
H-bond acc. $\leq$ 10
H-bond don. $\leq$ 5

Bioavailability Score ⓘ


Abbott Bioavailability
Score: Probability of F 0.56
> 10% in rat
implemented from
Martin YC. 2005 J.
Med. Chem.

Medicinal Chemistry

PAINS ⓘ

1 alert: quinone_A ⓘ

Pan Assay Interference
Structures:
implemented from
Baell JB. & Holloway
GA. 2010 J. Med.
Chem.




quinone_A


2 alerts: nitro_group, oxygen-nitrogen_single_bond ⓘ

Brenk ⓘ

Structural Alert:
implemented from
Brenk R. et al. 2008
ChemMedChem



nitro_group



aliphatic, not in a ring

oxygen-nitrogen_single_bon

Leadlikeness ⓘ

Leadlikeness:
implemented from
Teague SJ. 1999 Angew.
Chem. Int. Ed. Yes
 $250 \leq MW \leq 350$
 $XLOGP \leq 3.5$
Num. rotatable bonds $\leq$
7

Synthetic accessibility ⓘ

Synthetic accessibility
score: from 1 (very
easy) to 10 (very
difficult)
based on 1024
fragmental contributions 3.01
(FP2) modulated by size
and complexity penalties,
trained on 12'782'590
molecules and tested on
40 external molecules
($r^2 = 0.94$)

Topological Polar Surface Area: Calculated from Ertl P. et al. 2000 J. Med. Chem.	43.37 Å ²	<u>scale</u> Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Soluble
Log <i>P</i>_{o/w} (iLOGP) ⓘ	Lipophilicity	Log <i>S</i> (SILICOS-IT) ⓘ	
iLOGP: in-house physics-based method implemented from Daina A et al. 2014 J. Chem. Inf. Model.	2.52	SILICOS-IT: Fragmental method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos-it.com	-5.68
Log <i>P</i>_{o/w} (XLOGP3) ⓘ		Solubility Class ⓘ	5.50e-04 mg/ml ; 2.08e-06 mol/l
XLOGP3: Atomistic and knowledge-based method calculated by XLOGP program, version 3.2.2, courtesy of CCBG, Shanghai Institute of Organic Chemistry	3.27	Solubility class: Log <i>S</i> <u>scale</u> Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Moderately soluble
Log <i>P</i>_{o/w} (WLOGP) ⓘ			Pharmacokinetics
WLOGP: Atomistic method implemented from Wildman SA and Crippen GM. 1999 J. Chem. Inf. Model.	3.34	GI absorption ⓘ	
Log <i>P</i>_{o/w} (MLOGP) ⓘ		Gastrointestinal absorption: according to the white of the BOILED-Egg	High
MLOGP: Topological method implemented from Moriguchi I. et al. 1992 Chem. Pharm. Bull. Moriguchi I. et al. 1994 Chem. Pharm. Bull. Lipinski PA. et al. 2001 Adv. Drug. Deliv. Rev.	1.92	BBB permeant ⓘ	
Log <i>P</i>_{o/w} (SILICOS-IT) ⓘ		BBB permeation: according to the yolk of the BOILED-Egg	Yes
SILICOS-IT: Hybrid fragmental/topological method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos-it.com	3.85	P-gp substrate ⓘ	
Consensus Log <i>P</i>_{o/w} ⓘ		P-glycoprotein substrate: SVM model built on 1033 molecules (training set) and tested on 415 molecules (test set) 10-fold CV: ACC=0.72 / AUC=0.77 External: ACC=0.88 / AUC=0.94	No
Consensus Log <i>P</i>_{o/w}:		CYP1A2 inhibitor ⓘ	
		Cytochrome P450 1A2 inhibitor: SVM model built on 9145 molecules (training set) and tested on 3000 molecules (test set)	Yes

Average of all five predictions

2.98

10-fold CV: ACC=0.83 / AUC=0.90
External: ACC=0.84 / AUC=0.91

CYP2C19 inhibitor ⓘ

Cytochrome P450 2C19 inhibitor: SVM model built on 9272 molecules (training set) and tested on 3000 molecules (test set) Yes
10-fold CV: ACC=0.80 / AUC=0.86
External: ACC=0.80 / AUC=0.87

CYP2C9 inhibitor ⓘ

Cytochrome P450 2C9 inhibitor: SVM model built on 5940 molecules (training set) and tested on 2075 molecules (test set) Yes
10-fold CV: ACC=0.78 / AUC=0.85
External: ACC=0.71 / AUC=0.81

CYP2D6 inhibitor ⓘ

Cytochrome P450 2D6 inhibitor: SVM model built on 3664 molecules (training set) and tested on 1068 molecules (test set) No
10-fold CV: ACC=0.79 / AUC=0.85
External: ACC=0.81 / AUC=0.87

CYP3A4 inhibitor ⓘ

Cytochrome P450 3A4 inhibitor: SVM model built on 7518 molecules (training set) and tested on 2579 molecules (test set) Yes
10-fold CV: ACC=0.77 / AUC=0.85
External: ACC=0.78 / AUC=0.86

Log K_p (skin permeation) ⓘ

Skin permeation: QSPR model implemented from -5.59 cm/s

Potts RO and Guy RH.
1992 Pharm. Res.

Druglikeness

Lipinski ⓘ

Lipinski (Pfizer) filter:
implemented from

Lipinski CA. et al. 2001

Adv. Drug Deliv. Rev.

Yes; 0 violation

MW $\leq$ 500

MLOGP $\leq$ 4.15

N or O $\leq$ 10

NH or OH $\leq$ 5

Ghose ⓘ

Ghose filter:

implemented from

Ghose AK. et al. 1999 J.

Comb. Chem.

Yes

160 $\leq$ MW $\leq$ 480

-0.4 $\leq$ WLOGP $\leq$ 5.6

40 $\leq$ MR $\leq$ 130

20 $\leq$ atoms $\leq$ 70

Veber ⓘ

Veber (GSK) filter:

implemented from

Veber DF. et al. 2002 J.

Yes

Med. Chem.

Rotatable bonds $\leq$ 10

TPSA $\leq$ 140

Egan ⓘ

Egan (Pharmacia)

filter: implemented

from

Egan WJ. et al. 2000 J.

Yes

Med. Chem.

WLOGP $\leq$ 5.88

TPSA $\leq$ 131.6

Muegge ⓘ

Muegge (Bayer) filter:

implemented from

Muegge I. et al. 2001 J.

Med. Chem.

200 $\leq$ MW $\leq$ 600

-2 $\leq$ XLOGP $\leq$ 5

TPSA $\leq$ 150

Yes

Num. rings $\leq$ 7

Num. carbon $>$ 4

Num. heteroatoms $>$ 1

Num. rotatable bonds $\leq$

15

H-bond acc. $\leq$ 10

H-bond don. $\leq$ 5

Bioavailability Score ⓘ


Abbott Bioavailability
Score: Probability of F 0.56
> 10% in rat
implemented from
Martin YC. 2005 J.
Med. Chem.

Medicinal Chemistry

PAINS ⓘ

1 alert: quinone_A ⓘ

Pan Assay Interference
Structures:
implemented from
Baell JB. & Holloway
GA. 2010 J. Med.
Chem.



Brenk ⓘ


Structural Alert:
implemented from 0 alert
Brenk R. et al. 2008
ChemMedChem

Leadlikeness ⓘ

Leadlikeness:
implemented from
Teague SJ. 1999 Angew.
Chem. Int. Ed. Yes
 $250 \leq MW \leq 350$
 $XLOGP \leq 3.5$
Num. rotatable bonds $\leq$
7

Synthetic accessibility ⓘ

Synthetic accessibility
score: from 1 (very
easy) to 10 (very
difficult)
based on 1024
fragmental contributions 2.90
(FP2) modulated by size
and complexity penalties,
trained on 12'782'590
molecules and tested on
40 external molecules
($r^2 = 0.94$)



Enter a list of SMILES here:

```
Clc1cccc(c1)OC3=CC(=O)c2ccccc2C3=O
```

Fill with an example

Clear



Run!

Show BOILED-Egg

Retrieve data:


 Powered by
ChemAxon

Molecule 1



SMILES
S Clc1cccc(c1)OC1=CC(=O)c2c(C1=O)cccc2

Physicochemical Properties

Formula	C16H9ClO3
Molecular weight	284.69 g/mol
Num. heavy atoms	20
Num. arom. heavy atoms	12
Fraction Csp3	0.00
Num. rotatable bonds	2
Num. H-bond acceptors	3
Num. H-bond donors	0
Molar Refractivity	75.27
TPSA	

Water Solubility

Log S (ESOL)

ESOL: Topological method implemented from Delaney JS. 2004 J. Chem. Inf. Model.

-4.15

Solubility
Class

2.03e-02 mg/ml ; 7.12e-05 mol/l

Solubility class: Log S scale

Insoluble < -10 < Poorly Moderately soluble
< -6 < Moderately < -4
< Soluble < -2 Very < 0
< Highly

Log S (Ali)

Ali: Topological method implemented from Ali J. et al. 2012 J. Chem. Inf. Model.

-4.14

Solubility
Class

2.08e-02 mg/ml ; 7.32e-05 mol/l

Solubility class: Log S

Topological Polar Surface Area: Calculated from Ertl P. et al. 2000 J. Med. Chem.	43.37 Å ²	<u>scale</u> Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Moderately soluble
Lipophilicity		Log S (SILICOS-IT) ⓘ	
Log $P_{o/w}$ (iLOGP) ⓘ		SILICOS-IT: Fragmental method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos-it.com	
iLOGP: in-house physics-based method implemented from Daina A et al. 2014 J. Chem. Inf. Model.	2.53		-5.90
Log $P_{o/w}$ (XLOGP3) ⓘ		Solubility Class ⓘ	3.55e-04 mg/ml ; 1.25e-06 mol/l
XLOGP3: Atomistic and knowledge-based method calculated by XLOGP program, version 3.2.2, courtesy of CCBG, Shanghai Institute of Organic Chemistry	3.54	Solubility class: Log S <u>scale</u> Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Moderately soluble
Log $P_{o/w}$ (WLOGP) ⓘ			Pharmacokinetics
WLOGP: Atomistic method implemented from Wildman SA and Crippen GM. 1999 J. Chem. Inf. Model.	3.68	GI absorption ⓘ	
Log $P_{o/w}$ (MLOGP) ⓘ		Gastrointestinal absorption: according to the white of the BOILED-Egg	High
MLOGP: Topological method implemented from Moriguchi I. et al. 1992 Chem. Pharm. Bull. Moriguchi I. et al. 1994 Chem. Pharm. Bull. Lipinski PA. et al. 2001 Adv. Drug. Deliv. Rev.	2.19	BBB permeant ⓘ	
Log $P_{o/w}$ (SILICOS-IT) ⓘ		BBB permeation: according to the yolk of the BOILED-Egg	Yes
SILICOS-IT: Hybrid fragmental/topological method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos-it.com	3.98	P-gp substrate ⓘ	
Consensus Log $P_{o/w}$ ⓘ		P-glycoprotein substrate: SVM model built on 1033 molecules (training set) and tested on 415 molecules (test set) 10-fold CV: ACC=0.72 / AUC=0.77 External: ACC=0.88 / AUC=0.94	No
Consensus Log $P_{o/w}$:		CYP1A2 inhibitor ⓘ	
		Cytochrome P450 1A2 inhibitor: SVM model built on 9145 molecules (training set) and tested on 3000 molecules (test set)	Yes

Average of all five predictions

3.18

10-fold CV: ACC=0.83 / AUC=0.90
External: ACC=0.84 / AUC=0.91

CYP2C19 inhibitor ⓘ

Cytochrome P450 2C19 inhibitor: SVM model built on 9272 molecules (training set) and tested on 3000 molecules (test set) Yes
10-fold CV: ACC=0.80 / AUC=0.86
External: ACC=0.80 / AUC=0.87

CYP2C9 inhibitor ⓘ

Cytochrome P450 2C9 inhibitor: SVM model built on 5940 molecules (training set) and tested on 2075 molecules (test set) Yes
10-fold CV: ACC=0.78 / AUC=0.85
External: ACC=0.71 / AUC=0.81

CYP2D6 inhibitor ⓘ

Cytochrome P450 2D6 inhibitor: SVM model built on 3664 molecules (training set) and tested on 1068 molecules (test set) No
10-fold CV: ACC=0.79 / AUC=0.85
External: ACC=0.81 / AUC=0.87

CYP3A4 inhibitor ⓘ

Cytochrome P450 3A4 inhibitor: SVM model built on 7518 molecules (training set) and tested on 2579 molecules (test set) Yes
10-fold CV: ACC=0.77 / AUC=0.85
External: ACC=0.78 / AUC=0.86

Log K_p (skin permeation) ⓘ

Skin permeation: QSPR model implemented from -5.52 cm/s

Potts RO and Guy RH.
1992 Pharm. Res.

Druglikeness

Lipinski ⓘ

Lipinski (Pfizer) filter:
implemented from
Lipinski CA. et al. 2001
Adv. Drug Deliv. Rev. Yes; 0 violation
MW $\leq$ 500
MLOGP $\leq$ 4.15
N or O $\leq$ 10
NH or OH $\leq$ 5

Ghose ⓘ

Ghose filter:
implemented from
Ghose AK. et al. 1999 J.
Comb. Chem. Yes
160 $\leq$ MW $\leq$ 480
-0.4 $\leq$ WLOGP $\leq$ 5.6
40 $\leq$ MR $\leq$ 130
20 $\leq$ atoms $\leq$ 70

Veber ⓘ

Veber (GSK) filter:
implemented from
Veber DF. et al. 2002 J. Yes
Med. Chem.
Rotatable bonds $\leq$ 10
TPSA $\leq$ 140

Egan ⓘ

Egan (Pharmacia)
filter: implemented
from
Egan WJ. et al. 2000 J. Yes
Med. Chem.
WLOGP $\leq$ 5.88
TPSA $\leq$ 131.6

Muegge ⓘ

Muegge (Bayer) filter:
implemented from
Muegge I. et al. 2001 J.
Med. Chem.
200 $\leq$ MW $\leq$ 600
-2 $\leq$ XLOGP $\leq$ 5
TPSA $\leq$ 150 Yes
Num. rings $\leq$ 7
Num. carbon $>$ 4
Num. heteroatoms $>$ 1
Num. rotatable bonds $\leq$
15
H-bond acc. $\leq$ 10
H-bond don. $\leq$ 5

Bioavailability Score ⓘ


Abbott Bioavailability
Score: Probability of F 0.56
> 10% in rat
implemented from
Martin YC. 2005 J.
Med. Chem.

Medicinal Chemistry

PAINS ⓘ

1 alert: quinone_A ⓘ

Pan Assay Interference
Structures:
implemented from
Baell JB. & Holloway
GA. 2010 J. Med.
Chem.



Brenk ⓘ

Structural Alert:
implemented from 0 alert
Brenk R. et al. 2008
ChemMedChem

Leadlikeness ⓘ

Leadlikeness:
implemented from
Teague SJ. 1999 Angew.
Chem. Int. Ed. No; 1 violation: XLOGP3>3.5
 $250 \leq MW \leq 350$
 $XLOGP \leq 3.5$
Num. rotatable bonds $\leq$
7

Synthetic accessibility ⓘ

Synthetic accessibility
score: from 1 (very
easy) to 10 (very
difficult)
based on 1024
fragmental contributions 2.83
(FP2) modulated by size
and complexity penalties,
trained on 12'782'590
molecules and tested on
40 external molecules
($r^2 = 0.94$)

Topological Polar Surface Area: Calculated from Ertl P. et al. 2000 J. Med. Chem.	52.60 Å ²	<u>scale</u> Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Moderately soluble
Log $P_{o/w}$ (iLOGP) ⓘ	Lipophilicity	Log S (SILICOS-IT) ⓘ	
iLOGP: in-house physics-based method implemented from Daina A et al. 2014 J. Chem. Inf. Model.	3.05	SILICOS-IT: Fragmental method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos-it.com	-6.25
Log $P_{o/w}$ (XLOGP3) ⓘ		Solubility Class ⓘ	1.80e-04 mg/ml ; 5.62e-07 mol/l
XLOGP3: Atomistic and knowledge-based method calculated by XLOGP program, version 3.2.2, courtesy of CCBG, Shanghai Institute of Organic Chemistry	4.23	Solubility class: Log S <u>scale</u> Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Poorly soluble
Log $P_{o/w}$ (WLOGP) ⓘ			Pharmacokinetics
WLOGP: Atomistic method implemented from Wildman SA and Crippen GM. 1999 J. Chem. Inf. Model.	3.77	GI absorption ⓘ	
Log $P_{o/w}$ (MLOGP) ⓘ		Gastrointestinal absorption: according to the white of the BOILED-Egg	High
MLOGP: Topological method implemented from Moriguchi I. et al. 1992 Chem. Pharm. Bull. Moriguchi I. et al. 1994 Chem. Pharm. Bull. Lipinski PA. et al. 2001 Adv. Drug. Deliv. Rev.	1.97	BBB permeant ⓘ	
Log $P_{o/w}$ (SILICOS-IT) ⓘ		BBB permeation: according to the yolk of the BOILED-Egg	Yes
SILICOS-IT: Hybrid fragmental/topological method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos-it.com	4.64	P-gp substrate ⓘ	
Consensus Log $P_{o/w}$ ⓘ		P-glycoprotein substrate: SVM model built on 1033 molecules (training set) and tested on 415 molecules (test set) 10-fold CV: ACC=0.72 / AUC=0.77 External: ACC=0.88 / AUC=0.94	No
Consensus Log $P_{o/w}$:		CYP1A2 inhibitor ⓘ	
		Cytochrome P450 1A2 inhibitor: SVM model built on 9145 molecules (training set) and tested on 3000 molecules (test set)	Yes

Average of all five predictions

3.53

10-fold CV: ACC=0.83 / AUC=0.90
External: ACC=0.84 / AUC=0.91

CYP2C19 inhibitor ⓘ

Cytochrome P450 2C19 inhibitor: SVM model built on 9272 molecules (training set) and tested on 3000 molecules (test set) Yes
10-fold CV: ACC=0.80 / AUC=0.86
External: ACC=0.80 / AUC=0.87

CYP2C9 inhibitor ⓘ

Cytochrome P450 2C9 inhibitor: SVM model built on 5940 molecules (training set) and tested on 2075 molecules (test set) Yes
10-fold CV: ACC=0.78 / AUC=0.85
External: ACC=0.71 / AUC=0.81

CYP2D6 inhibitor ⓘ

Cytochrome P450 2D6 inhibitor: SVM model built on 3664 molecules (training set) and tested on 1068 molecules (test set) No
10-fold CV: ACC=0.79 / AUC=0.85
External: ACC=0.81 / AUC=0.87

CYP3A4 inhibitor ⓘ

Cytochrome P450 3A4 inhibitor: SVM model built on 7518 molecules (training set) and tested on 2579 molecules (test set) Yes
10-fold CV: ACC=0.77 / AUC=0.85
External: ACC=0.78 / AUC=0.86

Log K_p (skin permeation) ⓘ

Skin permeation: QSPR model implemented from -5.25 cm/s

Potts RO and Guy RH.
1992 Pharm. Res.

Druglikeness

Lipinski ⓘ

Lipinski (Pfizer) filter:
implemented from

Lipinski CA. et al. 2001

Adv. Drug Deliv. Rev.

Yes; 0 violation

MW $\leq$ 500

MLOGP $\leq$ 4.15

N or O $\leq$ 10

NH or OH $\leq$ 5

Ghose ⓘ

Ghose filter:

implemented from

Ghose AK. et al. 1999 J.

Comb. Chem.

Yes

160 $\leq$ MW $\leq$ 480

-0.4 $\leq$ WLOGP $\leq$ 5.6

40 $\leq$ MR $\leq$ 130

20 $\leq$ atoms $\leq$ 70

Veber ⓘ

Veber (GSK) filter:

implemented from

Veber DF. et al. 2002 J.

Yes

Med. Chem.

Rotatable bonds $\leq$ 10

TPSA $\leq$ 140

Egan ⓘ

Egan (Pharmacia)

filter: implemented

from

Egan WJ. et al. 2000 J.

Yes

Med. Chem.

WLOGP $\leq$ 5.88

TPSA $\leq$ 131.6

Muegge ⓘ

Muegge (Bayer) filter:

implemented from

Muegge I. et al. 2001 J.

Med. Chem.

200 $\leq$ MW $\leq$ 600

-2 $\leq$ XLOGP $\leq$ 5

TPSA $\leq$ 150

Yes

Num. rings $\leq$ 7

Num. carbon $>$ 4

Num. heteroatoms $>$ 1

Num. rotatable bonds $\leq$

15

H-bond acc. $\leq$ 10

H-bond don. $\leq$ 5

Bioavailability Score ⓘ


Abbott Bioavailability Score: Probability of F 0.56
> 10% in rat
implemented from
Martin YC. 2005 J.
Med. Chem.

Medicinal Chemistry

PAINS ⓘ

1 alert: quinone_A ⓘ


Pan Assay Interference
Structures:
implemented from
Baell JB. & Holloway
GA. 2010 J. Med.
Chem.



Brenk ⓘ

1 alert: isolated_alkene ⓘ

Structural Alert:
implemented from
Brenk R. et al. 2008
ChemMedChem




Leadlikeness ⓘ

Leadlikeness:
implemented from
Teague SJ. 1999 Angew.
Chem. Int. Ed.
250 ≤ MW ≤ 350
XLOGP ≤ 3.5
Num. rotatable bonds ≤
7

No; 1 violation: XLOGP3>3.5

Synthetic accessibility ⓘ

Synthetic accessibility
score: from 1 (very
easy) to 10 (very
difficult)
based on 1024
fragmental contributions 3.21
(FP2) modulated by size
and complexity penalties,
trained on 12'782'590
molecules and tested on
40 external molecules
($r^2 = 0.94$)



Enter a list of SMILES here:

Cc1cc(cc(C)c1Cl)OC3=CC(=O)c2ccccc2C3=O

Fill with an example

Clear



Run!

Show BOILED-Egg

Retrieve data:

Powered by
ChemAxon

Molecule 1



SMILES
S O=C1C(=CC(=O)c2c1cccc2)Oc1cc(C)c(c(c1)C)Cl

Physicochemical Properties

Formula	C18H13ClO3
Molecular weight	312.75 g/mol
Num. heavy atoms	22
Num. arom. heavy atoms	12
Fraction Csp3	0.11
Num. rotatable bonds	2
Num. H-bond acceptors	3
Num. H-bond donors	0
Molar Refractivity	85.20
TPSA	

Water Solubility

Log S (ESOL)

ESOL: Topological method implemented from
Delaney JS. 2004 J. Chem. Inf. Model.

-4.73

Solubility
Class

5.76e-03 mg/ml ; 1.84e-05 mol/l

Solubility class: Log S scale

Insoluble < -10 < Poorly Moderately soluble
< -6 < Moderately < -4
< Soluble < -2 Very < 0
< Highly

Log S (Ali)

Ali: Topological method implemented from
Ali J. et al. 2012 J. Chem. Inf. Model.

-4.88

Solubility
Class

4.10e-03 mg/ml ; 1.31e-05 mol/l

Solubility class: Log S

Topological Polar

Surface Area: 43.37 Å²
 Calculated from
 Ertl P. et al. 2000 J.
 Med. Chem.

Lipophilicity

Log $P_{o/w}$ (iLOGP) ②

iLOGP: in-house
 physics-based method
 implemented from
 Daina A et al. 2014 J.
 Chem. Inf. Model. 2.96

Log $P_{o/w}$ (XLOGP3) ②

XLOGP3: Atomistic
 and knowledge-based
 method calculated by
 XLOGP program,
 version 3.2.2, courtesy
 of CCBG, Shanghai
 Institute of Organic
 Chemistry 4.26

Log $P_{o/w}$ (WLOGP) ②

WLOGP: Atomistic
 method implemented
 from
 Wildman SA and
 Crippen GM. 1999 J.
 Chem. Inf. Model. 4.30

Log $P_{o/w}$ (MLOGP) ②

MLOGP: Topological
 method implemented
 from
 Moriguchi I. et al. 1992
 Chem. Pharm. Bull. 2.66
 Moriguchi I. et al. 1994
 Chem. Pharm. Bull.
 Lipinski PA. et al. 2001
 Adv. Drug. Deliv. Rev.

Log $P_{o/w}$ (SILICOS-IT) ②

SILICOS-IT: Hybrid
 fragmental/topological
 method calculated by
 FILTER-IT program,
 version 1.0.2, courtesy
 of SILICOS-IT. 4.99
<http://www.silicos-it.com>

Consensus Log $P_{o/w}$ ②

Consensus Log $P_{o/w}$:

scale

Insoluble < -10 < Poorly
 < -6 < Moderately < -4
 < Soluble < -2 Very < 0
 < Highly

Moderately soluble

Log S (SILICOS-IT) ②

SILICOS-IT:
 Fragmental method
 calculated by
 FILTER-IT program,
 version 1.0.2, courtesy
 of SILICOS-IT,
<http://www.silicos-it.com> -6.67

Solubility 6.75e-05 mg/ml ; 2.16e-07 mol/l
 Class ②

Solubility class: Log S
 scale
 Insoluble < -10 < Poorly
 < -6 < Moderately < -4
 < Soluble < -2 Very < 0
 < Highly

Pharmacokinetics

GI absorption ②

Gastrointestinal
 absorption: according
 to the white of the
 BOILED-Egg High

BBB permeant ②

BBB permeation:
 according to the yolk of
 the BOILED-Egg Yes

P-gp substrate ②

P-glycoprotein
 substrate: SVM model
 built on 1033 molecules
 (training set)
 and tested on 415
 molecules (test set) No
 10-fold CV: ACC=0.72 /
 AUC=0.77
 External: ACC=0.88 /
 AUC=0.94

CYP1A2 inhibitor ②

Cytochrome P450 1A2
 inhibitor: SVM model
 built on 9145 molecules
 (training set)
 and tested on 3000
 molecules (test set) Yes

Average of all five predictions

3.83

10-fold CV: ACC=0.83 / AUC=0.90
External: ACC=0.84 / AUC=0.91

CYP2C19 inhibitor ⓘ

Cytochrome P450 2C19 inhibitor: SVM model built on 9272 molecules (training set) and tested on 3000 molecules (test set) Yes
10-fold CV: ACC=0.80 / AUC=0.86
External: ACC=0.80 / AUC=0.87

CYP2C9 inhibitor ⓘ

Cytochrome P450 2C9 inhibitor: SVM model built on 5940 molecules (training set) and tested on 2075 molecules (test set) Yes
10-fold CV: ACC=0.78 / AUC=0.85
External: ACC=0.71 / AUC=0.81

CYP2D6 inhibitor ⓘ

Cytochrome P450 2D6 inhibitor: SVM model built on 3664 molecules (training set) and tested on 1068 molecules (test set) No
10-fold CV: ACC=0.79 / AUC=0.85
External: ACC=0.81 / AUC=0.87

CYP3A4 inhibitor ⓘ

Cytochrome P450 3A4 inhibitor: SVM model built on 7518 molecules (training set) and tested on 2579 molecules (test set) Yes
10-fold CV: ACC=0.77 / AUC=0.85
External: ACC=0.78 / AUC=0.86

Log K_p (skin permeation) ⓘ

Skin permeation: QSPR model implemented from -5.18 cm/s

Potts RO and Guy RH.
1992 Pharm. Res.

Druglikeness

Lipinski ⓘ

Lipinski (Pfizer) filter:
implemented from

Lipinski CA. et al. 2001

Adv. Drug Deliv. Rev.

Yes; 0 violation

MW $\leq$ 500

MLOGP $\leq$ 4.15

N or O $\leq$ 10

NH or OH $\leq$ 5

Ghose ⓘ

Ghose filter:

implemented from

Ghose AK. et al. 1999 J.

Comb. Chem.

Yes

160 $\leq$ MW $\leq$ 480

-0.4 $\leq$ WLOGP $\leq$ 5.6

40 $\leq$ MR $\leq$ 130

20 $\leq$ atoms $\leq$ 70

Veber ⓘ

Veber (GSK) filter:

implemented from

Veber DF. et al. 2002 J.

Yes

Med. Chem.

Rotatable bonds $\leq$ 10

TPSA $\leq$ 140

Egan ⓘ

Egan (Pharmacia)

filter: implemented

from

Egan WJ. et al. 2000 J.

Yes

Med. Chem.

WLOGP $\leq$ 5.88

TPSA $\leq$ 131.6

Muegge ⓘ

Muegge (Bayer) filter:

implemented from

Muegge I. et al. 2001 J.

Med. Chem.

200 $\leq$ MW $\leq$ 600

-2 $\leq$ XLOGP $\leq$ 5

TPSA $\leq$ 150

Yes

Num. rings $\leq$ 7

Num. carbon $>$ 4

Num. heteroatoms $>$ 1

Num. rotatable bonds $\leq$

15

H-bond acc. $\leq$ 10

H-bond don. $\leq$ 5

Bioavailability Score ⓘ


Abbott Bioavailability
Score: Probability of F 0.56
> 10% in rat
implemented from
Martin YC. 2005 J.
Med. Chem.

Medicinal Chemistry

PAINS ⓘ

1 alert: quinone_A ⓘ

Pan Assay Interference
Structures:
implemented from
Baell JB. & Holloway
GA. 2010 J. Med.
Chem.



Brenk ⓘ

Structural Alert:
implemented from 0 alert
Brenk R. et al. 2008
ChemMedChem

Leadlikeness ⓘ

Leadlikeness:
implemented from
Teague SJ. 1999 Angew.
Chem. Int. Ed. No; 1 violation: XLOGP3>3.5
 $250 \leq MW \leq 350$
 $XLOGP \leq 3.5$
Num. rotatable bonds $\leq$
7

Synthetic accessibility ⓘ

Synthetic accessibility
score: from 1 (very
easy) to 10 (very
difficult)
based on 1024
fragmental contributions 2.99
(FP2) modulated by size
and complexity penalties,
trained on 12'782'590
molecules and tested on
40 external molecules
($r^2 = 0.94$)

Topological Polar Surface Area: <u>Calculated from Ertl P. et al. 2000 J. Med. Chem.</u>	43.37 Å ²	<u>scale</u> Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Moderately soluble
Log <i>P</i>_{o/w} (iLOGP) ⓘ		Log <i>S</i> (SILICOS-IT) ⓘ	
iLOGP: in-house physics-based method implemented from Daina A et al. 2014 J. Chem. Inf. Model.	2.70	SILICOS-IT: <u>Fragmental method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos-it.com</u>	-6.50
Log <i>P</i>_{o/w} (XLOGP3) ⓘ		Solubility	1.00e-04 mg/ml ; 3.13e-07 mol/l
XLOGP3: Atomistic and knowledge-based method calculated by XLOGP program, version 3.2.2, courtesy of CCBG, Shanghai Institute of Organic Chemistry	4.16	Class ⓘ	
Log <i>P</i>_{o/w} (WLOGP) ⓘ		Solubility class: Log <i>S</i> scale Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Poorly soluble
WLOGP: Atomistic method implemented from Wildman SA and Crippen GM. 1999 J. Chem. Inf. Model.	4.34		Pharmacokinetics
Log <i>P</i>_{o/w} (MLOGP) ⓘ		GI absorption ⓘ	
MLOGP: Topological method implemented from Moriguchi I. et al. 1992 Chem. Pharm. Bull. Moriguchi I. et al. 1994 Chem. Pharm. Bull. Lipinski PA. et al. 2001 Adv. Drug. Deliv. Rev.	2.70	Gastrointestinal absorption: according to the white of the BOILED-Egg	High
Log <i>P</i>_{o/w} (SILICOS-IT) ⓘ		BBB permeant ⓘ	
SILICOS-IT: Hybrid fragmental/topological method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos-it.com	4.60	BBB permeation: according to the yolk of the BOILED-Egg	Yes
Consensus Log <i>P</i>_{o/w} ⓘ		P-gp substrate ⓘ	
Consensus Log <i>P</i>_{o/w}:		P-glycoprotein substrate: SVM model built on 1033 molecules (training set) and tested on 415 molecules (test set) 10-fold CV: ACC=0.72 / AUC=0.77 External: ACC=0.88 / AUC=0.94	No
		CYP1A2 inhibitor ⓘ	
		Cytochrome P450 1A2 inhibitor: SVM model built on 9145 molecules (training set) and tested on 3000 molecules (test set)	Yes

Average of all five predictions

3.70

10-fold CV: ACC=0.83 / AUC=0.90
External: ACC=0.84 / AUC=0.91

CYP2C19 inhibitor ⓘ

Cytochrome P450 2C19 inhibitor: SVM model built on 9272 molecules (training set) and tested on 3000 molecules (test set) Yes
10-fold CV: ACC=0.80 / AUC=0.86
External: ACC=0.80 / AUC=0.87

CYP2C9 inhibitor ⓘ

Cytochrome P450 2C9 inhibitor: SVM model built on 5940 molecules (training set) and tested on 2075 molecules (test set) Yes
10-fold CV: ACC=0.78 / AUC=0.85
External: ACC=0.71 / AUC=0.81

CYP2D6 inhibitor ⓘ

Cytochrome P450 2D6 inhibitor: SVM model built on 3664 molecules (training set) and tested on 1068 molecules (test set) No
10-fold CV: ACC=0.79 / AUC=0.85
External: ACC=0.81 / AUC=0.87

CYP3A4 inhibitor ⓘ

Cytochrome P450 3A4 inhibitor: SVM model built on 7518 molecules (training set) and tested on 2579 molecules (test set) Yes
10-fold CV: ACC=0.77 / AUC=0.85
External: ACC=0.78 / AUC=0.86

Log K_p (skin permeation) ⓘ

Skin permeation: QSPR model implemented from -5.29 cm/s

Potts RO and Guy RH.
1992 Pharm. Res.

Druglikeness

Lipinski ⓘ

Lipinski (Pfizer) filter:
implemented from
Lipinski CA. et al. 2001
Adv. Drug Deliv. Rev. Yes; 0 violation
MW $\leq$ 500
MLOGP $\leq$ 4.15
N or O $\leq$ 10
NH or OH $\leq$ 5

Ghose ⓘ

Ghose filter:
implemented from
Ghose AK. et al. 1999 J.
Comb. Chem. Yes
160 $\leq$ MW $\leq$ 480
-0.4 $\leq$ WLOGP $\leq$ 5.6
40 $\leq$ MR $\leq$ 130
20 $\leq$ atoms $\leq$ 70

Veber ⓘ

Veber (GSK) filter:
implemented from
Veber DF. et al. 2002 J. Yes
Med. Chem.
Rotatable bonds $\leq$ 10
TPSA $\leq$ 140

Egan ⓘ

Egan (Pharmacia)
filter: implemented
from
Egan WJ. et al. 2000 J. Yes
Med. Chem.
WLOGP $\leq$ 5.88
TPSA $\leq$ 131.6

Muegge ⓘ

Muegge (Bayer) filter:
implemented from
Muegge I. et al. 2001 J.
Med. Chem.
200 $\leq$ MW $\leq$ 600
-2 $\leq$ XLOGP $\leq$ 5
TPSA $\leq$ 150 Yes
Num. rings $\leq$ 7
Num. carbon $>$ 4
Num. heteroatoms $>$ 1
Num. rotatable bonds $\leq$
15
H-bond acc. $\leq$ 10
H-bond don. $\leq$ 5

Bioavailability Score ⓘ


Abbott Bioavailability
Score: Probability of F 0.56
> 10% in rat
implemented from
Martin YC. 2005 J.
Med. Chem.

Medicinal Chemistry

PAINS ⓘ

1 alert: quinone_A ⓘ

Pan Assay Interference
Structures:
implemented from
Baell JB. & Holloway
GA. 2010 J. Med.
Chem.



Brenk ⓘ


Structural Alert:
implemented from 0 alert
Brenk R. et al. 2008
ChemMedChem

Leadlikeness ⓘ

Leadlikeness:
implemented from
Teague SJ. 1999 Angew.
Chem. Int. Ed. No; 1 violation: XLOGP3>3.5
 $250 \leq MW \leq 350$
 $XLOGP \leq 3.5$
Num. rotatable bonds $\leq$
7

Synthetic accessibility ⓘ

Synthetic accessibility
score: from 1 (very
easy) to 10 (very
difficult)
based on 1024
fragmental contributions 2.85
(FP2) modulated by size
and complexity penalties,
trained on 12'782'590
molecules and tested on
40 external molecules
($r^2 = 0.94$)


[Show BOILED-Egg](#)



Enter a list of SMILES here:

Clc3cc(Cl)cc(Cl)c3OC2=CC(=O)c1ccccc1C2=O
[Fill with an example](#)
[Clear](#)
[Run!](#)

Retrieve data:


 Powered by
ChemAxon

Molecule 1


 SMILES
S Clc1cc(Cl)c(c(c1)Cl)OC1=CC(=O)c2c(C1=O)cccc2

Physicochemical Properties

Formula	C16H7Cl3O3
Molecular weight	353.58 g/mol
Num. heavy atoms	22
Num. arom. heavy atoms	12
Fraction Csp3	0.00
Num. rotatable bonds	2
Num. H-bond acceptors	3
Num. H-bond donors	0
Molar Refractivity	85.29
TPSA	

Water Solubility

Log S (ESOL)

ESOL: Topological method implemented from Delaney JS. 2004 J. Chem. Inf. Model.

-5.32

 Solubility
Class

1.69e-03 mg/ml ; 4.77e-06 mol/l

Solubility class: Log S scale

Insoluble < -10 < Poorly Moderately soluble
< -6 < Moderately < -4
< Soluble < -2 Very < 0
< Highly

Log S (Ali)

Ali: Topological method implemented from Ali J. et al. 2012 J. Chem. Inf. Model.

-5.43

 Solubility
Class

1.31e-03 mg/ml ; 3.69e-06 mol/l

Solubility class: Log S

Topological Polar Surface Area: Calculated from Ertl P. et al. 2000 J. Med. Chem.	43.37 Å ²	<u>scale</u> Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Moderately soluble
Log <i>P</i>_{o/w} (iLOGP) ⓘ	Lipophilicity	Log <i>S</i> (SILICOS-IT) ⓘ	
iLOGP: in-house physics-based method implemented from Daina A et al. 2014 J. Chem. Inf. Model.	2.77	SILICOS-IT: Fragmental method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos-it.com	-7.10
Log <i>P</i>_{o/w} (XLOGP3) ⓘ		Solubility Class ⓘ	2.82e-05 mg/ml ; 7.96e-08 mol/l
XLOGP3: Atomistic and knowledge-based method calculated by XLOGP program, version 3.2.2, courtesy of CCBG, Shanghai Institute of Organic Chemistry	4.79	Solubility class: Log <i>S</i> scale Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Poorly soluble
Log <i>P</i>_{o/w} (WLOGP) ⓘ			Pharmacokinetics
WLOGP: Atomistic method implemented from Wildman SA and Crippen GM. 1999 J. Chem. Inf. Model.	4.99	GI absorption ⓘ	
Log <i>P</i>_{o/w} (MLOGP) ⓘ		Gastrointestinal absorption: according to the white of the BOILED-Egg	High
MLOGP: Topological method implemented from Moriguchi I. et al. 1992 Chem. Pharm. Bull. Moriguchi I. et al. 1994 Chem. Pharm. Bull. Lipinski PA. et al. 2001 Adv. Drug. Deliv. Rev.	3.20	BBB permeant ⓘ	
Log <i>P</i>_{o/w} (SILICOS-IT) ⓘ		BBB permeation: according to the yolk of the BOILED-Egg	Yes
SILICOS-IT: Hybrid fragmental/topological method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos-it.com	5.24	P-gp substrate ⓘ	
Consensus Log <i>P</i>_{o/w} ⓘ		P-glycoprotein substrate: SVM model built on 1033 molecules (training set) and tested on 415 molecules (test set) 10-fold CV: ACC=0.72 / AUC=0.77 External: ACC=0.88 / AUC=0.94	No
Consensus Log <i>P</i>_{o/w}:		CYP1A2 inhibitor ⓘ	
		Cytochrome P450 1A2 inhibitor: SVM model built on 9145 molecules (training set) and tested on 3000 molecules (test set)	Yes

Average of all five predictions

4.20

10-fold CV: ACC=0.83 / AUC=0.90
External: ACC=0.84 / AUC=0.91

CYP2C19 inhibitor ⓘ

Cytochrome P450 2C19 inhibitor: SVM model built on 9272 molecules (training set) and tested on 3000 molecules (test set) Yes
10-fold CV: ACC=0.80 / AUC=0.86
External: ACC=0.80 / AUC=0.87

CYP2C9 inhibitor ⓘ

Cytochrome P450 2C9 inhibitor: SVM model built on 5940 molecules (training set) and tested on 2075 molecules (test set) Yes
10-fold CV: ACC=0.78 / AUC=0.85
External: ACC=0.71 / AUC=0.81

CYP2D6 inhibitor ⓘ

Cytochrome P450 2D6 inhibitor: SVM model built on 3664 molecules (training set) and tested on 1068 molecules (test set) No
10-fold CV: ACC=0.79 / AUC=0.85
External: ACC=0.81 / AUC=0.87

CYP3A4 inhibitor ⓘ

Cytochrome P450 3A4 inhibitor: SVM model built on 7518 molecules (training set) and tested on 2579 molecules (test set) No
10-fold CV: ACC=0.77 / AUC=0.85
External: ACC=0.78 / AUC=0.86

Log K_p (skin permeation) ⓘ

Skin permeation: QSPR model implemented from -5.06 cm/s

Potts RO and Guy RH.
1992 Pharm. Res.

Druglikeness

Lipinski ⓘ

Lipinski (Pfizer) filter:
implemented from
Lipinski CA. et al. 2001
Adv. Drug Deliv. Rev. Yes; 0 violation
MW $\leq$ 500
MLOGP $\leq$ 4.15
N or O $\leq$ 10
NH or OH $\leq$ 5

Ghose ⓘ

Ghose filter:
implemented from
Ghose AK. et al. 1999 J.
Comb. Chem. Yes
160 $\leq$ MW $\leq$ 480
-0.4 $\leq$ WLOGP $\leq$ 5.6
40 $\leq$ MR $\leq$ 130
20 $\leq$ atoms $\leq$ 70

Veber ⓘ

Veber (GSK) filter:
implemented from
Veber DF. et al. 2002 J. Yes
Med. Chem.
Rotatable bonds $\leq$ 10
TPSA $\leq$ 140

Egan ⓘ

Egan (Pharmacia)
filter: implemented
from
Egan WJ. et al. 2000 J. Yes
Med. Chem.
WLOGP $\leq$ 5.88
TPSA $\leq$ 131.6

Muegge ⓘ

Muegge (Bayer) filter:
implemented from
Muegge I. et al. 2001 J.
Med. Chem.
200 $\leq$ MW $\leq$ 600
-2 $\leq$ XLOGP $\leq$ 5
TPSA $\leq$ 150 Yes
Num. rings $\leq$ 7
Num. carbon $>$ 4
Num. heteroatoms $>$ 1
Num. rotatable bonds $\leq$
15
H-bond acc. $\leq$ 10
H-bond don. $\leq$ 5

Bioavailability Score ⓘ


Abbott Bioavailability
Score: Probability of F 0.56
> 10% in rat
implemented from
Martin YC. 2005 J.
Med. Chem.

Medicinal Chemistry

PAINS ⓘ

1 alert: quinone_A ⓘ

Pan Assay Interference
Structures:
implemented from
Baell JB. & Holloway
GA. 2010 J. Med.
Chem.



Brenk ⓘ

Structural Alert:
implemented from 0 alert
Brenk R. et al. 2008
ChemMedChem

Leadlikeness ⓘ

Leadlikeness:
implemented from
Teague SJ. 1999 Angew. No; 2 violations: MW>350,
Chem. Int. Ed. XLOGP3>3.5
 $250 \leq MW \leq 350$
 $XLOGP \leq 3.5$
Num. rotatable bonds $\leq$
7

Synthetic accessibility ⓘ

Synthetic accessibility
score: from 1 (very
easy) to 10 (very
difficult)
based on 1024
fragmental contributions 2.87
(FP2) modulated by size
and complexity penalties,
trained on 12'782'590
molecules and tested on
40 external molecules
($r^2 = 0.94$)

Topological Polar Surface Area:	43.37 Å ²	<u>scale</u> Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Moderately soluble
Calculated from Ertl P. et al. 2000 J. Med. Chem.			
	Lipophilicity		
Log <i>P</i> _{o/w} (iLOGP) ⓘ		Log <i>S</i> (SILICOS-IT) ⓘ	
<u>iLOGP: in-house physics-based method implemented from</u> Daina A et al. 2014 J. Chem. Inf. Model.	3.11	<u>SILICOS-IT: Fragmental method calculated by</u> FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos-it.com	-7.68
Log <i>P</i> _{o/w} (XLOGP3) ⓘ		Solubility	1.01e-05 mg/ml ; 2.07e-08 mol/l
<u>XLOGP3: Atomistic and knowledge-based method calculated by</u> XLOGP program, version 3.2.2, courtesy of CCBG, Shanghai Institute of Organic Chemistry	4.98	Class ⓘ	
Log <i>P</i> _{o/w} (WLOGP) ⓘ		<u>Solubility class: Log <i>S</i> scale</u> Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Poorly soluble
<u>WLOGP: Atomistic method implemented from</u> Wildman SA and Crippen GM. 1999 J. Chem. Inf. Model.	5.32		Pharmacokinetics
Log <i>P</i> _{o/w} (MLOGP) ⓘ		GI absorption ⓘ	
<u>MLOGP: Topological method implemented from</u> Moriguchi I. et al. 1992 Chem. Pharm. Bull. Moriguchi I. et al. 1994 Chem. Pharm. Bull. Lipinski PA. et al. 2001 Adv. Drug. Deliv. Rev.	3.54	<u>Gastrointestinal absorption: according to the white of the BOILED-Egg</u>	High
Log <i>P</i> _{o/w} (SILICOS-IT) ⓘ		BBB permeant ⓘ	
<u>SILICOS-IT: Hybrid fragmental/topological method calculated by</u> FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos-it.com	5.35	<u>BBB permeation: according to the yolk of the BOILED-Egg</u>	Yes
Consensus Log <i>P</i> _{o/w} ⓘ		P-gp substrate ⓘ	
<u>Consensus Log <i>P</i>_{o/w}:</u>		<u>P-glycoprotein substrate: SVM model built on 1033 molecules (training set)</u> and tested on 415 molecules (test set) 10-fold CV: ACC=0.72 / AUC=0.77 External: ACC=0.88 / AUC=0.94	No
		CYP1A2 inhibitor ⓘ	
		<u>Cytochrome P450 1A2 inhibitor: SVM model built on 9145 molecules (training set)</u> and tested on 3000 molecules (test set)	Yes

Average of all five
predictions

4.46

10-fold CV: ACC=0.83 /
AUC=0.90
External: ACC=0.84 /
AUC=0.91

CYP2C19 inhibitor ⓘ

Cytochrome P450
2C19 inhibitor: SVM
model built on 9272
molecules (training set)
and tested on 3000
molecules (test set) Yes
10-fold CV: ACC=0.80 /
AUC=0.86
External: ACC=0.80 /
AUC=0.87

CYP2C9 inhibitor ⓘ

Cytochrome P450 2C9
inhibitor: SVM model
built on 5940 molecules
(training set)
and tested on 2075
molecules (test set) Yes
10-fold CV: ACC=0.78 /
AUC=0.85
External: ACC=0.71 /
AUC=0.81

CYP2D6 inhibitor ⓘ

Cytochrome P450 2D6
inhibitor: SVM model
built on 3664 molecules
(training set)
and tested on 1068
molecules (test set) No
10-fold CV: ACC=0.79 /
AUC=0.85
External: ACC=0.81 /
AUC=0.87

CYP3A4 inhibitor ⓘ

Cytochrome P450 3A4
inhibitor: SVM model
built on 7518 molecules
(training set)
and tested on 2579
molecules (test set) No
10-fold CV: ACC=0.77 /
AUC=0.85
External: ACC=0.78 /
AUC=0.86

Log K_p (skin
permeation) ⓘ

Skin permeation:
QSPR model -5.73 cm/s
implemented from

Potts RO and Guy RH.
1992 Pharm. Res.

Druglikeness

Lipinski

Lipinski (Pfizer) filter:
implemented from
Lipinski CA. et al. 2001
Adv. Drug Deliv. Rev. Yes; 0 violation
MW $\leq$ 500
MLOGP $\leq$ 4.15
N or O $\leq$ 10
NH or OH $\leq$ 5

Ghose

Ghose filter:
implemented from
Ghose AK. et al. 1999 J.
Comb. Chem. No; 1 violation: MW>480
160 $\leq$ MW $\leq$ 480
-0.4 $\leq$ WLOGP $\leq$ 5.6
40 $\leq$ MR $\leq$ 130
20 $\leq$ atoms $\leq$ 70

Veber

Veber (GSK) filter:
implemented from
Veber DF. et al. 2002 J. Yes
Med. Chem.
Rotatable bonds $\leq$ 10
TPSA $\leq$ 140

Egan

Egan (Pharmacia)
filter: implemented
from
Egan WJ. et al. 2000 J. Yes
Med. Chem.
WLOGP $\leq$ 5.88
TPSA $\leq$ 131.6

Muegge

Muegge (Bayer) filter:
implemented from
Muegge I. et al. 2001 J.
Med. Chem.
200 $\leq$ MW $\leq$ 600
-2 $\leq$ XLOGP $\leq$ 5
TPSA $\leq$ 150 Yes
Num. rings $\leq$ 7
Num. carbon $>$ 4
Num. heteroatoms $>$ 1
Num. rotatable bonds $\leq$
15
H-bond acc. $\leq$ 10
H-bond don. $\leq$ 5

Bioavailability Score ⓘ


Abbott Bioavailability
Score: Probability of F 0.56
> 10% in rat
implemented from
Martin YC. 2005 J.
Med. Chem.

Medicinal Chemistry

PAINS ⓘ

1 alert: quinone_A ⓘ

Pan Assay Interference
Structures:
implemented from
Baell JB. & Holloway
GA. 2010 J. Med.
Chem.



Brenk ⓘ


Structural Alert:
implemented from 0 alert
Brenk R. et al. 2008
ChemMedChem

Leadlikeness ⓘ

Leadlikeness:
implemented from
Teague SJ. 1999 Angew. No; 2 violations: MW>350,
Chem. Int. Ed. XLOGP3>3.5
 $250 \leq MW \leq 350$
 $XLOGP \leq 3.5$
Num. rotatable bonds $\leq$
7

Synthetic accessibility ⓘ

Synthetic accessibility
score: from 1 (very
easy) to 10 (very
difficult)
based on 1024
fragmental contributions 2.85
(FP2) modulated by size
and complexity penalties,
trained on 12'782'590
molecules and tested on
40 external molecules
($r^2 = 0.94$)



Enter a list of SMILES here:

Clc3cc(OC2=CC(=O)c1ccccc1C2=O)c(Cl)cc3Cl

Fill with an example

Clear



Run!

Show BOILED-Egg

Retrieve data:


 Powered by
ChemAxon

Molecule 1


 SMILES
S Clc1cc(Cl)c(cc1OC1=CC(=O)c2c(C1=O)cccc2)Cl

Physicochemical Properties

Formula	C16H7Cl3O3
Molecular weight	353.58 g/mol
Num. heavy atoms	22
Num. arom. heavy atoms	12
Fraction Csp3	0.00
Num. rotatable bonds	2
Num. H-bond acceptors	3
Num. H-bond donors	0
Molar Refractivity	85.29
TPSA	

Water Solubility

Log S (ESOL)

ESOL: Topological method implemented from
Delaney JS. 2004 J. Chem. Inf. Model.

-5.32

 Solubility
Class

1.69e-03 mg/ml ; 4.77e-06 mol/l

Solubility class: Log S scale

Insoluble < -10 < Poorly Moderately soluble
< -6 < Moderately < -4
< Soluble < -2 Very < 0
< Highly

Log S (Ali)

Ali: Topological method implemented from
Ali J. et al. 2012 J. Chem. Inf. Model.

-5.43

 Solubility
Class

1.31e-03 mg/ml ; 3.69e-06 mol/l

Solubility class: Log S

Topological Polar Surface Area: Calculated from Ertl P. et al. 2000 J. Med. Chem.	43.37 Å ²	<u>scale</u> Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Moderately soluble
Log <i>P</i>_{o/w} (iLOGP) ⓘ	Lipophilicity	Log <i>S</i> (SILICOS-IT) ⓘ	
iLOGP: in-house physics-based method implemented from Daina A et al. 2014 J. Chem. Inf. Model.	2.93	SILICOS-IT: Fragmental method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos-it.com	-7.10
Log <i>P</i>_{o/w} (XLOGP3) ⓘ		Solubility Class ⓘ	2.82e-05 mg/ml ; 7.96e-08 mol/l
XLOGP3: Atomistic and knowledge-based method calculated by XLOGP program, version 3.2.2, courtesy of CCBG, Shanghai Institute of Organic Chemistry	4.79	Solubility class: Log <i>S</i> <u>scale</u> Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Poorly soluble
Log <i>P</i>_{o/w} (WLOGP) ⓘ			Pharmacokinetics
WLOGP: Atomistic method implemented from Wildman SA and Crippen GM. 1999 J. Chem. Inf. Model.	4.99	GI absorption ⓘ	
Log <i>P</i>_{o/w} (MLOGP) ⓘ		Gastrointestinal absorption: according to the white of the BOILED-Egg	High
MLOGP: Topological method implemented from Moriguchi I. et al. 1992 Chem. Pharm. Bull. Moriguchi I. et al. 1994 Chem. Pharm. Bull. Lipinski PA. et al. 2001 Adv. Drug. Deliv. Rev.	3.20	BBB permeant ⓘ	
Log <i>P</i>_{o/w} (SILICOS-IT) ⓘ		BBB permeation: according to the yolk of the BOILED-Egg	Yes
SILICOS-IT: Hybrid fragmental/topological method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos-it.com	5.24	P-gp substrate ⓘ	
Consensus Log <i>P</i>_{o/w} ⓘ		P-glycoprotein substrate: SVM model built on 1033 molecules (training set) and tested on 415 molecules (test set) 10-fold CV: ACC=0.72 / AUC=0.77 External: ACC=0.88 / AUC=0.94	No
Consensus Log <i>P</i>_{o/w}:		CYP1A2 inhibitor ⓘ	
		Cytochrome P450 1A2 inhibitor: SVM model built on 9145 molecules (training set) and tested on 3000 molecules (test set)	Yes

Average of all five predictions

4.23

10-fold CV: ACC=0.83 / AUC=0.90
External: ACC=0.84 / AUC=0.91

CYP2C19 inhibitor ⓘ

Cytochrome P450 2C19 inhibitor: SVM model built on 9272 molecules (training set) and tested on 3000 molecules (test set) Yes
10-fold CV: ACC=0.80 / AUC=0.86
External: ACC=0.80 / AUC=0.87

CYP2C9 inhibitor ⓘ

Cytochrome P450 2C9 inhibitor: SVM model built on 5940 molecules (training set) and tested on 2075 molecules (test set) Yes
10-fold CV: ACC=0.78 / AUC=0.85
External: ACC=0.71 / AUC=0.81

CYP2D6 inhibitor ⓘ

Cytochrome P450 2D6 inhibitor: SVM model built on 3664 molecules (training set) and tested on 1068 molecules (test set) No
10-fold CV: ACC=0.79 / AUC=0.85
External: ACC=0.81 / AUC=0.87

CYP3A4 inhibitor ⓘ

Cytochrome P450 3A4 inhibitor: SVM model built on 7518 molecules (training set) and tested on 2579 molecules (test set) No
10-fold CV: ACC=0.77 / AUC=0.85
External: ACC=0.78 / AUC=0.86

Log K_p (skin permeation) ⓘ

Skin permeation: QSPR model implemented from -5.06 cm/s

Potts RO and Guy RH.
1992 Pharm. Res.

Druglikeness

Lipinski ⓘ

Lipinski (Pfizer) filter:
implemented from
Lipinski CA. et al. 2001
Adv. Drug Deliv. Rev. Yes; 0 violation
MW $\leq$ 500
MLOGP $\leq$ 4.15
N or O $\leq$ 10
NH or OH $\leq$ 5

Ghose ⓘ

Ghose filter:
implemented from
Ghose AK. et al. 1999 J.
Comb. Chem. Yes
160 $\leq$ MW $\leq$ 480
-0.4 $\leq$ WLOGP $\leq$ 5.6
40 $\leq$ MR $\leq$ 130
20 $\leq$ atoms $\leq$ 70

Veber ⓘ

Veber (GSK) filter:
implemented from
Veber DF. et al. 2002 J. Yes
Med. Chem.
Rotatable bonds $\leq$ 10
TPSA $\leq$ 140

Egan ⓘ

Egan (Pharmacia)
filter: implemented
from
Egan WJ. et al. 2000 J. Yes
Med. Chem.
WLOGP $\leq$ 5.88
TPSA $\leq$ 131.6

Muegge ⓘ

Muegge (Bayer) filter:
implemented from
Muegge I. et al. 2001 J.
Med. Chem.
200 $\leq$ MW $\leq$ 600
-2 $\leq$ XLOGP $\leq$ 5
TPSA $\leq$ 150 Yes
Num. rings $\leq$ 7
Num. carbon $>$ 4
Num. heteroatoms $>$ 1
Num. rotatable bonds $\leq$
15
H-bond acc. $\leq$ 10
H-bond don. $\leq$ 5

Bioavailability Score ⓘ


Abbott Bioavailability
Score: Probability of F 0.56
> 10% in rat
implemented from
Martin YC. 2005 J.
Med. Chem.

Medicinal Chemistry

PAINS ⓘ

1 alert: quinone_A ⓘ

Pan Assay Interference
Structures:
implemented from
Baell JB. & Holloway
GA. 2010 J. Med.
Chem.




quinone_A

Brenk ⓘ

1 alert: halogenated_ring_1 ⓘ

Structural Alert:
implemented from
Brenk R. et al. 2008
ChemMedChem



halogenated_ring_1

Leadlikeness ⓘ

Leadlikeness:
implemented from
Teague SJ. 1999 Angew.
Chem. Int. Ed.
250 ≤ MW ≤ 350
XLOGP ≤ 3.5
Num. rotatable bonds ≤
7

No; 2 violations: MW>350,
 XLOGP3>3.5

Synthetic accessibility ⓘ

Synthetic accessibility
score: from 1 (very
easy) to 10 (very
difficult)
based on 1024
fragmental contributions 2.93
(FP2) modulated by size
and complexity penalties,
trained on 12'782'590
molecules and tested on
40 external molecules
($r^2 = 0.94$)

Topological Polar Surface Area: Calculated from Ertl P. et al. 2000 J. Med. Chem.	43.37 Å ²	<u>scale</u> Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Moderately soluble
Log $P_{o/w}$ (iLOGP) ⓘ		Log S (SILICOS-IT) ⓘ	
iLOGP: in-house physics-based method implemented from Daina A et al. 2014 J. Chem. Inf. Model.	2.66	SILICOS-IT: Fragmental method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos-it.com	-6.50
Log $P_{o/w}$ (XLOGP3) ⓘ		Solubility	1.00e-04 mg/ml ; 3.13e-07 mol/l
XLOGP3: Atomistic and knowledge-based method calculated by XLOGP program, version 3.2.2, courtesy of CCBG, Shanghai Institute of Organic Chemistry	4.16	Class ⓘ	
Log $P_{o/w}$ (WLOGP) ⓘ		Solubility class: Log S scale Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Poorly soluble
WLOGP: Atomistic method implemented from Wildman SA and Crippen GM. 1999 J. Chem. Inf. Model.	4.34		Pharmacokinetics
Log $P_{o/w}$ (MLOGP) ⓘ		GI absorption ⓘ	
MLOGP: Topological method implemented from Moriguchi I. et al. 1992 Chem. Pharm. Bull. Moriguchi I. et al. 1994 Chem. Pharm. Bull. Lipinski PA. et al. 2001 Adv. Drug. Deliv. Rev.	2.70	Gastrointestinal absorption: according to the white of the BOILED-Egg	High
Log $P_{o/w}$ (SILICOS-IT) ⓘ		BBB permeant ⓘ	
SILICOS-IT: Hybrid fragmental/topological method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos-it.com	4.60	BBB permeation: according to the yolk of the BOILED-Egg	Yes
Consensus Log $P_{o/w}$ ⓘ		P-gp substrate ⓘ	
Consensus Log $P_{o/w}$:		P-glycoprotein substrate: SVM model built on 1033 molecules (training set) and tested on 415 molecules (test set) 10-fold CV: ACC=0.72 / AUC=0.77 External: ACC=0.88 / AUC=0.94	No
		CYP1A2 inhibitor ⓘ	
		Cytochrome P450 1A2 inhibitor: SVM model built on 9145 molecules (training set) and tested on 3000 molecules (test set)	Yes

Average of all five predictions

3.69

10-fold CV: ACC=0.83 / AUC=0.90
External: ACC=0.84 / AUC=0.91

CYP2C19 inhibitor ⓘ

Cytochrome P450 2C19 inhibitor: SVM model built on 9272 molecules (training set) and tested on 3000 molecules (test set) Yes
10-fold CV: ACC=0.80 / AUC=0.86
External: ACC=0.80 / AUC=0.87

CYP2C9 inhibitor ⓘ

Cytochrome P450 2C9 inhibitor: SVM model built on 5940 molecules (training set) and tested on 2075 molecules (test set) Yes
10-fold CV: ACC=0.78 / AUC=0.85
External: ACC=0.71 / AUC=0.81

CYP2D6 inhibitor ⓘ

Cytochrome P450 2D6 inhibitor: SVM model built on 3664 molecules (training set) and tested on 1068 molecules (test set) No
10-fold CV: ACC=0.79 / AUC=0.85
External: ACC=0.81 / AUC=0.87

CYP3A4 inhibitor ⓘ

Cytochrome P450 3A4 inhibitor: SVM model built on 7518 molecules (training set) and tested on 2579 molecules (test set) Yes
10-fold CV: ACC=0.77 / AUC=0.85
External: ACC=0.78 / AUC=0.86

Log K_p (skin permeation) ⓘ

Skin permeation: QSPR model implemented from -5.29 cm/s

Potts RO and Guy RH.
1992 Pharm. Res.

Druglikeness

Lipinski

Lipinski (Pfizer) filter:
implemented from

Lipinski CA. et al. 2001

Adv. Drug Deliv. Rev.

Yes; 0 violation

MW $\leq$ 500

MLOGP $\leq$ 4.15

N or O $\leq$ 10

NH or OH $\leq$ 5

Ghose

Ghose filter:

implemented from

Ghose AK. et al. 1999 J.

Comb. Chem.

Yes

160 $\leq$ MW $\leq$ 480

-0.4 $\leq$ WLOGP $\leq$ 5.6

40 $\leq$ MR $\leq$ 130

20 $\leq$ atoms $\leq$ 70

Veber

Veber (GSK) filter:

implemented from

Veber DF. et al. 2002 J.

Yes

Med. Chem.

Rotatable bonds $\leq$ 10

TPSA $\leq$ 140

Egan

Egan (Pharmacia)

filter: implemented

from

Egan WJ. et al. 2000 J.

Yes

Med. Chem.

WLOGP $\leq$ 5.88

TPSA $\leq$ 131.6

Muegge

Muegge (Bayer) filter:

implemented from

Muegge I. et al. 2001 J.

Med. Chem.

200 $\leq$ MW $\leq$ 600

-2 $\leq$ XLOGP $\leq$ 5

TPSA $\leq$ 150

Yes

Num. rings $\leq$ 7

Num. carbon $>$ 4

Num. heteroatoms $>$ 1

Num. rotatable bonds $\leq$

15

H-bond acc. $\leq$ 10

H-bond don. $\leq$ 5

Bioavailability Score ⓘ


Abbott Bioavailability
Score: Probability of F 0.56
> 10% in rat
implemented from
Martin YC. 2005 J.
Med. Chem.

Medicinal Chemistry

PAINS ⓘ

1 alert: quinone_A ⓘ

Pan Assay Interference
Structures:
implemented from
Baell JB. & Holloway
GA. 2010 J. Med.
Chem.



Brenk ⓘ

Structural Alert:
implemented from 0 alert
Brenk R. et al. 2008
ChemMedChem

Leadlikeness ⓘ

Leadlikeness:
implemented from
Teague SJ. 1999 Angew.
Chem. Int. Ed. No; 1 violation: XLOGP3>3.5
 $250 \leq MW \leq 350$
 $XLOGP \leq 3.5$
Num. rotatable bonds $\leq$
7

Synthetic accessibility ⓘ

Synthetic accessibility
score: from 1 (very
easy) to 10 (very
difficult)
based on 1024
fragmental contributions 2.87
(FP2) modulated by size
and complexity penalties,
trained on 12'782'590
molecules and tested on
40 external molecules
($r^2 = 0.94$)

Bioavailability Score ⓘ


Abbott Bioavailability
Score: Probability of F 0.56
> 10% in rat
implemented from
Martin YC. 2005 J.
Med. Chem.

Medicinal Chemistry

PAINS ⓘ

1 alert: quinone_A ⓘ

Pan Assay Interference
Structures:
implemented from
Baell JB. & Holloway
GA. 2010 J. Med.
Chem.



Brenk ⓘ

Structural Alert:
implemented from 0 alert
Brenk R. et al. 2008
ChemMedChem

Leadlikeness ⓘ

Leadlikeness:
implemented from
Teague SJ. 1999 Angew. No; 2 violations: MW>350,
Chem. Int. Ed. XLOGP3>3.5
 $250 \leq MW \leq 350$
 $XLOGP \leq 3.5$
Num. rotatable bonds $\leq$
7

Synthetic accessibility ⓘ

Synthetic accessibility
score: from 1 (very
easy) to 10 (very
difficult)
based on 1024
fragmental contributions 2.85
(FP2) modulated by size
and complexity penalties,
trained on 12'782'590
molecules and tested on
40 external molecules
($r^2 = 0.94$)

In order to avoid security-related warning messages when switching to secured connection, you may want either to:

- confirm the exception on the next page, or
- import our [CA key](#) in your web browser

Click [here](#) to proceed.

[Click2Drug](#)

[SwissDock](#)

[SwissParam](#)

[SwissSidechain](#)

[SwissBioisostere](#)

[SwissTargetPrediction](#)

[SwissADME](#)

[SwissSimilarity](#)

[About us](#)

[Swiss Institute of Bioinformatics](#)

SwissADME

- [Disclaimer](#)
- [Help](#)
- [FAQ](#)
- [Home](#)


This website allows you to compute physicochemical descriptors as well as to predict ADME parameters, pharmacokinetic properties, druglike nature and medicinal chemistry friendliness of one or multiple small molecules to support drug discovery.

The main article describing the web service and its underlying methodologies is [SwissADME: a free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules. *Sci. Rep.* \(2017\) 7:42717.](#)

For details about development and validation of iLOG, please refer to this article: [iLOGP: a simple, robust, and efficient description of *n*-octanol/water partition coefficient for drug design using the GB/SA approach. *J. Chem. Inf. Model.* \(2014\) 54\(12\):3284-3301.](#)

For details about development and validation of the BOILED-Egg, please refer to this article: [A BOILED-Egg to predict gastrointestinal absorption and brain penetration of small molecules. *ChemMedChem* \(2016\) 11\(11\):1117-1121.](#)

Developed and maintained by the [Molecular Modeling Group](#) of the SIB | Swiss Institute of Bioinformatics.


[Show BOILED-Egg](#)



Enter a list of SMILES here:

[O-][N+](=O)c1ccc(cc1)OC3=CC(=O)c2ccccc2C3=O
[Fill with an example](#)
[Clear](#)
[Run!](#)

Retrieve data:


 Powered by
ChemAxon

Molecule 1


 SMILES
O=C1C(=CC(=O)c2c1cccc2)Oc1ccc(cc1)[N+](=O)[O-]

Physicochemical Properties

Formula	C16H9NO5
Molecular weight	295.25 g/mol
Num. heavy atoms	22
Num. arom. heavy atoms	12
Fraction Csp3	0.00
Num. rotatable bonds	3
Num. H-bond acceptors	5
Num. H-bond donors	0
Molar Refractivity	79.08
TPSA	

Water Solubility

Log *S* (ESOL)

ESOL: Topological method implemented from
Delaney JS. 2004 J. Chem. Inf. Model.

-3.60

 Solubility
 Class

7.38e-02 mg/ml ; 2.50e-04 mol/l

Solubility class: Log *S* scale

Insoluble < -10 < Poorly Soluble
< -6 < Moderately < -4
< Soluble < -2 Very < 0
< Highly

Log *S* (Ali)

Ali: Topological method implemented from
Ali J. et al. 2012 J. Chem. Inf. Model.

-4.27

 Solubility
 Class

1.59e-02 mg/ml ; 5.40e-05 mol/l

Solubility class: Log *S*

Topological Polar Surface Area: Calculated from Ertl P. et al. 2000 J. Med. Chem.	89.19 Å ²	<u>scale</u> Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Moderately soluble
Lipophilicity Log $P_{o/w}$ (iLOGP) ⓘ		Log S (SILICOS-IT) ⓘ	
iLOGP: in-house physics-based method implemented from Daina A et al. 2014 J. Chem. Inf. Model.	2.01	SILICOS-IT: Fragmental method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos-it.com	-4.66
Log $P_{o/w}$ (XLOGP3) ⓘ		Solubility Class ⓘ	6.40e-03 mg/ml ; 2.17e-05 mol/l
XLOGP3: Atomistic and knowledge-based method calculated by XLOGP program, version 3.2.2, courtesy of CCBG, Shanghai Institute of Organic Chemistry	2.74	Solubility class: Log S <u>scale</u> Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Moderately soluble
Log $P_{o/w}$ (WLOGP) ⓘ			Pharmacokinetics
WLOGP: Atomistic method implemented from Wildman SA and Crippen GM. 1999 J. Chem. Inf. Model.	2.94	GI absorption ⓘ	
Log $P_{o/w}$ (MLOGP) ⓘ		Gastrointestinal absorption: according to the white of the BOILED-Egg	High
MLOGP: Topological method implemented from Moriguchi I. et al. 1992 Chem. Pharm. Bull. Moriguchi I. et al. 1994 Chem. Pharm. Bull. Lipinski PA. et al. 2001 Adv. Drug. Deliv. Rev.	1.48	BBB permeant ⓘ	
Log $P_{o/w}$ (SILICOS-IT) ⓘ		BBB permeation: according to the yolk of the BOILED-Egg	No
SILICOS-IT: Hybrid fragmental/topological method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos-it.com	1.15	P-gp substrate ⓘ	
Consensus Log $P_{o/w}$ ⓘ		P-glycoprotein substrate: SVM model built on 1033 molecules (training set) and tested on 415 molecules (test set) 10-fold CV: ACC=0.72 / AUC=0.77 External: ACC=0.88 / AUC=0.94	No
Consensus Log $P_{o/w}$:		CYP1A2 inhibitor ⓘ	
		Cytochrome P450 1A2 inhibitor: SVM model built on 9145 molecules (training set) and tested on 3000 molecules (test set)	Yes

Average of all five predictions

2.06

10-fold CV: ACC=0.83 / AUC=0.90
External: ACC=0.84 / AUC=0.91

CYP2C19 inhibitor ⓘ

Cytochrome P450 2C19 inhibitor: SVM model built on 9272 molecules (training set) and tested on 3000 molecules (test set) Yes
10-fold CV: ACC=0.80 / AUC=0.86
External: ACC=0.80 / AUC=0.87

CYP2C9 inhibitor ⓘ

Cytochrome P450 2C9 inhibitor: SVM model built on 5940 molecules (training set) and tested on 2075 molecules (test set) Yes
10-fold CV: ACC=0.78 / AUC=0.85
External: ACC=0.71 / AUC=0.81

CYP2D6 inhibitor ⓘ

Cytochrome P450 2D6 inhibitor: SVM model built on 3664 molecules (training set) and tested on 1068 molecules (test set) No
10-fold CV: ACC=0.79 / AUC=0.85
External: ACC=0.81 / AUC=0.87

CYP3A4 inhibitor ⓘ

Cytochrome P450 3A4 inhibitor: SVM model built on 7518 molecules (training set) and tested on 2579 molecules (test set) No
10-fold CV: ACC=0.77 / AUC=0.85
External: ACC=0.78 / AUC=0.86

Log K_p (skin permeation) ⓘ

Skin permeation: QSPR model implemented from -6.16 cm/s

Potts RO and Guy RH.
1992 Pharm. Res.

Druglikeness

Lipinski ⓘ

Lipinski (Pfizer) filter:
implemented from

Lipinski CA. et al. 2001

Adv. Drug Deliv. Rev.

Yes; 0 violation

MW $\leq$ 500

MLOGP $\leq$ 4.15

N or O $\leq$ 10

NH or OH $\leq$ 5

Ghose ⓘ

Ghose filter:

implemented from

Ghose AK. et al. 1999 J.

Comb. Chem.

Yes

160 $\leq$ MW $\leq$ 480

-0.4 $\leq$ WLOGP $\leq$ 5.6

40 $\leq$ MR $\leq$ 130

20 $\leq$ atoms $\leq$ 70

Veber ⓘ

Veber (GSK) filter:

implemented from

Veber DF. et al. 2002 J.

Yes

Med. Chem.

Rotatable bonds $\leq$ 10

TPSA $\leq$ 140

Egan ⓘ

Egan (Pharmacia)

filter: implemented

from

Egan WJ. et al. 2000 J.

Yes

Med. Chem.

WLOGP $\leq$ 5.88

TPSA $\leq$ 131.6

Muegge ⓘ

Muegge (Bayer) filter:

implemented from

Muegge I. et al. 2001 J.

Med. Chem.

200 $\leq$ MW $\leq$ 600

-2 $\leq$ XLOGP $\leq$ 5

TPSA $\leq$ 150

Yes

Num. rings $\leq$ 7

Num. carbon $>$ 4

Num. heteroatoms $>$ 1

Num. rotatable bonds $\leq$

15

H-bond acc. $\leq$ 10

H-bond don. $\leq$ 5

Bioavailability Score ⓘ


Abbott Bioavailability Score: Probability of F 0.56
> 10% in rat
implemented from
Martin YC. 2005 J.
Med. Chem.

Medicinal Chemistry

PAINS ⓘ

1 alert: quinone_A ⓘ

Pan Assay Interference
Structures:
implemented from
Baell JB. & Holloway
GA. 2010 J. Med.
Chem.




quinone_A


2 alerts: nitro_group, oxygen-nitrogen_single_bond ⓘ

Brenk ⓘ

Structural Alert:
implemented from
Brenk R. et al. 2008
ChemMedChem



nitro_group



aliphatic, not in a ring

oxygen-nitrogen_single_bon

Leadlikeness ⓘ

Leadlikeness:
implemented from
Teague SJ. 1999 Angew.
Chem. Int. Ed. Yes
250 ≤ MW ≤ 350
XLOGP ≤ 3.5
Num. rotatable bonds ≤
7

Synthetic accessibility ⓘ

Synthetic accessibility
score: from 1 (very
easy) to 10 (very
difficult)
based on 1024
fragmental contributions 2.82
(FP2) modulated by size
and complexity penalties,
trained on 12'782'590
molecules and tested on
40 external molecules
(R² = 0.94)

In order to avoid security-related warning messages when switching to secured connection, you may want either to:

- confirm the exception on the next page, or
- import our [CA key](#) in your web browser

Click [here](#) to proceed.

[Click2Drug](#)

[SwissDock](#)

[SwissParam](#)

[SwissSidechain](#)

[SwissBioisostere](#)

[SwissTargetPrediction](#)

[SwissADME](#)

[SwissSimilarity](#)

[About us](#)

[Swiss Institute of Bioinformatics](#)

SwissADME

- [Disclaimer](#)
- [Help](#)
- [FAQ](#)
- [Home](#)


This website allows you to compute physicochemical descriptors as well as to predict ADME parameters, pharmacokinetic properties, druglike nature and medicinal chemistry friendliness of one or multiple small molecules to support drug discovery.

The main article describing the web service and its underlying methodologies is [SwissADME: a free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules](#). *Sci. Rep.* (2017) 7:42717.

For details about development and validation of iLOG, please refer to this article: [iLOGP: a simple, robust, and efficient description of *n*-octanol/water partition coefficient for drug design using the GB/SA approach](#). *J. Chem. Inf. Model.* (2014) 54(12):3284-3301.

For details about development and validation of the BOILED-Egg, please refer to this article: [A BOILED-Egg to predict gastrointestinal absorption and brain penetration of small molecules](#). *ChemMedChem* (2016) 11(11):1117-1121.

Developed and maintained by the [Molecular Modeling Group](#) of the SIB | Swiss Institute of Bioinformatics.



Enter a list of SMILES here:

COc1ccc(cc1)OC3=CC(=O)c2ccccc2C3=O

Fill with an example

Clear



Run!

Show BOILED-Egg

Retrieve data:


 Powered by
ChemAxon

Molecule 1


 SMILES
S COc1ccc(cc1)OC1=CC(=O)c2c(C1=O)cccc2

Physicochemical Properties

Formula	C17H12O4
Molecular weight	280.27 g/mol
Num. heavy atoms	21
Num. arom. heavy atoms	12
Fraction Csp3	0.06
Num. rotatable bonds	3
Num. H-bond acceptors	4
Num. H-bond donors	0
Molar Refractivity	76.75
TPSA	

Water Solubility

Log S (ESOL)

ESOL: Topological method implemented from
Delaney JS. 2004 J. Chem. Inf. Model.

-3.62

 Solubility
Class

6.77e-02 mg/ml ; 2.42e-04 mol/l

Solubility class: Log S scale

Insoluble < -10 < Poorly Soluble
< -6 < Moderately < -4
< Soluble < -2 Very < 0
< Highly

Log S (Ali)

Ali: Topological method implemented from
Ali J. et al. 2012 J. Chem. Inf. Model.

-3.64

 Solubility
Class

6.36e-02 mg/ml ; 2.27e-04 mol/l

Solubility class: Log S

Topological Polar Surface Area:	52.60 Å ²	<u>scale</u> Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Soluble
Calculated from Ertl P. et al. 2000 J. Med. Chem.			
	Lipophilicity	Log S (SILICOS-IT) ?	
Log <i>P</i> _{o/w} (iLOGP) ?		SILICOS-IT: Fragmental method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos-it.com	-5.42
<u>iLOGP: in-house physics-based method implemented from Daina A et al. 2014 J. Chem. Inf. Model.</u>	2.57		
Log <i>P</i> _{o/w} (XLOGP3) ?		Solubility Class ?	1.07e-03 mg/ml ; 3.83e-06 mol/l
<u>XLOGP3: Atomistic and knowledge-based method calculated by XLOGP program, version 3.2.2, courtesy of CCBG, Shanghai Institute of Organic Chemistry</u>	2.88	<u>Solubility class: Log S scale</u> Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Moderately soluble
Log <i>P</i> _{o/w} (WLOGP) ?			Pharmacokinetics
<u>WLOGP: Atomistic method implemented from Wildman SA and Crippen GM. 1999 J. Chem. Inf. Model.</u>	3.04	GI absorption ?	
Log <i>P</i> _{o/w} (MLOGP) ?		<u>Gastrointestinal absorption: according to the white of the BOILED-Egg</u>	High
<u>MLOGP: Topological method implemented from Moriguchi I. et al. 1992 Chem. Pharm. Bull. Moriguchi I. et al. 1994 Chem. Pharm. Bull. Lipinski PA. et al. 2001 Adv. Drug. Deliv. Rev.</u>	1.34	BBB permeant ?	
Log <i>P</i> _{o/w} (SILICOS-IT) ?		<u>BBB permeation: according to the yolk of the BOILED-Egg</u>	Yes
<u>SILICOS-IT: Hybrid fragmental/topological method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos-it.com</u>	3.38	P-gp substrate ?	
Consensus Log <i>P</i> _{o/w} ?		<u>P-glycoprotein substrate: SVM model built on 1033 molecules (training set) and tested on 415 molecules (test set) 10-fold CV: ACC=0.72 / AUC=0.77 External: ACC=0.88 / AUC=0.94</u>	No
<u>Consensus Log <i>P</i>_{o/w}:</u>		CYP1A2 inhibitor ?	
		<u>Cytochrome P450 1A2 inhibitor: SVM model built on 9145 molecules (training set) and tested on 3000 molecules (test set)</u>	Yes

Average of all five predictions

2.64

10-fold CV: ACC=0.83 / AUC=0.90
External: ACC=0.84 / AUC=0.91

CYP2C19 inhibitor ⓘ

Cytochrome P450 2C19 inhibitor: SVM model built on 9272 molecules (training set) and tested on 3000 molecules (test set) Yes
10-fold CV: ACC=0.80 / AUC=0.86
External: ACC=0.80 / AUC=0.87

CYP2C9 inhibitor ⓘ

Cytochrome P450 2C9 inhibitor: SVM model built on 5940 molecules (training set) and tested on 2075 molecules (test set) Yes
10-fold CV: ACC=0.78 / AUC=0.85
External: ACC=0.71 / AUC=0.81

CYP2D6 inhibitor ⓘ

Cytochrome P450 2D6 inhibitor: SVM model built on 3664 molecules (training set) and tested on 1068 molecules (test set) No
10-fold CV: ACC=0.79 / AUC=0.85
External: ACC=0.81 / AUC=0.87

CYP3A4 inhibitor ⓘ

Cytochrome P450 3A4 inhibitor: SVM model built on 7518 molecules (training set) and tested on 2579 molecules (test set) Yes
10-fold CV: ACC=0.77 / AUC=0.85
External: ACC=0.78 / AUC=0.86

Log K_p (skin permeation) ⓘ

Skin permeation: QSPR model implemented from -5.96 cm/s

Potts RO and Guy RH.
1992 Pharm. Res.

Druglikeness

Lipinski ⓘ

Lipinski (Pfizer) filter:
implemented from
Lipinski CA. et al. 2001
Adv. Drug Deliv. Rev. Yes; 0 violation
MW $\leq$ 500
MLOGP $\leq$ 4.15
N or O $\leq$ 10
NH or OH $\leq$ 5

Ghose ⓘ

Ghose filter:
implemented from
Ghose AK. et al. 1999 J.
Comb. Chem. Yes
160 $\leq$ MW $\leq$ 480
-0.4 $\leq$ WLOGP $\leq$ 5.6
40 $\leq$ MR $\leq$ 130
20 $\leq$ atoms $\leq$ 70

Veber ⓘ

Veber (GSK) filter:
implemented from
Veber DF. et al. 2002 J. Yes
Med. Chem.
Rotatable bonds $\leq$ 10
TPSA $\leq$ 140

Egan ⓘ

Egan (Pharmacia)
filter: implemented
from
Egan WJ. et al. 2000 J. Yes
Med. Chem.
WLOGP $\leq$ 5.88
TPSA $\leq$ 131.6

Muegge ⓘ

Muegge (Bayer) filter:
implemented from
Muegge I. et al. 2001 J.
Med. Chem.
200 $\leq$ MW $\leq$ 600
-2 $\leq$ XLOGP $\leq$ 5
TPSA $\leq$ 150 Yes
Num. rings $\leq$ 7
Num. carbon $>$ 4
Num. heteroatoms $>$ 1
Num. rotatable bonds $\leq$
15
H-bond acc. $\leq$ 10
H-bond don. $\leq$ 5

Bioavailability Score ⓘ


Abbott Bioavailability
Score: Probability of F 0.56
> 10% in rat
implemented from
Martin YC. 2005 J.
Med. Chem.

Medicinal Chemistry

PAINS ⓘ

1 alert: quinone_A ⓘ

Pan Assay Interference
Structures:
implemented from
Baell JB. & Holloway
GA. 2010 J. Med.
Chem.



Brenk ⓘ

Structural Alert:
implemented from 0 alert
Brenk R. et al. 2008
ChemMedChem

Leadlikeness ⓘ

Leadlikeness:
implemented from
Teague SJ. 1999 Angew.
Chem. Int. Ed. Yes
 $250 \leq MW \leq 350$
 $XLOGP \leq 3.5$
Num. rotatable bonds $\leq$
7

Synthetic accessibility ⓘ

Synthetic accessibility
score: from 1 (very
easy) to 10 (very
difficult)
based on 1024
fragmental contributions 2.95
(FP2) modulated by size
and complexity penalties,
trained on 12'782'590
molecules and tested on
40 external molecules
($r^2 = 0.94$)

In order to avoid security-related warning messages when switching to secured connection, you may want either to:

- confirm the exception on the next page, or
- import our [CA key](#) in your web browser

Click [here](#) to proceed.

[Click2Drug](#)

[SwissDock](#)

[SwissParam](#)

[SwissSidechain](#)

[SwissBioisostere](#)

[SwissTargetPrediction](#)

[SwissADME](#)

[SwissSimilarity](#)

[About us](#)

[Swiss Institute of Bioinformatics](#)

SwissADME

- [Disclaimer](#)
- [Help](#)
- [FAQ](#)
- [Home](#)


This website allows you to compute physicochemical descriptors as well as to predict ADME parameters, pharmacokinetic properties, druglike nature and medicinal chemistry friendliness of one or multiple small molecules to support drug discovery.

The main article describing the web service and its underlying methodologies is [SwissADME: a free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules](#). *Sci. Rep.* (2017) 7:42717.

For details about development and validation of iLOG, please refer to this article: [iLOGP: a simple, robust, and efficient description of *n*-octanol/water partition coefficient for drug design using the GB/SA approach](#). *J. Chem. Inf. Model.* (2014) 54(12):3284-3301.

For details about development and validation of the BOILED-Egg, please refer to this article: [A BOILED-Egg to predict gastrointestinal absorption and brain penetration of small molecules](#). *ChemMedChem* (2016) 11(11):1117-1121.

Developed and maintained by the [Molecular Modeling Group](#) of the SIB | Swiss Institute of Bioinformatics.



Enter a list of SMILES here:

O=C4C=C(Oc1ccc(cc1)c2ccccc2)C(=O)c3ccccc34

Fill with an example

Clear



Run!

Show BOILED-Egg

Retrieve data:


 Powered by
ChemAxon

Molecule 1


 SMILES
S O=C1C(=CC(=O)c2c1cccc2)Oc1ccc(cc1)c1ccccc1

Physicochemical Properties

Formula	C22H14O3
Molecular weight	326.34 g/mol
Num. heavy atoms	25
Num. arom. heavy atoms	18
Fraction Csp3	0.00
Num. rotatable bonds	3
Num. H-bond acceptors	3
Num. H-bond donors	0
Molar Refractivity	95.69
TPSA	

Water Solubility

Log S (ESOL)

ESOL: Topological method implemented from
Delaney JS. 2004 J. Chem. Inf. Model.

-5.05

 Solubility
Class

2.89e-03 mg/ml ; 8.87e-06 mol/l

Solubility class: Log S scale

Insoluble < -10 < Poorly Moderately soluble
< -6 < Moderately < -4
< Soluble < -2 Very < 0
< Highly

Log S (Ali)

Ali: Topological method implemented from
Ali J. et al. 2012 J. Chem. Inf. Model.

-5.16

 Solubility
Class

2.24e-03 mg/ml ; 6.87e-06 mol/l

Solubility class: Log S

Topological Polar Surface Area: Calculated from Ertl P. et al. 2000 J. Med. Chem.	43.37 Å ²	<u>scale</u> Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Moderately soluble
Log <i>P</i>_{o/w} (iLOGP) ⓘ		Log <i>S</i> (SILICOS-IT) ⓘ	
iLOGP: in-house physics-based method implemented from Daina A et al. 2014 J. Chem. Inf. Model.	2.89	SILICOS-IT: Fragmental method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos-it.com	-7.79
Log <i>P</i>_{o/w} (XLOGP3) ⓘ		Solubility	5.35e-06 mg/ml ; 1.64e-08 mol/l
XLOGP3: Atomistic and knowledge-based method calculated by XLOGP program, version 3.2.2, courtesy of CCBG, Shanghai Institute of Organic Chemistry	4.53	Class ⓘ	
Log <i>P</i>_{o/w} (WLOGP) ⓘ		Solubility class: Log <i>S</i> scale Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Poorly soluble
WLOGP: Atomistic method implemented from Wildman SA and Crippen GM. 1999 J. Chem. Inf. Model.	4.70		Pharmacokinetics
Log <i>P</i>_{o/w} (MLOGP) ⓘ		GI absorption ⓘ	
MLOGP: Topological method implemented from Moriguchi I. et al. 1992 Chem. Pharm. Bull. Moriguchi I. et al. 1994 Chem. Pharm. Bull. Lipinski PA. et al. 2001 Adv. Drug. Deliv. Rev.	2.85	Gastrointestinal absorption: according to the white of the BOILED-Egg	High
Log <i>P</i>_{o/w} (SILICOS-IT) ⓘ		BBB permeant ⓘ	
SILICOS-IT: Hybrid fragmental/topological method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos-it.com	4.92	BBB permeation: according to the yolk of the BOILED-Egg	Yes
Consensus Log <i>P</i>_{o/w} ⓘ		P-gp substrate ⓘ	
Consensus Log <i>P</i>_{o/w}:		P-glycoprotein substrate: SVM model built on 1033 molecules (training set) and tested on 415 molecules (test set) 10-fold CV: ACC=0.72 / AUC=0.77 External: ACC=0.88 / AUC=0.94	No
		CYP1A2 inhibitor ⓘ	
		Cytochrome P450 1A2 inhibitor: SVM model built on 9145 molecules (training set) and tested on 3000 molecules (test set)	Yes

Average of all five predictions

3.98

10-fold CV: ACC=0.83 / AUC=0.90
External: ACC=0.84 / AUC=0.91

CYP2C19 inhibitor ⓘ

Cytochrome P450 2C19 inhibitor: SVM model built on 9272 molecules (training set) and tested on 3000 molecules (test set) Yes
10-fold CV: ACC=0.80 / AUC=0.86
External: ACC=0.80 / AUC=0.87

CYP2C9 inhibitor ⓘ

Cytochrome P450 2C9 inhibitor: SVM model built on 5940 molecules (training set) and tested on 2075 molecules (test set) Yes
10-fold CV: ACC=0.78 / AUC=0.85
External: ACC=0.71 / AUC=0.81

CYP2D6 inhibitor ⓘ

Cytochrome P450 2D6 inhibitor: SVM model built on 3664 molecules (training set) and tested on 1068 molecules (test set) No
10-fold CV: ACC=0.79 / AUC=0.85
External: ACC=0.81 / AUC=0.87

CYP3A4 inhibitor ⓘ

Cytochrome P450 3A4 inhibitor: SVM model built on 7518 molecules (training set) and tested on 2579 molecules (test set) Yes
10-fold CV: ACC=0.77 / AUC=0.85
External: ACC=0.78 / AUC=0.86

Log K_p (skin permeation) ⓘ

Skin permeation: QSPR model implemented from -5.07 cm/s

Potts RO and Guy RH.
1992 Pharm. Res.

Druglikeness

Lipinski ⓘ

Lipinski (Pfizer) filter:
implemented from

Lipinski CA. et al. 2001

Adv. Drug Deliv. Rev.

Yes; 0 violation

MW $\leq$ 500

MLOGP $\leq$ 4.15

N or O $\leq$ 10

NH or OH $\leq$ 5

Ghose ⓘ

Ghose filter:
implemented from

Ghose AK. et al. 1999 J.

Comb. Chem.

Yes

160 $\leq$ MW $\leq$ 480

-0.4 $\leq$ WLOGP $\leq$ 5.6

40 $\leq$ MR $\leq$ 130

20 $\leq$ atoms $\leq$ 70

Veber ⓘ

Veber (GSK) filter:
implemented from

Veber DF. et al. 2002 J.

Yes

Med. Chem.

Rotatable bonds $\leq$ 10

TPSA $\leq$ 140

Egan ⓘ

Egan (Pharmacia)
filter: implemented
from

Egan WJ. et al. 2000 J.

Yes

Med. Chem.

WLOGP $\leq$ 5.88

TPSA $\leq$ 131.6

Muegge ⓘ

Muegge (Bayer) filter:
implemented from

Muegge I. et al. 2001 J.

Med. Chem.

200 $\leq$ MW $\leq$ 600

-2 $\leq$ XLOGP $\leq$ 5

TPSA $\leq$ 150

Yes

Num. rings $\leq$ 7

Num. carbon $>$ 4

Num. heteroatoms $>$ 1

Num. rotatable bonds $\leq$

15

H-bond acc. $\leq$ 10

H-bond don. $\leq$ 5

Bioavailability Score ⓘ


Abbott Bioavailability
Score: Probability of F 0.56
> 10% in rat
implemented from
Martin YC. 2005 J.
Med. Chem.

Medicinal Chemistry

PAINS ⓘ

1 alert: quinone_A ⓘ

Pan Assay Interference
Structures:
implemented from
Baell JB. & Holloway
GA. 2010 J. Med.
Chem.



Brenk ⓘ

Structural Alert:
implemented from 0 alert
Brenk R. et al. 2008
ChemMedChem

Leadlikeness ⓘ

Leadlikeness:
implemented from
Teague SJ. 1999 Angew.
Chem. Int. Ed. No; 1 violation: XLOGP3>3.5
 $250 \leq MW \leq 350$
 $XLOGP \leq 3.5$
Num. rotatable bonds $\leq$
7

Synthetic accessibility ⓘ

Synthetic accessibility
score: from 1 (very
easy) to 10 (very
difficult)
based on 1024
fragmental contributions 3.14
(FP2) modulated by size
and complexity penalties,
trained on 12'782'590
molecules and tested on
40 external molecules
($r^2 = 0.94$)

In order to avoid security-related warning messages when switching to secured connection, you may want either to:

- confirm the exception on the next page, or
- import our [CA key](#) in your web browser

Click [here](#) to proceed.

[Click2Drug](#)

[SwissDock](#)

[SwissParam](#)

[SwissSidechain](#)

[SwissBioisostere](#)

[SwissTargetPrediction](#)

[SwissADME](#)

[SwissSimilarity](#)

[About us](#)

[Swiss Institute of Bioinformatics](#)

SwissADME

- [Disclaimer](#)
- [Help](#)
- [FAQ](#)
- [Home](#)


This website allows you to compute physicochemical descriptors as well as to predict ADME parameters, pharmacokinetic properties, druglike nature and medicinal chemistry friendliness of one or multiple small molecules to support drug discovery.

The main article describing the web service and its underlying methodologies is [SwissADME: a free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules. *Sci. Rep.* \(2017\) 7:42717.](#)

For details about development and validation of iLOG, please refer to this article: [iLOGP: a simple, robust, and efficient description of *n*-octanol/water partition coefficient for drug design using the GB/SA approach. *J. Chem. Inf. Model.* \(2014\) 54\(12\):3284-3301.](#)

For details about development and validation of the BOILED-Egg, please refer to this article: [A BOILED-Egg to predict gastrointestinal absorption and brain penetration of small molecules. *ChemMedChem* \(2016\) 11\(11\):1117-1121.](#)

Developed and maintained by the [Molecular Modeling Group](#) of the SIB | Swiss Institute of Bioinformatics.



Enter a list of SMILES here:

```
CC(C)c3ccc(C)cc3OC2=CC(=O)c1ccccc1C2=O
```

Fill with an example

Clear



Run!

Show BOILED-Egg

Retrieve data:


 Powered by
 ChemAxon

Molecule 1



SMILES
S Cc1ccc(c(c1)OC1=CC(=O)c2c(C1=O)cccc2)C(C)C

Physicochemical Properties

Formula	C20H18O3
Molecular weight	306.36 g/mol
Num. heavy atoms	23
Num. arom. heavy atoms	12
Fraction Csp3	0.20
Num. rotatable bonds	3
Num. H-bond acceptors	3
Num. H-bond donors	0
Molar Refractivity	89.80
TPSA	

Water Solubility

Log S (ESOL)

ESOL: Topological method implemented from Delaney JS. 2004 J. Chem. Inf. Model.

-4.88

Solubility
Class

4.08e-03 mg/ml ; 1.33e-05 mol/l

Solubility class: Log S scale

Insoluble < -10 < Poorly Moderately soluble
< -6 < Moderately < -4
< Soluble < -2 Very < 0
< Highly

Log S (Ali)

Ali: Topological method implemented from Ali J. et al. 2012 J. Chem. Inf. Model.

-5.32

Solubility
Class

1.47e-03 mg/ml ; 4.80e-06 mol/l

Solubility class: Log S

Topological Polar Surface Area: Calculated from Ertl P. et al. 2000 J. Med. Chem.	43.37 Å ²	<u>scale</u> Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Moderately soluble
Log $P_{o/w}$ (iLOGP) ⓘ	Lipophilicity	Log S (SILICOS-IT) ⓘ	
iLOGP: in-house physics-based method implemented from Daina A et al. 2014 J. Chem. Inf. Model.	3.06	SILICOS-IT: Fragmental method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos-it.com	-6.49
Log $P_{o/w}$ (XLOGP3) ⓘ		Solubility Class ⓘ	9.94e-05 mg/ml ; 3.24e-07 mol/l
XLOGP3: Atomistic and knowledge-based method calculated by XLOGP program, version 3.2.2, courtesy of CCBG, Shanghai Institute of Organic Chemistry	4.68	Solubility class: Log S <u>scale</u> Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Poorly soluble
Log $P_{o/w}$ (WLOGP) ⓘ			Pharmacokinetics
WLOGP: Atomistic method implemented from Wildman SA and Crippen GM. 1999 J. Chem. Inf. Model.	4.46	GI absorption ⓘ	
Log $P_{o/w}$ (MLOGP) ⓘ		Gastrointestinal absorption: according to the white of the BOILED-Egg	High
MLOGP: Topological method implemented from Moriguchi I. et al. 1992 Chem. Pharm. Bull. Moriguchi I. et al. 1994 Chem. Pharm. Bull. Lipinski PA. et al. 2001 Adv. Drug. Deliv. Rev.	2.62	BBB permeant ⓘ	
Log $P_{o/w}$ (SILICOS-IT) ⓘ		BBB permeation: according to the yolk of the BOILED-Egg	Yes
SILICOS-IT: Hybrid fragmental/topological method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos-it.com	4.95	P-gp substrate ⓘ	
Consensus Log $P_{o/w}$ ⓘ		P-glycoprotein substrate: SVM model built on 1033 molecules (training set) and tested on 415 molecules (test set) 10-fold CV: ACC=0.72 / AUC=0.77 External: ACC=0.88 / AUC=0.94	No
Consensus Log $P_{o/w}$:		CYP1A2 inhibitor ⓘ	
		Cytochrome P450 1A2 inhibitor: SVM model built on 9145 molecules (training set) and tested on 3000 molecules (test set)	Yes

Average of all five predictions

3.95

10-fold CV: ACC=0.83 / AUC=0.90
External: ACC=0.84 / AUC=0.91

CYP2C19 inhibitor ⓘ

Cytochrome P450 2C19 inhibitor: SVM model built on 9272 molecules (training set) and tested on 3000 molecules (test set) Yes
10-fold CV: ACC=0.80 / AUC=0.86
External: ACC=0.80 / AUC=0.87

CYP2C9 inhibitor ⓘ

Cytochrome P450 2C9 inhibitor: SVM model built on 5940 molecules (training set) and tested on 2075 molecules (test set) Yes
10-fold CV: ACC=0.78 / AUC=0.85
External: ACC=0.71 / AUC=0.81

CYP2D6 inhibitor ⓘ

Cytochrome P450 2D6 inhibitor: SVM model built on 3664 molecules (training set) and tested on 1068 molecules (test set) No
10-fold CV: ACC=0.79 / AUC=0.85
External: ACC=0.81 / AUC=0.87

CYP3A4 inhibitor ⓘ

Cytochrome P450 3A4 inhibitor: SVM model built on 7518 molecules (training set) and tested on 2579 molecules (test set) Yes
10-fold CV: ACC=0.77 / AUC=0.85
External: ACC=0.78 / AUC=0.86

Log K_p (skin permeation) ⓘ

Skin permeation: QSPR model implemented from -4.85 cm/s

Potts RO and Guy RH.
1992 Pharm. Res.

Druglikeness

Lipinski

Lipinski (Pfizer) filter:
implemented from
Lipinski CA. et al. 2001
Adv. Drug Deliv. Rev. Yes; 0 violation
MW $\leq$ 500
MLOGP $\leq$ 4.15
N or O $\leq$ 10
NH or OH $\leq$ 5

Ghose

Ghose filter:
implemented from
Ghose AK. et al. 1999 J.
Comb. Chem. Yes
160 $\leq$ MW $\leq$ 480
-0.4 $\leq$ WLOGP $\leq$ 5.6
40 $\leq$ MR $\leq$ 130
20 $\leq$ atoms $\leq$ 70

Veber

Veber (GSK) filter:
implemented from
Veber DF. et al. 2002 J. Yes
Med. Chem.
Rotatable bonds $\leq$ 10
TPSA $\leq$ 140

Egan

Egan (Pharmacia)
filter: implemented
from
Egan WJ. et al. 2000 J. Yes
Med. Chem.
WLOGP $\leq$ 5.88
TPSA $\leq$ 131.6

Muegge

Muegge (Bayer) filter:
implemented from
Muegge I. et al. 2001 J.
Med. Chem.
200 $\leq$ MW $\leq$ 600
-2 $\leq$ XLOGP $\leq$ 5
TPSA $\leq$ 150 Yes
Num. rings $\leq$ 7
Num. carbon $>$ 4
Num. heteroatoms $>$ 1
Num. rotatable bonds $\leq$
15
H-bond acc. $\leq$ 10
H-bond don. $\leq$ 5

Bioavailability Score ⓘ


Abbott Bioavailability
Score: Probability of F 0.56
> 10% in rat
implemented from
Martin YC. 2005 J.
Med. Chem.

Medicinal Chemistry

PAINS ⓘ

1 alert: quinone_A ⓘ

Pan Assay Interference
Structures:
implemented from
Baell JB. & Holloway
GA. 2010 J. Med.
Chem.



Brenk ⓘ

Structural Alert:
implemented from 0 alert
Brenk R. et al. 2008
ChemMedChem

Leadlikeness ⓘ

Leadlikeness:
implemented from
Teague SJ. 1999 Angew.
Chem. Int. Ed. No; 1 violation: XLOGP3>3.5
 $250 \leq MW \leq 350$
 $XLOGP \leq 3.5$
Num. rotatable bonds $\leq$
7

Synthetic accessibility ⓘ

Synthetic accessibility
score: from 1 (very
easy) to 10 (very
difficult)
based on 1024
fragmental contributions 3.31
(FP2) modulated by size
and complexity penalties,
trained on 12'782'590
molecules and tested on
40 external molecules
($r^2 = 0.94$)

In order to avoid security-related warning messages when switching to secured connection, you may want either to:

- confirm the exception on the next page, or
- import our [CA key](#) in your web browser

Click [here](#) to proceed.

[Click2Drug](#)

[SwissDock](#)

[SwissParam](#)

[SwissSidechain](#)

[SwissBioisostere](#)

[SwissTargetPrediction](#)

[SwissADME](#)

[SwissSimilarity](#)

[About us](#)

[Swiss Institute of Bioinformatics](#)

SwissADME

- [Disclaimer](#)
- [Help](#)
- [FAQ](#)
- [Home](#)


This website allows you to compute physicochemical descriptors as well as to predict ADME parameters, pharmacokinetic properties, druglike nature and medicinal chemistry friendliness of one or multiple small molecules to support drug discovery.

The main article describing the web service and its underlying methodologies is [SwissADME: a free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules. *Sci. Rep.* \(2017\) 7:42717.](#)

For details about development and validation of iLOG, please refer to this article: [iLOGP: a simple, robust, and efficient description of *n*-octanol/water partition coefficient for drug design using the GB/SA approach. *J. Chem. Inf. Model.* \(2014\) 54\(12\):3284-3301.](#)

For details about development and validation of the BOILED-Egg, please refer to this article: [A BOILED-Egg to predict gastrointestinal absorption and brain penetration of small molecules. *ChemMedChem* \(2016\) 11\(11\):1117-1121.](#)

Developed and maintained by the [Molecular Modeling Group](#) of the SIB | Swiss Institute of Bioinformatics.



Enter a list of SMILES here:

Clc4ccc(Oc3ccc(Cl)cc3OC2=CC(=O)c1ccccc1C2=O)c(Cl)c4

Fill with an example

Clear



Run!

Show BOILED-Egg

Retrieve data:


 Powered by
ChemAxon

Molecule 1



SMILE Clc1ccc(c(c1)OC1=CC(=O)c2c(C1=O)cccc2)Oc1ccc(cc1Cl)Cl

Physicochemical Properties

Formula C₂₂H₁₁Cl₃O₄
 Molecular weight 445.68 g/mol
 Num. heavy atoms 29
 Num. arom. heavy atoms 18
 Fraction Csp³ 0.00
 Num. rotatable bonds 4
 Num. H-bond acceptors 4
 Num. H-bond donors 0
 Molar Refractivity 111.80
 TPSA

Water Solubility

Log *S* (ESOL)

ESOL: Topological method implemented from Delaney JS. 2004 J. Chem. Inf. Model.

-6.96

Solubility Class

4.93e-05 mg/ml ; 1.11e-07 mol/l

Solubility class: Log *S* scale

Insoluble < -10 < Poorly Poorly soluble
< -6 < Moderately < -4
< Soluble < -2 Very < 0
< Highly

Log *S* (Ali)

Ali: Topological method implemented from Ali J. et al. 2012 J. Chem. Inf. Model.

-7.50

Solubility Class

1.39e-05 mg/ml ; 3.13e-08 mol/l

Solubility class: Log *S*

Topological Polar Surface Area: <u>Calculated from Ertl P. et al. 2000 J. Med. Chem.</u>	52.60 Å ²	<u>scale</u> Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Poorly soluble
Log <i>P</i>_{o/w} (iLOGP) ⓘ	Lipophilicity	Log <i>S</i> (SILICOS-IT) ⓘ	
iLOGP: in-house physics-based method implemented from Daina A et al. 2014 J. Chem. Inf. Model.	3.66	SILICOS-IT: <u>Fragmental method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos-it.com</u>	-9.28
Log <i>P</i>_{o/w} (XLOGP3) ⓘ		Solubility Class ⓘ	2.36e-07 mg/ml ; 5.28e-10 mol/l
XLOGP3: Atomistic and knowledge-based method calculated by XLOGP program, version 3.2.2, courtesy of CCBG, Shanghai Institute of Organic Chemistry	6.60	Solubility class: Log <i>S</i> scale Insoluble < -10 < Poorly < -6 < Moderately < -4 < Soluble < -2 Very < 0 < Highly	Poorly soluble
Log <i>P</i>_{o/w} (WLOGP) ⓘ			Pharmacokinetics
WLOGP: Atomistic method implemented from Wildman SA and Crippen GM. 1999 J. Chem. Inf. Model.	6.78	GI absorption ⓘ	
Log <i>P</i>_{o/w} (MLOGP) ⓘ		Gastrointestinal absorption: according to the white of the BOILED-Egg	High
MLOGP: Topological method implemented from Moriguchi I. et al. 1992 Chem. Pharm. Bull. Moriguchi I. et al. 1994 Chem. Pharm. Bull. Lipinski PA. et al. 2001 Adv. Drug. Deliv. Rev.	3.97	BBB permeant ⓘ	
Log <i>P</i>_{o/w} (SILICOS-IT) ⓘ		BBB permeation: according to the yolk of the BOILED-Egg	No
SILICOS-IT: Hybrid fragmental/topological method calculated by FILTER-IT program, version 1.0.2, courtesy of SILICOS-IT, http://www.silicos-it.com	6.34	P-gp substrate ⓘ	
Consensus Log <i>P</i>_{o/w} ⓘ		P-glycoprotein substrate: SVM model built on 1033 molecules (training set) and tested on 415 molecules (test set) 10-fold CV: ACC=0.72 / AUC=0.77 External: ACC=0.88 / AUC=0.94	No
Consensus Log <i>P</i>_{o/w}:		CYP1A2 inhibitor ⓘ	
		Cytochrome P450 1A2 inhibitor: SVM model built on 9145 molecules (training set) and tested on 3000 molecules (test set)	No

Average of all five predictions

5.47

10-fold CV: ACC=0.83 / AUC=0.90
External: ACC=0.84 / AUC=0.91

CYP2C19 inhibitor ⓘ

Cytochrome P450 2C19 inhibitor: SVM model built on 9272 molecules (training set) and tested on 3000 molecules (test set) Yes
10-fold CV: ACC=0.80 / AUC=0.86
External: ACC=0.80 / AUC=0.87

CYP2C9 inhibitor ⓘ

Cytochrome P450 2C9 inhibitor: SVM model built on 5940 molecules (training set) and tested on 2075 molecules (test set) Yes
10-fold CV: ACC=0.78 / AUC=0.85
External: ACC=0.71 / AUC=0.81

CYP2D6 inhibitor ⓘ

Cytochrome P450 2D6 inhibitor: SVM model built on 3664 molecules (training set) and tested on 1068 molecules (test set) No
10-fold CV: ACC=0.79 / AUC=0.85
External: ACC=0.81 / AUC=0.87

CYP3A4 inhibitor ⓘ

Cytochrome P450 3A4 inhibitor: SVM model built on 7518 molecules (training set) and tested on 2579 molecules (test set) No
10-fold CV: ACC=0.77 / AUC=0.85
External: ACC=0.78 / AUC=0.86

Log K_p (skin permeation) ⓘ

Skin permeation: QSPR model implemented from -4.33 cm/s

Potts RO and Guy RH.
1992 Pharm. Res.

Druglikeness

Lipinski

Lipinski (Pfizer) filter:
implemented from

Lipinski CA. et al. 2001

Adv. Drug Deliv. Rev.

Yes; 0 violation

MW $\leq$ 500

MLOGP $\leq$ 4.15

N or O $\leq$ 10

NH or OH $\leq$ 5

Ghose

Ghose filter:

implemented from

Ghose AK. et al. 1999 J.

Comb. Chem.

No; 1 violation: WLOGP>5.6

160 $\leq$ MW $\leq$ 480

-0.4 $\leq$ WLOGP $\leq$ 5.6

40 $\leq$ MR $\leq$ 130

20 $\leq$ atoms $\leq$ 70

Veber

Veber (GSK) filter:

implemented from

Veber DF. et al. 2002 J.

Yes

Med. Chem.

Rotatable bonds $\leq$ 10

TPSA $\leq$ 140

Egan

Egan (Pharmacia)

filter: implemented

from

Egan WJ. et al. 2000 J.

No; 1 violation: WLOGP>5.88

Med. Chem.

WLOGP $\leq$ 5.88

TPSA $\leq$ 131.6

Muegge

Muegge (Bayer) filter:

implemented from

Muegge I. et al. 2001 J.

Med. Chem.

200 $\leq$ MW $\leq$ 600

-2 $\leq$ XLOGP $\leq$ 5

TPSA $\leq$ 150

No; 1 violation: XLOGP3>5

Num. rings $\leq$ 7

Num. carbon $>$ 4

Num. heteroatoms $>$ 1

Num. rotatable bonds $\leq$

15

H-bond acc. $\leq$ 10

H-bond don. $\leq$ 5

Bioavailability Score ⓘ


Abbott Bioavailability
Score: Probability of F 0.56
> 10% in rat
implemented from
Martin YC. 2005 J.
Med. Chem.

Medicinal Chemistry

PAINS ⓘ

1 alert: quinone_A ⓘ

Pan Assay Interference
Structures:
implemented from
Baell JB. & Holloway
GA. 2010 J. Med.
Chem.



Brenk ⓘ

Structural Alert:
implemented from 0 alert
Brenk R. et al. 2008
ChemMedChem

Leadlikeness ⓘ

Leadlikeness:
implemented from
Teague SJ. 1999 Angew. No; 2 violations: MW>350,
Chem. Int. Ed. XLOGP3>3.5
 $250 \leq MW \leq 350$
 $XLOGP \leq 3.5$
Num. rotatable bonds $\leq$
7

Synthetic accessibility ⓘ

Synthetic accessibility
score: from 1 (very
easy) to 10 (very
difficult)
based on 1024
fragmental contributions 3.34
(FP2) modulated by size
and complexity penalties,
trained on 12'782'590
molecules and tested on
40 external molecules
($r^2 = 0.94$)