

High-throughput discovery of hydrogen evolution electrocatalysts in the complex solid solution system Co-Cr-Fe-Mo-Nit

Simon Schumacher,^{a‡} Sabrina Baha,^{b‡} Alan Savan,^b Corina Andronescu,^{a*} and Alfred Ludwig^{bc*}

^a Chemical Technology III, Faculty of Chemistry and CENIDE Center for Nanointegration, University Duisburg-Essen, Carl-Benz Straße 199, 47057 Duisburg, Germany, corina.andronescu@uni-due.de

^b Materials Discovery and Interfaces, Institute for Materials, Ruhr University Bochum, Universitätsstraße 150, D-44801 Bochum, Germany, alfred.ludwig@ruhr-uni-bochum.de

^c Center for Interface-Dominated High Performance Materials, Ruhr University Bochum, Universitätsstraße 150, D-44801 Bochum, Germany.

† Electronic Supplementary Information (ESI) available. See DOI: 10.1039/x0xx00000x

‡ Equal contributions.




Fig. S1: Comparison of the colormaps of the current at -250 mV vs RHE, the Tafel slope in log(*i*)-range 2.5 – 3.5 and the R^2 -values of the linear fitting of the corresponding Tafel slopes




Fig. S2: Tafel-plots from the LSVs with the highest (red) and the lowest current (blue) at -250 mV vs RHE and the center MA (green) from each ML